



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

June 12, 2020

Eric D. Smith  
Director, Regulatory Affairs  
PBI/Gordon Corporation  
1217 W. 12<sup>th</sup> Street  
Kansas City, MO 64101

Subject: Label Amendment – Marketing Statements, Resistance Management Language  
Product Name: EH-1369 Professional  
EPA Registration Number: 2217-907  
Application Date: March 26, 2018  
Decision Number: 540196

Dear Mr. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 2217-907  
Decision No. 540196

with FIFRA section 6. If you have any questions, please contact Kable Bo Davis by phone at 703-306-0415, or via email at [davis.kable@epa.gov](mailto:davis.kable@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read 'MO' followed by a flourish, and the word 'for' written to the right.

Mindy Ondish  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

|                                      |       |   |           |
|--------------------------------------|-------|---|-----------|
| 2,4-D<br>Mecoprop-p<br>Dichlorprop-p | GROUP | 4 | HERBICIDE |
|--------------------------------------|-------|---|-----------|

# EH-1369 Professional

## EPA Reg. No. 2217-907

**ACTIVE INGREDIENTS:**


|   |                 |
|---|-----------------|
| 2,4-D, dimethylamine salt .....         | 0.723%          |
| Mecoprop-p, dimethylamine salt .....    | 0.148%          |
| Dichlorprop-P, dimethylamine salt ..... | 0.088%          |
| <b>OTHER INGREDIENTS:</b> .....         | <u>99.041%</u>  |
| <b>TOTAL</b> .....                      | <b>100.000%</b> |

**THIS PRODUCT CONTAINS:**

- 12.02 lb. 2,4-dichlorophenoxyacetic acid equivalent per ton, or 0.601%.
- 2.44 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton, or 0.122%
- 1.48 lb. (+)-(R)-2-(2,4-dichlorophenoxy) propionic acid equivalent per ton, or 0.074%.

Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/Gordon Corporation.



**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

See attached booklet [inside pages] for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements and Directions for Use, including First Aid, and Storage and Disposal.

**Guaranteed Analysis:** See appendix for example.

**Net Contents:** \_\_\_\_\_

**EPA Est. No.** \_\_\_\_\_

**Company Name and Address:**

PBI/Gordon Corporation  
P.O. Box 860350  
Shawnee, KS 66286



# **PRECAUTIONARY STATEMENTS**

## **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes or clothing.

## **Personal Protective Equipment (PPE)**

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt
- long pants and
- shoes plus socks.

## **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **User Safety Recommendations**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **First Aid**

|  |   |
|--|---|
| <b>If in eyes:</b>   | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If on skin or on clothing:</b>  | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If swallowed:</b>   | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information. |   |

## **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **[Environmental Hazards (alternate per PRN 2008-1)**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **USE RESTRICTIONS:**

- Do not enter or allow people (or pets) to enter the treated area until dusts have settled.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not over water the treated area to the point of runoff or apply when raining or when rain is expected that day.

## **Weed Resistance Management**

For resistance management, this product contains Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 4. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 4 within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different

group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.

- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 800-884-3179.

#### **WHERE TO USE:**

EH-1369 ORP is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p and dichlorprop-p. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass.

EH-1369 ORP is intended to be applied by noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

#### **WHEN TO APPLY:**

EH-1369 ORP may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

If over-seeding, wait 4 weeks after grass seeds sprout before applying this product.

EH-1369 ORP may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

#### **HOW MUCH TO APPLY:**

**For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass and zoysiagrass:** Apply 140 to 174 pounds of EH-1369 ORP per acre of established turfgrass. The equivalent application rates are 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass.

**For St. Augustinegrass, centipedegrass, hybrid bermudagrass, buffalograss, and colonial bentgrass:** Apply 110 to 140 pounds of EH-1369 ORP per acre of established turfgrass. The equivalent application rates are 2.5 to 3.2 pounds of product per 1,000 square feet of turfgrass.

Cultivars vary in tolerance to this product. Do not apply this product to 'Floritam' St. Augustinegrass.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

A sequential or follow-up application may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be 30 days.

**LIMITATIONS FOR USE ON TURFGRASS:**

- The maximum application rate is 4.0 pounds of product/1,000 sq.ft. per application (1.05 lb 2,4-D ae, 0.13 lb 2,4-DP-p ae and 0.21 lb MCP-P-p ae /acre per application).
- The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications.
- The maximum seasonal rate is 8.0 pounds of product/1,000 sq.ft. (2.10 lb 2,4-D ae, 0.26 lb 2,4-DP-p ae and 0.42 lb MCP-P-p ae/acre per year).
- Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers. Do not use this product on 'Floritam' St. Augustinegrass.
- Do not apply this product to newly seeded grasses until well-established or after the third mowing.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

**APPLICATION SUGGESTIONS:**

1. Spring and fall applications are recommended for the best control. However, EH-1369 ORP can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

**CALIBRATION INSTRUCTIONS:**

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

To provide proper distribution, calibrate spreader before application.

- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.
- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 3.8 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 260 square feet and adjust to apply at the rate of 1 pound of product per 260 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]



| Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. | Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. |
|----------------|-------------------------------------|--|----------------|-------------------------------------|--|
| 15             | 2.5                                 | 6,000                                  | 28             | 2.5                                 | 11,200                                 |
|                | 3.0                                 | 5,000                                  |                | 3.0                                 | 9,300                                  |
|                | 3.2                                 | 4,700                                  |                | 3.2                                 | 8,800                                  |
|                | 3.4                                 | 4,400                                  |                | 3.4                                 | 8,200                                  |
|                | 3.6                                 | 4,200                                  |                | 3.6                                 | 7,800                                  |
| 16             | 4.0                                 | 3,750                                  | 36             | 4.0                                 | 7,000                                  |
|                | 2.5                                 | 6,400                                  |                | 2.5                                 | 14,400                                 |
|                | 3.0                                 | 5,300                                  |                | 3.0                                 | 12,000                                 |
|                | 3.2                                 | 5,000                                  |                | 3.2                                 | 11,300                                 |
|                | 3.4                                 | 4,700                                  |                | 3.4                                 | 10,500                                 |
| 17             | 3.6                                 | 4,400                                  | 37             | 3.6                                 | 10,000                                 |
|                | 4.0                                 | 4,000                                  |                | 4.0                                 | 9,000                                  |
|                | 2.5                                 | 6,800                                  |                | 2.5                                 | 14,800                                 |
|                | 3.0                                 | 5,700                                  |                | 3.0                                 | 12,300                                 |
|                | 3.2                                 | 5,300                                  |                | 3.2                                 | 11,600                                 |
| 18             | 3.4                                 | 5,000                                  | 39             | 3.4                                 | 10,900                                 |
|                | 3.6                                 | 4,700                                  |                | 3.6                                 | 10,300                                 |
|                | 4.0                                 | 4,250                                  |                | 4.0                                 | 9,250                                  |
|                | 2.5                                 | 7,200                                  |                | 2.5                                 | 15,600                                 |
|                | 3.0                                 | 6,000                                  |                | 3.0                                 | 13,000                                 |
| 20             | 3.2                                 | 5,600                                  | 40             | 3.2                                 | 12,200                                 |
|                | 3.4                                 | 5,300                                  |                | 3.4                                 | 11,500                                 |
|                | 3.6                                 | 5,000                                  |                | 3.6                                 | 10,800                                 |
|                | 4.0                                 | 4,500                                  |                | 4.0                                 | 9,750                                  |
|                | 2.5                                 | 8,000                                  |                | 2.5                                 | 16,000                                 |
| 24             | 3.0                                 | 6,700                                  | 42             | 3.0                                 | 13,300                                 |
|                | 3.2                                 | 6,400                                  |                | 3.2                                 | 12,500                                 |
|                | 3.4                                 | 5,800                                  |                | 3.4                                 | 11,800                                 |
|                | 3.6                                 | 5,700                                  |                | 3.6                                 | 11,100                                 |
|                | 4.0                                 | 5,000                                  |                | 4.0                                 | 10,000                                 |
| 25             | 2.5                                 | 9,600                                  | 45             | 2.5                                 | 16,800                                 |
|                | 3.0                                 | 8,000                                  |                | 3.0                                 | 14,000                                 |
|                | 3.2                                 | 7,500                                  |                | 3.2                                 | 13,100                                 |
|                | 3.4                                 | 7,100                                  |                | 3.4                                 | 12,400                                 |
|                | 3.6                                 | 6,700                                  |                | 3.6                                 | 11,700                                 |
| 26             | 4.0                                 | 6,000                                  | 50             | 4.0                                 | 10,500                                 |
|                | 2.5                                 | 10,000                                 |                | 2.5                                 | 18,000                                 |
|                | 3.0                                 | 8,300                                  |                | 3.0                                 | 15,000                                 |
|                | 3.2                                 | 7,800                                  |                | 3.2                                 | 14,100                                 |
|                | 3.4                                 | 7,400                                  |                | 3.4                                 | 13,200                                 |
| 26             | 3.6                                 | 6,900                                  | 50             | 3.6                                 | 12,500                                 |
|                | 4.0                                 | 6,250                                  |                | 4.0                                 | 11,300                                 |
|                | 2.5                                 | 8,000                                  |                | 2.5                                 | 20,000                                 |
|                | 3.0                                 | 8,700                                  |                | 3.0                                 | 16,700                                 |
|                | 3.2                                 | 8,100                                  |                | 3.2                                 | 15,600                                 |
| 26             | 3.4                                 | 7,600                                  | 50             | 3.4                                 | 14,700                                 |
|                | 3.6                                 | 7,200                                  |                | 3.6                                 | 13,900                                 |
|                | 4.0                                 | 6,500                                  |                | 4.0                                 | 12,500                                 |

**[Optional Table**

| <b>Suggested Spreader Settings (Calibration Gauge Setting Number)</b><br>(Recommended rates in lb/1,000 sq.ft.) |                |                   |                 |                                   |
|---|----------------|-------------------|-----------------|-----------------------------------|
| <b>Scotts® Spreader Selection</b>   | <b>3.2 Low</b> | <b>3.6 Medium</b> | <b>4.0 High</b> | <b>Width of Coverage (in ft.)</b> |
| Scotts Landscaper Pro(TM) Spreader  | L              | M                 | N               | 13                                |



|   |       |       |      |    |
|---|-------|-------|------|----|
| Scotts Landscaper Pro(TM) Spreader Stainless  | L     | M     | N    | 13 |
| Scotts Deluxe EdgeGuard® Spreader   | 5     | 5.5   | 6.25 | 5  |
| AccuPro(TM) 2000 Rotary (cone 4)  | L     | M     | N    | 13 |
| SR-2000(TM) Rotary (cone 4)   | L     | M     | N    | 13 |
|   |       |       |      |    |
| Lbs. of N/1000 sq.ft.   | 0.7   | 0.896 | 1.12 |    |
| Lbs. of P2O5/1000 sq.ft.  | 0.075 | 0.096 | 0.12 |    |
| AccuPro(TM) and SR-2000(TM) are trademarks of The Andersons Agriservices, Inc.<br>To prevent staining, sweep or blow stray particles from concrete. |       |       |      |    |

**DISTRIBUTION AND SPREADER PATTERNS:**

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

**OR**

[Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply with a drop spreader or by hand. Application by drop spreader may damage the coated portion of this product.]

**SINGLE PASS DISTRIBUTION:**

For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.

1. For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage.
2. First, spread the borders of the lawn. Then apply this product to the center with straight line passes
3. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

**OR**

**[SINGLE PASS DISTRIBUTION:**

For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.

1. For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage.
2. Apply to the outer edges of the lawn first. Then apply this product to the center with straight line passes
3. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.]

Controls (Number) weeds. (Listing of weeds may include the following.)

| <b>BROADLEAF WEEDS</b>    |                 |                     |                        |
|---------------------------|-----------------|---------------------|------------------------|
| Annual yellow sweetclover | Cockle          | Musk thistle        | Spiny cocklebur        |
| Aster                     | Cocklebur       | Mustard             | Spiny sowthistle       |
| Austrian fieldcress       | Common mullein  | Narrowleaf plantain | Spotted catsear        |
| Bedstraw                  | Creeping jenny  | Narrowleaf vetch    | Spotted spurge         |
| Beggarticks               | Cudweed         | Nettle              | Spurweed               |
| Betony, Florida           | Curly dock      | Orange hawkweed     | Stinging nettle        |
| Bindweed, field           | Daisy, English  | Oriental cocklebur  | Strawberry, India mock |
| Bird vetch                | Daisy fleabane  | Oxalis              | Tall nettle            |
| Bitter wintercress        | Daisy, oxeye    | Parsley-piert       | Tall vervain           |
| Bittercress, hairy        | Dandelion       | Parsnip             | Tansy ragwort          |
| Bitterweed                | Dichondra       | Pearlwort           | Tansy mustard          |
| Black-eyed Susan          | Dogbane         | Pennycress          | Tanweed                |
| Black medic               | Dogfennel       | Pennywort           | Thistle                |
| Black mustard             | Dollarweed      | Pepperweed          | Trailing crownvetch    |
| Blackseed plantain        | Elderberry      | Pigweed             | Tumble mustard         |
| Blessed thistle           | False dandelion | Pineywoods bedstraw | Tumble pigweed         |

## BROADLEAF WEEDS

|                      |                    |                             |                         |
|----------------------|--------------------|-----------------------------|-------------------------|
| Bloodflower milkweed | Falseflax          | Plains coreopsis (tickseed) | Velvetleaf              |
| Blue lettuce         | False sunflower    | Plantain                    | Venice mallow           |
| Blue vervain         | Fiddleneck         | Poison ivy                  | Virginia buttonweed     |
| Bracted plantain     | Florida pusley     | Poison oak                  | Virginia creeper        |
| Brassbuttons         | Frenchweed         | Pokeweed                    | Virginia pepperweed     |
| Bristly oxtongue     | Galinsoga          | Poorjoe                     | Wavyleaf bullthistle    |
| Broadleaf dock       | Goathead           | Prairie sunflower           | Western clematis        |
| Broadleaf plantain   | Goldenrod          | Prickly lettuce             | Western salsify         |
| Broomweed            | Ground ivy         | Prickly sida                | White mustard           |
| Buckhorn             | Gumweed            | Prostrate knotweed          | Wild aster              |
| Buckhorn plantain    | Hairy fleabane     | Prostrate pigweed           | Wild buckwheat          |
| Bulbous buttercup    | Hawkweed           | Prostrate spurge            | Wild carrot             |
| Bull thistle         | Healall            | Prostrate vervain           | Wild four-o'clock       |
| Bullnettle           | Heartleaf drymary  | Puncturevine                | Wild garlic             |
| Burclover            | Heath aster        | Purslane, common            | Wild geranium           |
| Burdock              | Hedge bindweed     | Ragweed                     | Wild lettuce            |
| Burning nettle       | Hedge mustard      | Red sorrel                  | Wild marigold           |
| Burweed              | Hemp               | Redroot pigweed             | Wild mustard            |
| Buttercup            | Henbit             | Redstem filaree             | Wild onion              |
| Buttonweed           | Hoary cress        | Rough cinquefoil            | Wild parsnip            |
| Canada thistle       | Hoary plantain     | Rough fleabane              | Wild radish             |
| Carolina geranium    | Hoary vervain      | Russian pigweed             | Wild rape               |
| Carpetweed           | Horsenettle        | Russian thistle             | Wild strawberry         |
| Catchweed bedstraw   | Jimsonweed         | Scarlet pimpernel           | Wild sweet potato       |
| Catnip               | Knawel             | Scotch thistle              | Wild vetch              |
| Catsear              | Knotweed           | Sheep sorrel                | Woodsorrel              |
| Chickweed, common    | Kochia             | Shepherdspurse              | Woolly croton           |
| Chickweed, mouse ear | Lambsquarters      | Slender plantain            | Woolly morningglory     |
| Chicory              | Lespedeza          | Smallflower galinsoga       | Woolly plantain         |
| Cinquefoil           | Mallow             | Smooth dock                 | Wormseed                |
| Clover, crimson      | Matchweed          | Smooth pigweed              | Yarrow                  |
| Clover, hop          | Mexicanweed        | Sorrel                      | Yellow rocket           |
| Clover, red          | Milk vetch         | Sowthistle                  | Yellowflower pepperweed |
| Clover, strawberry   | Morningglory       | Spanishneedles              |                         |
| Clover, sweet        | Mouse ear hawkweed | Speedwell                   |                         |
| Clover, white        | Mugwort            | Spiny amaranth              |                         |

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **LIMITED WARRANTY AND DISCLAIMER**

**IMPORTANT:** Read this Limited Warranty and Disclaimer before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this Limited Warranty and Disclaimer in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the directions for use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.** The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

## **APPENDIX**

### **1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- (Fifteen, Sixteen, Seventeen, Eighteen, Twenty, or Fifty) Pounds Covers Up To X,000 Square Feet.
- Controls (desired weed species from attached listing may be inserted here e.g. dandelion, chickweed, knotweed, plantain, henbit, spurge) and over (200+) (more) broadleaf weeds.
- Helps build a stronger, deeper root system.
- Feeds your lawn with time-released nitrogen for extended greening.
- Encourages early Spring green-up.
- Timed-release fertilizer
- Use when broadleaf weeds are actively growing.
- Feeds for up to [3 months] [time period as designated by subregistrant].
- Quick greening & extended feeding for up to 8 weeks.
- Nitrogen (N) contains timed-release for growing a thick green lawn.

- Phosphorous (P) improves root growth & plant storage capacity.
- Potassium (K) improves plant hardiness & drought resistance.
- For a thick, green weed-free lawn.
- Greens fast & lasts.
- Formulated with Scotts® Poly-S® controlled release fertilizer, lasting up to 3 months.
- APPLY ANYTIME WHEN WEEDS ARE ACTIVELY GROWING.
- Scotts Landscaper Pro(TM) Turf Fertilizer with Post Emergent Weed Control controls over 200 Broadleaf weeds while providing proper nutrition for the turf.
- [N18] [NITROGEN] [For vigorous blade, leaf and stem growth and dark green color]
- [P0] [PHOSPHATE] [Contains zero Phosphorus]
- [K9][POTASH][Promotes strong roots and improves turf hardiness]
- Kills 200+ listed weeds, including dandelion, clover and other common listed weeds

**2. State mandated fertilizer labeling requirements:**

**[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “State Restrictions”]**

| <b>GUARANTEED ANALYSIS XX-X-X</b>   |     |
|---|-----|
| Total Nitrogen (N) .....  | XX% |
| x.x% ammoniacal nitrogen  |     |
| x.x% urea nitrogen*   |     |
| Available Phosphate (P2O5) .....  | X%  |
| Soluble Potash (K2O) .....  | X%  |
| Sulfur (S) .....  | X%  |
| x.x% free sulfur (S)  |     |
| Iron (Fe).....  | X%  |
| Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.  |     |
| *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.  |     |
| This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx |     |

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “[When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.]Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”

- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet [and herbicide to control listed weeds]. Please consult your local extension office for questions regarding a regular fertilizer program [suitable for your geographic region]."

## **DOCUMENT CONTROL INFORMATION**

**1. Unique Label Identifier:** 002217-00907.20200611.amend-proposed-clean

**2. Reason for Issue:** add additional ad claims, add missing lb/gal information to substatement of ingredient statement, reorganize label by moving referral statement to front panel and state mandated fertilizer statements to appendix, addition of weed resistance management