

2217-910

10/9/2009

Tarket
10/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

PBI / Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101

OCT 9 2009

Subject: EPA Reg. 2217-910 / EH-1458 Herbicide Dicamba RED Amendment

The labeling referred to above is in compliance with the Dicamba RED and is amended under the RED provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- 2) Please note: Final product reregistration can't be considered until after all active ingredients in this product are eligible for reregistration.
- 3) Delete "General" from "General Precautions".
- 4) Delete "recommended" from "30 day intervals are recommended" on page 6.
- 5) To the Warranty section add "to the extent consistent with applicable law" in front of "the manufacturer neither makes nor" and "in no case".

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supersedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2 of 10

ACCEPTED
with COMMENTS
In EPA Letter Dated:
OCT 9 2009

EH-1458 HERBICIDE

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EPA Reg. No. 2217-910

2217-910

For Residential Use Only

ACTIVE INGREDIENT:

2,4-D, 2-ethylhexyl ester	4.01%
Mecoprop-p acid	0.49%
Dicamba acid	0.27%
Carfentrazone-ethyl	0.16%
INERT INGREDIENTS:	95.07%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

1.40 lb 0.226 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 2.66%.
Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through the skin. Avoid contact with skin, eyes, or clothing. When using this product, wear long-sleeved shirt and long pants, socks, shoes, and gloves.

Personal Hygiene Statements

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

First Aid	
If on skin:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed: (optional for Toxicity Category IV)	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)]

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Physical and Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


GENERAL PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.



- [Optional Icon] Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans.
- Do not apply when windy
- To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn or garden area only.

PRODUCT FACTS:	
<ul style="list-style-type: none"> Pint Containers: Treats up to 5,300 square feet of southern turfgrass and/or up to 2,650 square feet of northern turfgrass. Quart Containers: Treats up to 10,600 square feet of southern [sensitive] turfgrass [lawns] and/or up to 5,300 square feet of northern [hardy] turfgrass [lawns]. Gallon Containers: Treats up to 42,600 square feet and/or up to 21,300 square feet of northern turfgrass. 	
KILLS WEEDS:	Dandelion, clover, chickweed plus over xxx additional broadleaf weeds

WHERE TO USE:	Intended for use only on home lawns.
AMOUNT TO USE:	<p>For Southern [Sensitive] turf [lawns]: [bentgrass, hybrid bermudagrass, buffalograss, St. Augustinegrass (except 'Floratam'), and centipedegrass]</p> <p>Use: 3.0 fluid ounces (6 Tablespoons)/1,000 square feet</p> <p>Or</p> <p>Use: 1.5 fl.oz. per gallon of water per 500 sq.ft.</p> <hr/> <p>For Northern [Hardy] turf [lawns]: [Kentucky bluegrass, fescues, perennial ryegrass, bahiagrass, common bermudagrass, and zoysiagrass]</p> <p>Use: 6.0 fluid ounces (12 Tablespoons)/1,000 square feet</p> <p>Or</p> <p>Use: 3.0 fl.oz. per gallon of water per 500 sq.ft.</p>
	<p>Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx • http://www.company name.com</p>

WHERE TO USE:

EH-1458 Herbicide is intended for use only on residential (home) lawns including Kentucky bluegrass, perennial ryegrass, fescues, common bermudagrass, hybrid bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass and St. Augustinegrass (except 'Floratam').

A. For [hand-operated] tank sprayers

[SPOT TREATMENTS WITH HAND-OPERATED TANK SPRAYERS, PUMP-UP SPRAYERS OR COMPRESSED AIR SPRAYERS:]

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre.

Spot treat broadleaf weeds with this product as they appear. Repeat treatment for hard-to-control weeds in 2 to 3 weeks 30 days. Do not exceed specified dosages for any area.

The maximum application rate is 3.0 fl.oz. per 500 sq.ft. per application (0.09 lb MCPP-p acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass, and St. Augustinegrass (except 'Floratam'): Mix 1.5 fluid ounce (3 Tablespoons) of product per gallon of water to cover 500 square feet of lawn.

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass: Mix 3.0 fluid ounces (6 Tablespoons) of product per gallon of water to cover 500 square feet of lawn.

[Equal measures: 1 Tablespoon (Tbs.) = 3 Teaspoons (tsp.) = 1/2 fluid ounce (fl.oz.)]

B. For Ortho® Dial-N-Spray® hose-end sprayers

[optional - THREE EASY STEPS TO MIX AND APPLY

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.

2. Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying.
3. See below for FILLING the Ortho® Dial-N-Spray® hose-end sprayer. Spray evenly over the measured area.]

[Narrative format:]

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (do not use on 'Floritam' St Augustinegrass)

- Determine the total lawn area to be sprayed.
- RATE PER GALLON: Set the dial to 1.5 oz.
- FILLING: Add 3.0 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar.
- Spray evenly over the measured area.

OR

- RATE PER GALLON: Set the dial to 1.5 oz.
- FILLING: Add 15.0 fl.oz of product into sprayer jar. Do not add water to jar.
- Spray evenly over 5,000 sq.ft. Any unused product can be poured back into its original container.

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass:

- Determine the total lawn area to be sprayed.
- RATE PER GALLON: Set the dial to 2.5 oz.
- FILLING: Add 6.0 fl.oz of product into sprayer jar for each 1,000 sq.ft. of lawn. Do not add water to jar.
- Spray evenly over the measured area.

OR

- RATE PER GALLON: Set the dial to 2.5 oz.
- FILLING: Add 30 fl.oz of product into sprayer jar. Do not add water to jar.
- Spray evenly over 5,000 sq ft. Any unused product can be poured back into its original container.

[Tabular Format]

Lawn Grass	Sprayer Type	Dial Setting	Product To Measure	Area to Treat
Bentgrass, hybrid bermudagrass, buffalograss, centipedegrass, and St. Augustinegrass (do not use on 'Floritam' St. Augustinegrass)	Tank Sprayer	NA	1.5 fl.oz. (3 Tbs.) into 1 gallon of water	500 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	1.5 oz.	Add 3.0 fl.oz. (6 Tbs.) into sprayer jar for each 1,000 square feet. Do not add water to jar.	1,000 sq.ft.
Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass, and zoysiagrass	Tank Sprayer	NA	3.0 fl.oz. (6Tbs.) into 1 gallon of water.	500 sq.ft.
	Ortho® Dial-N-Spray® Or Ortho Plant & Weed Sprayette®	2.5 oz.	Add 6.0 fl.oz. into sprayer jar for each 1,000 square feet. Do not add water to the jar.	1,000 sq.ft.

[Equal Measures: 1 Tablespoons = 3 Teaspoons (tsp.) = 1/2 fl.oz.]

C. Ready-to-use hose-end sprayers

[optional: PRODUCT FACTS

THREE EASY STEPS TO MIX AND APPLY or HOW TO USE READY-SPRAY® SYSTEM

1. Measure the total (entire) lawn area to be sprayed. For rectangular areas, multiply the length by width to obtain total square feet.
2. See below for directions for the ready-to-use hose-end sprayer.
3. Spray evenly over entire lawn.

CONNECT

1. Connect Sprayer to hose.
2. Turn on Water.

SPRAY

1. To begin spraying, point nozzle in the direction you want to spray.
2. Bend small plastic tab back and turn knob clockwise to ON position.
3. Spray evenly over measured area.

FINISH

1. To STOP spraying, rotate the knob counterclockwise to the OFF position.
2. Turn OFF water.
3. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob counterclockwise to the OFF position.
4. Disconnect Sprayer from hose.]

AMOUNT TO SPRAY [USE]

Use the amount indicated for your type of lawn:

For bentgrass, hybrid bermudagrass, buffalograss, centipedegrass and St. Augustinegrass (except 'Floratum'): (Quart size): This container covers approximately 10,600 square feet. (Pint size): This container covers approximately 5,300 square feet

For Kentucky bluegrass, bahiagrass, common bermudagrass, fescues, perennial ryegrass and zoysiagrass: (Quart size): This container covers approximately 5,300 square feet. (Pint size): This container covers approximately 2,650 square feet

BROADLEAF WEEDS CONTROLLED:			
EH-1458 Herbicide will control or suppress the following broadleaf weeds. A partial list is presented below.			
Amaranth, purple	Cudweed	Mustard, black	Speedwell, corn
Amaranth, slender	Daisy, English	Mustard, blue	Speedwell, ivyleaf
Aster, heath	Daisy, oxeye	Mustard, hedge	Speedwell, Persian
Aster, white heath	Dandelion, common	Mustard, Indian	Speedwell, purslane
Aster, purple	Dandelion, false	Mustard, tansy	Speedwell, slender
Aster, white prairie	Day-flower	Mustard, tumble	Speedwell, snow
Artichoke	Deadnettle	Mustard, white	Speedwell, thyme-leaf
Austrian fieldcress	Dichondra	Mustard, wormseed	Spiny amaranth
Bedstraw, catch-weed	Dock, broadleaf	Nettle, stinging	Spurge, spotted
Bedstraw, smooth	Dock, curly	Nettle, tall	Spurge, prostrate
Bedstraw, northern	Dogbane	Nettle, wood	Star of Bethlehem
Bedstraw, pineywoods	Dog-fennel	Parsley-piert	Strawberry, Indian mock
Beggar-ticks, hairy	Dollarweed (Pennywort)	Parsnip	Tansy ragwort
Betony, Florida	Dove-weed	Pearlwort	Tanweed
Bindweed, field	Elderberry	Pennycress	Thistle, blessed
Bindweed, hedge	Falseflax	Pepperweed, perennial	Thistle, bull
Bitter Wintercress	False sunflower	Pepperweed, Virginia	Thistle, Canada
Bitter-cress, hairy	Fiddleneck	(Peppergrass)	Thistle, Flodman's
Bitterweed	Filaree, redstem	Pigweed, prostrate	Thistle, musk
Black-eyed Susan	Filaree, whitestem	Pigweed, redroot	Thistle, plumeless
Black medic	Fleabane, daisy	Pigweed, Russian	Thistle, prairie
Blood-flower milkweed	(annual)	Pigweed, smooth	Thistle, Russian
Blue lettuce	Fleabane, rough	Pigweed, tumble	Thistle, Scotch
Brassbuttons	Galinsoga, hairy	Pineapple weed	Thistle, wavy-leaf
Bristly oxtongue	Galinsoga, smallflower	Plains coreopsis (tickseed)	Thistle, yellow
Broomweed	Goldenrod	Plantain, black-seed	Trailing crown-vetch
Buckhorn	Ground ivy	Plantain, bracted	Velvetleaf
Bull-nettle	Groundsel	Plantain, broadleaf	Vervain, blue
Bur-clover	Gumweed	Plantain, buckhorn	Vervain, hoary
Burdock	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Burweed, lawn	Hawkweed, meadow	Plantain, narrow-leaf	Vervain, tall
Buttercup, bulbous	Hawkweed, mouse-ear	Plantain, slender	Vetch, bird

BROADLEAF WEEDS CONTROLLED:			
EH-1458 Herbicide will control or suppress the following broadleaf weeds. A partial list is presented below.			
Buttercup, creeping	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Buttonweed, common (poorjoe)	Hawkweed, yellow	Poison ivy	Vetch, milk
Buttonweed, Virginia	Healall	Poison oak	Vetch, narrow-leaf
Carolina geranium	Heart-leaf drymary	Pokeweed	Vetch, wild
Carpetweed	Hemp	Prairie sunflower	Virginia creeper
Catnip	Henbit	Prickly lettuce (compass plant)	Western clematis
Catsear, spotted (false dandelion)	Hoary-cress	Prickly sida	Western salsify
Chamber bitter	Horsenettle	Prostrate knotweed	Wild aster
Chickweed, common	Horseweed	Puncturevine	Wild buckweat
Chickweed, mouse-ear	Jimsonweed	Purslane, common	Wild carrot
Chickweed, sticky	Johnny-jumpup violet	Pusley, Brazilian	Wild four-o'clock
Chicory	Knawel	Pusley, Florida	Wild garlic
Cinquefoil	Knotweed, prostrate	Pusley, large flower	Wild lettuce
Clover, crimson	Kochia	Ragweed, bur	Wild marigold
Clover, Alyce	Lambsquarters	Ragweed, common	Wild mustard
Clover, hop	Lespedeza	Ragweed, lance-leaf	Wild onion
Clover, rabbit-foot	Mallow, alkali	Ragweed, western	Wild parsnip
Clover, red	Mallow, bristly	Redstem filaree	Wild radish
Clover, strawberry	Mallow, common	Rough cinquefoil	Wild rape
Clover, yellow sweet	Mallow, dwarf	Scarlet pimpernel	Wild strawberry
Clover, white sweet	Mallow, Venice	Shepherd's purse	Wild sweet potato
Clover, white	Marcela	Smartweed, ladysthumb	Wild violet
Cockle	Matchweed	Smartweed, pale	Woodsorrel, creeping
Cocklebur	Mexican-weed	Smartweed, Pennsylvania	Woodsorrel, violet
Cocklebur, spiny	Moneywort	Smooth chaff-flower	Woodsorrel, yellow (<i>Oxalis</i>)
Common mullein	Morningglory, bigroot	Smooth dock	Woolly croton
Corn spurry	Morningglory, ivyleaf	Sorrel, red (sheep)	Wormseed
Creeping beggar-weed	Morningglory, red	Sowthistle, annual	Yarrow, common
Creeping Jenny (creeping Charlie)	Morningglory, tall	Sowthistle, spiny	Yarrow, western
Cupid's shaving brush	Morningglory, wolly	Spanish needles	Yarrow, yellow
	Mugwort	Speedwell, bi-lobed	Yellow Rocket
	Mustard, ball	Speedwell, common	

[Optional text for further use clarification]

WHEN TO USE

Spray when broadleaf weeds are young and actively growing for best results.

EH-1458 Herbicide may be applied in the spring, summer or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Broadcast (entire areas) applications or follow-up applications at 2 to 6 week 30 day intervals are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.

- **For Established Lawns:** Apply when daytime temperatures are below 90°F.
- **For Newly Seeded Areas:** Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- **For Newly Sodded, Sprigged or Plugged areas:** Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
- **Reseeding:** Treated areas can be reseeded at 2 weeks after the application of this product.
- **Limitations:**
 - The maximum per-application rate is 6.0 fl.oz. of product per 1,000 sq.ft per application, and the maximum number of broadcast applications is two (2) per year with a minimum of 30 days

between applications. The maximum seasonal rate is 12.0 fl.oz. of product per 1,000 sq.ft. excluding spot treatments.

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- **Irrigation:** Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigations (watering) before and after the application. Avoid applications of this product when turf is under stress since injury may result.
- **Mowing:** Avoid mowing within 1 to 2 days before and after the application of this product.
- **Non-target plants:** Do not apply this product to carpetgrass, dichondra, food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. ~~Avoid spraying exposed roots of ornamentals and trees.~~

[end optional text]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

~~**CONTAINER DISPOSAL:** If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partially filled: Call your local solid waste agency [or toll free number (1-800-XXX-XXXX)] for disposal instructions. Never place unused product down any indoor or outdoor drain.~~

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____

- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds.
- Kills weeds - not lawn grasses.
- Kills weeds - not lawns.
- For lawns containing Kentucky bluegrass, fescues, bentgrass, perennial ryegrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam'), and buffalograss.
- Can be applied with hose-end or pump-up (tank) sprayers.
- Kills (broadleaf) weeds.
- For northern and southern grasses.
- Won't harm lawn grasses when used as directed.
- Kills weeds in your lawn.
- Formulated for use on northern and sensitive southern lawns.
- Kills major broadleaf lawn weeds: List of the weed species.
- Helps provide a beautiful lawn.
- For cool and warm season lawn grasses.
- Kills major broadleaf lawn weeds including dandelion, chickweed, and knotweed in lawns.
- For use on grass lawns especially Kentucky bluegrass, fescues, perennial ryegrass, bentgrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass (except 'Floritam') and buffalograss.
- Can be used on St. Augustinegrass (except 'Floritam'), bentgrass, and buffalograss.
- A Trimec® Turf Herbicide Formulation.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- "One quart covers up to xxxx square feet (of southern turfgrass)" or "One pint covers up to xxxx square feet (of southern turfgrass)".
- Treats up to (Number) square feet.
- Kills over xxx broadleaf lawn weeds including dandelion and others.
- Makes up to (Number) gallons of spray.
- Now for northern and southern lawns.
- New formula for northern and southern grasses/lawns.
- Kills major broadleaf weeds including dandelion, clover, oxalis, chickweed, knotweed plantain, henbit, spurge and others.
- Concentrate
- Specially formulated for residential use (home lawns)
 - Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge and many others.
 - One easy step to a weed free lawn - Helps promote a beautiful lawn.
 - Simple easy to use/one step application
 - Simple and easy.
 - Use (Brand Name) to kill broadleaf weeds in lawns including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge, and many others. Won't harm bahiagrass, bentgrass, bermudagrass, bluegrass, buffalograss, centipedegrass, fescues, ryegrass or zoysiagrass when used as directed.. Can also be used on St. Augustinegrass except 'Floritam'.
 - 1 pint covers 2,650 to 5,300 sq.ft. (1 quart covers 5,300 to 10,600 sq.ft.) depending on lawn. See inside booklet for complete instructions. Specially formulated for residential use.
 - CONVENIENT.
 - ECONOMICAL - Use to cover small or large areas.
 - Rapid and effective weed control for common and troublesome weed species in turfgrass.
 - Superior cool weather performance.
 - Fast acting with evidence of injury within hours. Generally, the injury symptoms can be noticed in hours and plant death can occur within 7 to 14 days.
 - Kills the weeds, not the lawn.
 - Kills broadleaf weeds

- Won't harm your lawn when used as directed.
- Kills hard to control weeds.
- Kills over xxx tough lawn weeds.
- Maximum weed control
- Kills weeds, not lawns
- The performance leader
- Treats up to xxx sq.ft.
- Tough on weeds, easy on lawns
- Results in 24 hours
- Kills over 250 weeds
- Kills the toughest weeds
- Rain-Proof(TM) - Rain or watering more than 1 hour after application will not wash away effectiveness
- Rain-Proof(TM) in 1 hour
- Rain Fast in 1 hour
- **Notice:** Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.
- **Guarantee:** If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.
- TRIMEC® is a registered trademark of PBI/Gordon Corporation.
- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group
- Kills the Root
- Results in Hours
- Kills Major Broadleaf Weeds
- ORTHO® & YOUR ENVIRONMENT Your yard is a place for your family and pets to enjoy. That's why Ortho® products are designed with care to provide effective solutions to weed problems. For best results, please follow instructions for appropriate usage, storage and disposal.
- Do not allow people or pets to enter treated area until sprays have dried.
- New Formula [for use for limited time during product introduction]

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00910.20090708.amend-proposed.doc
2. Reason for Issue: Corrected Reg No.