

2217-913

08/05/2008

1/12



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number: 2217-913  
Date of Issuance: AUG 5 2008

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance:  
**conditional**

Name of Pesticide Product:  
**EH-1451 Weed & Feed**

Name and Address of Registrant (include ZIP Code):

**PBI/Gordon Corporation**  
**1217 West 12<sup>th</sup> Street**  
**P.O. Box 014090**  
**Kansas City, MO 64101**

**Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.**

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
2. Submit the data listed below:
  - a. Within one year of the date on this registration notice, submit Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies.
3. Make the following label changes:
  - a. Under Hazards to Humans and Domestic Animals, add bolded statement "Causes moderate eye irritation. Avoid contact with eyes or clothing. **[Wear protective eyewear.] Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.** Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals."

Signature of Approving Official:  
Joanne I. Miller  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
*Joanne I. Miller*

Date: AUG 5 2008

- b. Under the Storage and Disposal section, change the bullets to read **Pesticide Storage and Pesticide Disposal**.
- c. Revise the EPA Registration Number to read, "**EPA Reg. No. 2217-913.**"

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

3/12

ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
AUG 5 2008

# EH-1451 WEED AND FEED

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

## EPA Reg. No. 2217-XXX

2217-913

**ACTIVE INGREDIENT:**

2,4-D, 2-ethyhexyl ester .....	1.108%
Mecoprop-p .....	0.167%
Dicamba .....	0.071%
Dithiopyr .....	0.160%
<b>OTHER INGREDIENTS:</b> .....	<b>98.494%</b>
<b>TOTAL</b> .....	<b>100.000%</b>

**THIS PRODUCT CONTAINS:**

- 14.70 lbs 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735% (CAS 1928-43-4)
  - 3.34 lbs (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167% (CAS 16484-77-8)
  - 1.42 lbs 3,6-dichloro-o-anisic acid equivalent per ton or 0.071% (CAS 1918-00-9)
  - 3.20 lbs dithiopyr per ton or 0.160% (CAS 97886-45-8)
- Isomer Specific By AOAC Methods.

<b>GUARANTEED FERTILIZER ANALYSIS</b>	
Total Nitrogen (N) .....	x.x%
x.x% Ammoniacal Nitrogen	
x.x% Water Insoluble Nitrogen	
x.x% Urea Nitrogen	
x.x% Other Water Soluble Nitrogen	
Available Phosphate (P2O5) .....	x.x%
Soluble Potash (K2O) .....	x.x%
Iron (Fe) .....	x.x%
Nutrient Sources: [Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer.]	

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

**For residential use by homeowners:**  
**New York State:** Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year (0.25 lb. of dithiopyr per acre per year).

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**



4/12

## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.</p>	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.

### **General Precautions and Restrictions:**

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Do not allow people or pets to enter the treated area until dusts have settled.

### **PRODUCT DESCRIPTION**

EH-1451 Weed & Feed is formulated (designed) to provide control of most annual grasses and certain broadleaf weeds commonly found in home lawns.

EH-1451 Weed & Feed contains three components: a preemergent herbicide, a lawn fertilizer, and a post emergent broadleaf weed killer.

- The preemergent herbicide can be applied to established turfgrass to control crabgrass goosegrass, foxtails, and other undesirable annual grasses.
- The fertilizer contains slow release nutrients that may extend the availability for turfgrass uptake.
- The post emergent broadleaf herbicide will control over 250 lawn weeds including dandelion, chickweed, thistle and clover. A partial list of broadleaf weeds is presented on this label.

**WHERE TO USE**

**EH-1451 Weed & Feed may be used on the following established turf species:**

**Established Cool Season Turfgrass:**

- Kentucky bluegrass
- perennial ryegrass
- fine fescues
- tall fescue

**Established Warm Season Turfgrass:**

- bahiagrass
- bermudagrass
- buffalograss
- zoysiagrass and

Do not use EH-1451 Weed & Feed on dichondra, carpetgrass or colonial bentgrass.

Do not use EH-1451 Weed & Feed on St. Augustinegrass in Florida.

Do not use this product on turfgrass where desirable clovers are present.

**Guidelines to follow:**

The use of EH-1451 Weed & Feed on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

**Other Use Precautions:**

- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches.
- Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans.
- Do not apply when windy.
- Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Do not exceed specified dosage for any area.
- Do not use clippings from treated areas for composting or for mulching vegetables or fruit trees.

6/12

<b>WEEDS CONTROLLED</b>			
When applied as directed, this product will control and/or suppress* the following weeds:			
<b>Annual Grass Weeds:</b>			
Annual bluegrass* ( <i>Poa annua</i> ) Annual ryegrass	Barnyardgrass* Crabgrass	Foxtail (green and yellow)	Goosegrass* (crowfootgrass)
<b>A partial listing of established broadleaf weeds:</b> [see appendix for complete list or include here]			
Bedstraw Beggarticks Black medic Broadleaf plantain Buckhorn plantain Burclover Burdock Carpetweed Chickweed, common Chickweed, mouseear Chicory Clover, hop Clover, strawberry Clover, white	Curly dock Daisy, oxeye Dandelion Dichondra False dandelion Ground ivy Healall Henbit Knotweed Lambsquarters Lespedeza Mallow Matchweed Momingglory	Mustard Narrowleaf plantain Oxalis Parsley-piert Pearlwort Pennycress Pepperweed Pigweed Plantain Poison ivy Poison oak Pokeweed Purslane, common Ragweed	Red sorrel Redroot pigweed Sheep sorrel Shepherdspurse Slender plantain Smooth dock Smooth pigweed Spurge Thistle Woodsorrel Yarrow and other broadleaf weeds
* for acceptable control of barnyardgrass and goosegrass, two applications made at a 6 to 10 week interval are needed			

**WHEN TO APPLY**

**A. Established lawns:**


This product may be applied as a single application or as split applications in the spring, early summer, or fall.

For the best results in the spring, apply this product one to two (1 to 2) weeks before the expected germination of crabgrass. When the period of germination for crabgrass and other weeds listed on this label extends into early summer, then a second application can be made at 6 to 10 weeks after the initial application.

**For crabgrass and other listed grasses:** This product is a preemergent herbicide to control crabgrass (smooth, southern and large) and other susceptible annual grasses. This product may also provide early post emergent control of crabgrass. However, post emergent crabgrass control will be obtained only when this product is applied before the first tiller stage of crabgrass.

[start optional growth stage description]

The graphics below indicate the growth stages of crabgrass, and the optimum timing for application of this product to assure control:



BEST Grassy weed control	GOOD Grassy weed control	POOR Grassy weed control
1 to 2 weeks before germination	Stages 9, 10, 11, or 12 above	Stage 13 or greater

[end optional growth stage description]

**For broadleaf weed control:**

Best results are obtained during the spring and fall when broadleaf weeds are present, young and actively growing under adequate soil moisture and favorable temperatures (50° to 80°F).

**HOW TO APPLY**

This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain; poor results will occur if it is applied to dry lawns. The lawn should not be watered or disturbed for 24 hours after application. Delay mowing for 3 days after application.

Apply this product when the wind is calm. Avoid spreading granules onto flowers, vegetables, and ornamental shrubbery or desirable plants.

**HOW MUCH TO USE**

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

**For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass:**

Apply 2.5 to 3.6 pounds of product per 1,000 square feet of established turfgrass.

**For hybrid bermudagrass, bahiagrass and buffalograss:**

Apply 2.0 to 3.0 pounds of product per 1,000 square feet of established turfgrass.

**For residential use by homeowners:**

**New York State:** Do not exceed one application at the recommended rate of 3.6 pounds of product per 1,000 sq.ft. per year (0.25 lb. active ingredient of dithiopyr per acre per year).

**Other States:** The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application (1.15 lbs 2,4-D ae, 0.26 lb MCP-p acid equivalent, 0.11 lb dicamba acid equivalent, and 0.25 lb dithiopyr per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 7.2 pounds of product per 1,000 sq.ft. (2.30 lbs 2,4-D acid equivalent, 0.52 lb MCP-p acid equivalent, 0.22 lb dicamba acid equivalent, and 0.50 lb. dithiopyr per acre).

**SUGGESTED SPREADER SETTINGS**

Note to reader: A listing of spreaders and suggested settings for the application rate of 2.0 to 3.6 lbs. per 1,000 sq. ft. will be presented on the container labels.

The above spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

**CALIBRATION INSTRUCTIONS** [examples]:

- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

<p><b>STORAGE AND DISPOSAL</b></p> <p><b>STORAGE:</b> Keep in original container in a cool, dry area, away from extreme heat and direct sunlight. Do not store near food feed, seeds or other pesticides.</p> <p><b>DISPOSAL: If empty:</b> Do not reuse this container. Place in trash or offer for recycling if available. <b>If partly filled:</b> Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.</p>
--

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

**APPENDIX**

**1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

**2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- Northern and Southern Grasses Bluegrass, Fescue, Zoysia, Bermuda, Bahia, Perennial Ryegrass
- Treats Northern and Southern Grasses
- Kills xx+ Weed; Types, Even the Roots..
- Kills xx+ Weeds
- Especially effective on hard-to-kill weeds
- Kills broadleaf weeds in lawns



- Kills major broadleaf weeds, even the roots. And [brand name] is formulated to kill only the weeds, not the lawn. Use as directed to kill major broadleaf weeds in your lawn.
- Kills dandelion and many other broadleaf weeds.
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion.
- Tough weed killer
- Kills even the tough[est] broadleaf weeds [like clover, dandelion, oxalis]
- Destroys the hard-to-kill broadleaf weeds [like clover, dandelion; oxalis]
- Kills [major] broadleaf weeds [like clover, dandelion, oxalis]
- Provides broadleaf control
- Controls [Kills] clover, dandelion, plantain, chickweed and [other broadleaf,weeds] [others as listed]
- Clobbers clover
- Kills the root
- Kills weed roots!
- Kills more weeds
- Kills tough lawn weeds
- Consistently fast on: tough weeds like clover, plantain, ground ivy and spurge [various other weeds listed]
- Starts working on contact
- [cool][all] weather performance
- Proven [all] [cool] weather performance
- For home lawn care
- For home lawns only
- For a beautiful weed-free lawn
- Helps keep lawns weed-free
- Kills the weeds, not the lawn
- Kills weeds, not lawn grasses
- Kills [down to] the root, [not the lawn]
- For use on lawns
- Targets weeds, leaves [grasses][the lawn] unharmed
- Guaranteed [Results] or Your Money Back\*
  - If you are not satisfied with this product for any reason, send your written request for a refund to the address at left, together with the original dated store receipt showing the price you paid. Please allow 6 to 8 weeks for processing.
- Controls Dandelion, White Clover, Plantain, Chickweed, Spurge and Many Other Listed Weeds
- Greens in 72 Hours & Promotes Thick, Green Growth
- Contains Composite Timed-Release (CTR) Nitrogen for Extended Feeding!
- Kills Over xxx Weed Types
- Kills Weeds While Feeding Your Lawn
- Encourages Quick Greening & Provides Extended Nitrogen Feeding up to 2 Months
- Kills xx+ Weed Types Including Roots
- Especially Effective on Hard-to-Kill Weeds
- Kills major broadleaf weeds including roots. And (brand name) is formulated to kill only the weeds, not the lawn. Use as directed to kill major broadleaf weeds in your lawn.
- For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass
- Sta-Green Turf Note – Control Those Pesky Weeds
  - Many broadleaf weeds (dandelion, chickweed, henbit) germinate in the fall but are not visible in your lawn until spring. Applying (product name) in the fall can provide effective results. If needed, you can make a follow-up application in the spring.
- Project Guide – For a thicker, greener, more beautiful lawn, follow (Product Program). It's easy to follow and can be started at anytime by simply applying the product that corresponds with the current season!
- (Product Name) is specially formulated to complement the phosphorus already present in some soils and give your turf the additional primary nutrients needed for a good turf feeding. A soil test can help

you determine if your soil has an adequate supply of phosphorus and if this is the best product for your soil type. Proper watering and mowing practices along with regular feedings will help promote a beautiful green lawn.

- Prevents Crabgrass and Fertilizes
- Weed 'n Feed Plus Crabgrass Preventer
- Eliminates Lawn Weeds - One application does it all - crabgrass, dandelion, viney weeds, even clover.

**3. Trademark acknowledgement statements:**

- DIMENSION® is a registered trademark of Dow AgroSciences, L.L.C.
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**4. Fertilizer Labeling Requirements in States:**

- Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>.
- Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing \_\_\_ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.
- If the use rate applies less than 1 pound nitrogen per 1,000 sq.ft. per application, the following statement must appear: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq.ft.

**Fertilizer Grades:**

- Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5, and K2O include the following:

18-0-6	22-4-11	26-4-4	30-4-4
19-6-8	23-3-7	26-4-6	32-3-3
22-3-3	25-3-3	28-0-3	32-4-8
22-3-6	25-4-8	29-3-4	24-4-8
22-3-17	26-3-6	30-3-3	

[Controls][Kills][253][more than ###] weeds! (List of weeds may include the following):

Amaranth, purple	Cupid shaving Brush	Mustard, ball	Speedwell, common
Amaranth, slender	Cudweed	Mustard, black	Speedwell, corn
Aster, heath	Daisy, English	Mustard, blue	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, hedge	Speedwell, persian
Aster, purple	Dandelion, common	Mustard, indian,	Speedwell, purslane
Aster, white prairie	Dandelion, false	Mustard, tansy	Speedwell, slender
Artichoke	Day flower	Mustard, tumble	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, white	Speedwell, thymeleaf
Bedstraw, Catchweed	Dichondra	Mustard, wormseed	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Nettle, stinging (burning)	Spurge, spotted
Bedstraw, Northern	Dock, curly	Nettle, tall	Spurge, prostrate
Bedstraw, pineywoods	Dogbane	Nettle, wood	Spurweed
Beggarticks, hairy	Dogfennel	Parsley-piert	Star of Bethlehem
Betony, Florida	Dollarweed (Pennywort)	Parsnip	Strawberry, India mock
Bindweed, Field	Dove weed	Pearlwort	Tansy ragwort
Bindweed, Hedge	Elderberry	Pennycress (frenchweed)	Tanweed
Bitter Wintercress	Falseflax	Pepperweed, perennial	Thistle, blessed
Bittercress, hairy	False sunflower	Pepperweed, virginia	Thistle, bull
Bitterweed	Fiddleneck	(Peppergrass)	Thistle, canada
Black-eyed Susan	Filaree, redstem	Pigweed, prostrate	Thistle, flodman (prairie)
Black Medic	Filaree, whitestem	Pigweed, redroot	Thistle, musk
Blood flower milkweed	Fleabane, daisy (annual)	Pigweed, russian	Thistle, platte
Blue lettuce	Fleabane, rough	Pigweed, smooth	Thistle, plumeless
Brassbuttons	Galinsoga, hairy	Pigweed, tumble	Thistle, russian
Bristly ox tongue	Galinsoga, smallflower	Pineapple weed	Thistle, scotch
Broomweed	Goldenrod	Plains coreopsis (tickseed)	Thistle, wavyleaf
Buckhorn	Ground ivy	Plantain, blackseed	Thistle, yellow (horrible)
Bullnettle	Groundsel	Plantain, bracted	Trailing crownvetch
Burclover	Gumweed	Plantain, broadleaf	Velvetleaf
Burdock	Hairy fleabane	Plantain, buckhorn	Vervain, blue
Burweed, lawn (spurweed)	Hawkweed, meadow	Plantain, hoary	Vervain, hoary
Buttercup, bulbous	Hawkweed, mouseear	Plantain, narrowleaf	Vervain, prostrate
Buttercup, creeping	Hawkweed, orange	Plantain, slender	Vervain, tall
Buttonweed, common (poorjoe)	Hawkweed, yellow	Plantain, woolly	Vetch, bird
Buttonweed, virginia	Healall	Poison ivy	Vetch, hairy
Carolina geranium (wild)	Heartleaf drymary	Poison oak	Vetch, milk
Carpetweed	Hemp	Pokeweed	Vetch, narrowleaf
Catnip	Henbit	Prairie Sunflower	Vetch, wild
Catsear, spotted (false dandelion)	Hoary cress	Prickly lettuce (compass plant)	Virginia creeper
Chamber bitter	Horsenettle	Prickly sida	Western clematis
Chickweed, common	Horseweed	Prostrate knotweed	Western salsify
Chickweed, mouseear	Jimsonweed	Puncturevine	Wild aster
Chickweed, sticky	Johnny-Jumpup violet	Purslane, common	Wild buckweat
Chicory	Knawel	Pusley, brazil	Wild carrot
Cinquefoil	Knotweed, prostrate	Pusley, florida	Wild four-o'clock
Clover, crimson	Kochia	Pusley, large flower	Wild garlic
Clover, alyce	Lambsquarters	Ragweed, bur	Wild lettuce
Clover, hop	Lespedeza	Ragweed, common	Wild marigold
Clover, rabbit foot	Mallow, alkali	Ragweed, lancleaf	Wild mustard (yellow rocket)
Clover, red	Mallow, bristly	Ragweed, western	Wild onion
Clover, strawberry	Mallow, common	Redstem filaree	Wild parsnip
Clover, yellow sweet	Mallow, dwarf	Rough cinquefoil	Wild radish
Clover, white sweet	Mallow, venice	Scarlet pimpernel	Wild rape
Clover, white	Marcela	Shepardspurse	Wild strawberry
Cockle	Matchweed	Smartweed, ladythumb	Wild sweet potato
Cocklebur	Mexicanweed	Smartweed, pale	Wild violet
Cocklebur, oriental	Moneywort	Smartweed, pennsylvania	Woodsorrel, creeping
Cocklebur, spiny	Morningglory, bigroot	Smooth chaff-flower	Woodsorrel, violet
Common mullein	Morningglory, ivyleaf	Smoothdock	Woodsorrel, yellow (Oxalis)
Corn Spurry	Morningglory, mexican	Sorrel, red (sheep)	Woolly croton
Creeping Beggarweed	Morningglory, red	Sowthistle, annual	Wormseed
Creeping Jenny (creeping charlie)	Morningglory, tall	Sowthistle, spiny	Yarrow, common
	Morningglory, wolly	Spanish needles	Yarrow, western
	Mugwort	Speedwell, bi-lobed	Yarrow, yellow

## **DOCUMENT CONTROL INFORMATION**

**1. Unique Label Identifier:** 002217-00XXX.20080211.doc

**2. Previous Version:** None

**3. Reason for Issue:** New Registration