



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 5, 2023

Eric D. Smith
Director
Regulatory Affairs
PBI/Gorgon Corporation
Post Office Box 014090
Kansas City, Missouri 64101

Subject: Registration Review Label Mitigation for Quinclorac
Product Name: EH-1465 Herbicide
EPA Registration Number: 2217-919
Application Date: January 31, 2019
Decision Number: 592120

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Quinclorac Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

Page 2 of 2
EPA Reg. No. 2217-919
Decision No. 592120

approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


If you have any questions about this letter, please contact Caleb Carr via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure



2,4-D QUINCLORAC DICAMBA	GROUP	4	HERBICIDE
--------------------------------	-------	---	-----------

EH-1465 HERBICIDE

EPA Reg. No. 2217-919

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	22.82%
Quinclorac	7.56%
Dicamba, dimethylamine salt	2.09%
INERT INGREDIENTS:	<u>67.53%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 1.78 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 18.95%
 - 0.71 lbs. 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 7.56%
 - 0.17 lbs. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.74%
- Isomer Specific By AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet [inside pages] for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.

Net Contents: _____
EPA Est. No. _____

Company Name and Address:
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed. Do not get in eyes or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators, and other handlers must wear:

- protective eyewear,
- long-sleeved shirt and long pants,
- shoes and socks, plus
- waterproof gloves (except for applicators using ground boom equipment) and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-888-800-5556 for emergency medical treatment information.

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls,
- waterproof gloves, and
- shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

PRODUCT DESCRIPTION:

EH-1465 Herbicide is a product that contains three (3) active ingredients including quinclorac that broaden the spectrum of weed control. EH-1465 Herbicide is a versatile herbicide that will control many troublesome grassy and broadleaf weeds in established turfgrass when used as directed.

WEED RESISTANCE MANAGEMENT

For resistance management, this product contains Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 877-800-5556.

WHERE TO USE:

This product provides selective broadleaf control in cool-season and warm-season turfgrass in five (5) use sites. See table 1 for turf grass species that can be treated with EH-1465 Herbicide.

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
- **Non-cropland sites** include Highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations.
- **Agricultural site:** Commercial sod production

Prohibitions of Sites:

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.
- **DO NOT** apply EH-1465 Herbicide to the following:
 - Bahiagrass
 - Bentgrass, (creeping, 'Seaside', 'Colonial')
 - Carpetgrass
 - Centipedegrass
 - Dichondra
 - St. Augustinegrass
 - Greens and collars at golf courses
 - Lawns with desirable clovers or legumes
 - Ornamentals (flowers, trees, groundcovers, landscape beds and shrubs)

Turfgrass tolerance:

- Turfgrass tolerance to this product may vary, and temporary turfgrass yellowing may occur on fine fescues and bermudagrass (see specific Bermudagrass sections).
- Tolerant turf species listed on this label may exhibit temporary turf injury. The best tolerance occurs under optimal conditions for the turfgrass. Adverse environmental conditions may reduce the selectivity on the turfgrass. Injury may occur under marginal conditions (e.g. low temperatures and drought stress) or under extreme conditions (e.g. high temperatures and high humidity). To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Under any of these stress conditions, to the extent consistent with applicable law, any turf damage caused by the use of this product is beyond the control of PBI/Gordon Corporation and all risk is assumed by the buyer and/or user.
- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Use adjuvants and spray additives or tank-mix combinations only when your experience indicates that the tank mixture will not result in objectionable turf injury. [optional text: See "SPRAY PREPARATION: Additions of adjuvants (and/or Tank Mixtures) to improve Grassy Weed Control:" section of this label. End of optional text]
- Do not broadcast apply this product when temperatures are above 90°F; some injury can also be expected with spot treatments when air temperatures exceed 90°F.

CULTURAL TIPS:

For newly seeded areas:

Delay application of this product to grass seedlings until after the second or third mowing, or 28 days after emergence.

For newly sodded, sprigged, or plugged areas:

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application. If dry conditions exist, a scheduled irrigation or watering 24

hours before and 24 hours after application is recommended. If rainfall does not occur in 2 to 7 days after application, irrigation of at least 1/2 inch is recommended.

Mowing:

Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury. Clippings from the first three mowings should be left on the treated area. Do not use these clippings as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

SPRAY PREPARATION:

Mixing with Water:

Add one-half the required amount of water to the spray tank, then add EH-1465 Herbicide slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying.

When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Mixing with Liquid fertilizers:

In certain applications, liquid fertilizer may replace part of the water as a tank-mix.

ALWAYS PREMIX EH-1465 Herbicide WITH WATER BEFORE ADDING TO FLUID FERTILIZERS. For liquid nitrogen solutions such as UAN or urea solutions, use a premix of 1 part of this product with 4 parts of water or use a premix with a 1:4 ratio of product to water. For other fluid fertilizers such as suspensions, use a premix of 1 part of this product with 50 to 60 parts of water.

Use suitable sources and rates of fertilizer based upon local recommendations. Refer to the mixing directions on the labels of the liquid fertilizers. Always perform a jar test for compatibility before large scale mixing.

The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

[Optional text]

Additions of Adjuvants (and/or Tank Mixtures) to Improve Grassy Weed Control:

Methylated seed oil is a suitable adjuvant for post-emergent applications and may cause slight yellowing to desirable turf. Chelated irons or sprayable nitrogen can reduce the slight yellowing that may occur on some turfgrass species. Methylated seed oil used must meet the following criteria:

- Be non-phytotoxic
- Contain only EPA-approved ingredients
- Provide good mixing quality in the jar test, and
- Be used only when your experience indicates that this tank mixture will not result in objectionable turfgrass injury.

Do not include additives when tank-mixing with oil-based products or EC (Emulsifiable Concentrate) products as this may cause phytotoxicity (yellowing) of desirable turfgrass. Other additives or adjuvants may be used, however, certain additives may reduce the selectivity on the turfgrass. Under some environmental conditions some of these products may cause phytotoxicity (yellowing) of desirable turfgrass. Use these adjuvants, spray additives or tank-mix combinations, only when your experience indicates that the tank mixture will not result in objectionable turf injury. **[end optional text]**

GROUND EQUIPMENT:

Spray equipment: Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate flat fan nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Hand-operated sprayers including backpack sprayers and pump-up sprayers are appropriate for small turfgrass areas. Calibration and proper application are essential when using this product.

Cleaning spray equipment: Clean sprayer before and after using this product. Use soap, household ammonia, detergent and water, or a spray tank cleaner and rinse thoroughly. Cross-contamination may cause physical incompatibility (mixing problems) or result in turf injury.

Spray distribution:

- The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator.
- Uniform applications are essential when using this product. Over-application, excessive overlaps, or rates above those specified on this label can cause turf injury.
- Avoid spray overlaps with hand-held equipment: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage. To avoid excessive spray pattern overlaps, a spray colorant may be used.
- This product may cause injury to susceptible/non-target plants at the use site by contacting the foliage, stems, or roots. To prevent injury to susceptible crops and other desirable broadleaf plants including but not limited to cotton, legumes, tobacco, tomatoes, garden/vegetable crops, and ornamentals (flowers, trees, and shrubs) avoid contact with the spray solution, spray droplets, and spray mist (fine droplets).

Chemigation: Do not apply this product through any type of irrigation system.

Aerial application: Do not apply as an aerial application.

Spray Drift Management

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or very coarse spray, apply only as a coarse or very coarse spray (ASABE standard S572.1).

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a medium or coarser spray (ASABE standard S572.1).

Wind Speed

Do not apply when wind speeds are greater than 10 mph at the application site. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Additional requirements for ground boom application: Do not release spray at a height higher than 30 inches above the ground.

POST EMERGENT CONTROL OF GRASSY WEEDS

EH-1465 Herbicide can provide post emergent control and suppression of certain grassy weeds in turfgrass. Apply this product to weeds that are young and actively growing for best results (see Table 1). Applications under adequate soil moisture conditions are preferred to summer treatments as generally, applications in the summer to older, drought stressed weeds are less effective.

Broadcast or spot treatments with one or two applications will control or suppress these grassy weeds:

- Crabgrass, (large and smooth)
- Foxtail, (green, yellow and giant)
- Signalgrass, (broadleaf)
- Barnyardgrass

Weed control is affected by the spray volume, timing and the weed growth stages (see Table 2). Generally, apply this product to actively-growing weeds during the early growth stages for the best results. Crabgrass can be controlled during mature (late season) growth stages.

POST EMERGENT CONTROL OF BROADLEAF WEEDS

Apply this product to broadleaf weeds that are young and actively growing for the best results. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective. Fall applications provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

USE RATES AND SPRAY VOLUMES:

Use rates and spray volumes of EH-1465 Herbicide are presented in Table 3.

Use the lower application rate (5.5 pints/Acre) for bermudagrass and for light (sparse) infestations of crabgrass and annual grasses in the early growth stages under conditions of active growth. Use the higher application rates within the specified range (up to 6.75 pints/Acre) for hard-to-control weeds such as dense infestations, perennial weeds or for weeds beyond the appropriate growth stages.

Spot treatments during the spring and summer are suitable for sparse infestations or as a follow-up treatment to a broadcast application on an "as-needed" basis.

SPRAY SCHEDULE: Use a single application for light infestations and two (2) applications for dense infestations.

Applications: Second or follow-up applications as either broadcast or spot treatments should be made 30 days after the initial application on ornamental turfgrass and are recommended for more mature weeds, for dense infestations and for adverse environmental conditions.

Other situations that may need two broadcast or follow-up treatments include the following:

- Under certain conditions, applications of this product at the 3 to 4 tiller stage of the annual grasses may not provide complete control.
- All weed grasses do not germinate at the same time. The period of germination for crabgrass and annual grasses can extend into the summer after the initial application of this product and results may be poor and erratic.
- Dense infestations of weeds may prevent thorough spray coverage of the target weeds.
- Biotypes of large and smooth crabgrass in California have shown varied response to quinclorac. If control failure occurs following a second application, do not reapply this product. Change to a herbicide with a different mode of action [such as Gordon's Trimec Plus Post-Emergent Grass & Broadleaf Herbicide].

Extremes in environmental conditions, ie. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. Under dry conditions, the expression of herbicide symptoms is generally delayed, and weeds hardened off by drought may be less susceptible to this product.

If objectionable turf injury occurs with the first application, then avoid making the second application of this product until the turfgrass recovery is complete.

Do not broadcast apply this product when temperatures are above 90°F; temporary turfgrass discoloration can also be expected with spot treatments when air temperatures exceed 90°F.

State Restrictions:

Arizona: Do not use this product on sod farms in Arizona.

TABLE 1. Established turf species
<p>Cool-Season Turf: Kentucky Bluegrass, Perennial ryegrass, Tall Fescue, Annual Bluegrass (<i>Poa annua</i>), Rough Bluegrass (<i>Poa trivialis</i>), Annual Ryegrass</p>
<p>Warm-Season Turf: Bermudagrass:</p> <ul style="list-style-type: none"> • Apply only when Bermudagrass is actively growing. • Expect temporary discoloration. • Some Bermudagrass hybrids may be more susceptible to discoloration. • It is impossible to test all environmental conditions and all Bermudagrass hybrids. We suggest testing this product on a small area and observe the treated area for 30 days (during normal growing conditions) to determine the acceptability of turf discoloration. • Some stunting of the Bermudagrass should be expected and turf generally recovers in 21 days. • Do not apply in the fall during fall-to-winter transition period. • Do not apply in the spring during winter-to-spring transition period. • To avoid turf injury, use only on Bermudagrass that is not under stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. • For optimum results: <ul style="list-style-type: none"> ○ Irrigate 24 hours before and 24 hours after application with 1/2 inch of water. ○ The addition of nitrogen fertilizer or chelated iron (such as FeRROMECE® Plus MICROS) may reduce some turf discoloration. ○ Spray in the morning hours and avoid applications during extreme hot or dry conditions. ○ Equipment calibration is essential and avoid spray overlaps. ○ Do not use higher pressure equipment, spray pressure should be 40 psi or lower. • Use only the lower rate shown on Table 3 (5.5 pts/Acre or 2.0 fl.oz./1,000 sq.ft.)

Turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur on fine fescues

TABLE 2. Leaf stages of grassy weeds for optimum control with this product.

Weed species	1 to 3 leaf	4 to 5 leaf (1 tiller)	6 leaf (2 tillers)	7 to 8 leaf or more (3 to 4 tillers or more)	Mature (late season)
Crabgrass, (large and smooth)	X	X	X	Footnote 1	X
Barnyardgrass Foxtail, (green, yellow and giant) Signalgrass, (broadleaf)	X	X	X	Footnote 1 (reduced control)	Footnote 1 (reduced control)

¹ Second or follow-up applications on ornamental turfgrass may be required.

Table 3. Use rates for grassy weeds and broadleaf weeds in turf.

Application schedule	Rates	Normal Spray Volume*	Maximum number of broadcast applications	Application Interval
Cool-season grasses: Kentucky Bluegrass, Perennial ryegrass, Tall Fescue, Annual Bluegrass (<i>Poa annua</i>), Rough Bluegrass (<i>Poa trivialis</i>), Annual Ryegrass	5.5 to 6.75 pints/A or 2.0 to 2.5 fl.oz./1000 sq. ft.	50 to 220 gal/A or 1.2 to 5.0 gal/1000 sq.ft.	2	30 days
Bermudagrass	5.5 pints/A or 2.0 fl.oz./1000 sq.ft.	50 to 220 gal/A or 1.2 to 5.0 gal/1000 sq.ft.	2	30 days

Low Volume Spray Equipment [such as PermaGreen Equipment and backpack sprayers]:

Equipment should be calibrated to apply at least 20 gallons of total spray volume per acre (0.45 gallons/1,000 sq.ft.). Use this lower spray volume only when your experience indicates that this volume results in effective weed coverage, weed control and will not result in objectionable turf injury.

*Note: Use the higher spray volumes for dense weed populations

Limitations on broadcast treatments for ornamental turfgrass and sod farms::

The maximum application rate is 6.75 pints of product per acre per application (1.50 lb 2,4-D ae, 0.59 lb quinclorac ai, and 0.14 lb dicamba ae per acre per application).

For ornamental turfgrass, the maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. For sod farms, the maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications.

The maximum seasonal rate is 13.5 pints of product per acre (3.0 lbs 2,4-D ae, 1.18 lb quinclorac ai, and 0.28 lb dicamba ae per acre), excluding spot treatments.

SPOT TREATMENTS FOR GRASSY WEEDS AND BROADLEAF WEEDS:

- Calibration and proper application are essential when using this product. Spray coverage should be uniform and complete.
- Over applications can result in turfgrass injury.
- **Cool-season grasses listed above:** Mix (Use) 2.5 fl.oz. of this product with 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. For low volume spray equipment see Table 3. Apply to actively growing weeds during the early growth stages as shown in Table 1.
- **For Bermudagrass:** Mix (Use) 2.0 fl.oz. of this product with 1.0 gallon of water for treatment of approximately 1,000 sq.ft. of turfgrass. For low volume spray equipment see Table 3. Apply to actively growing weeds during the early growth stages as shown in Table 1. Do not use more than 2.0 fl.oz./1000 sq.ft. on Bermudagrass.

WEEDS CONTROLLED:

EH-1465 Herbicide will control or suppress the following. Apply when weeds are young and actively growing.

Aster, white heath & white prairie	jenny)	Poison oak
Barnyardgrass	Field oxeye-daisy (*creeping oxeye)	Prickly lettuce (*compass plant)
Bedstraw	Filaree, whitestem & redstem	Puncturevine
Beggarweed, creeping	Florida betony	Purple cudweed
Bindweed	Florida pusley	Purslane
Black medic	Foxtail1 (green, yellow and giant)	Ragweed
Broadleaf plantain	Ground ivy	Redweed
Buckhorn plantain	Groundsel	Red sorrel (*sheep sorrel)
Bull thistle	Geranium (Carolina)	Shepherdspurse
Burdock, common	Hawkweed	Signalgrass1 (broadleaf)
Buttercup, creeping	Healall	Speedwell *Veronica (common, slender, and thymeleaf)
Carpetweed	Henbit	Spurge
Chickweed, common	Innocence (Blue-eyed Mary)	Thistles
Chicory	Knotweed	Virginia buttonweed1
Cinquefoil	Lambsquarters	White clover (*Dutch clover, honeysuckle clover, white trefoil & purplewort)
Clovers	Lespedeza, common	Wild carrot
Crabgrass1, 2 (large and smooth)	Mallow, common	Wild garlic
Curly dock	Matchweed	Wild geranium
Dandelion	Morningglory spp.	Wild lettuce
Dayflower	Mouseear chickweed	Wild mustard
Deadnettle	Old world diamond flower	Wild onion
Dock	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Wild strawberry
Dogfennel	Parsley-piert	Wild Violet1
Dollarweed (*pennywort)	Pennsylvania smartweed	Yarrow
English Daisy1	Pepperweed	Yellow rocket
False dandelion (*spotted catsear & common catsear)	Pigweed	
Field bindweed (*morningglory & creeping	Pineappleweed	
	Plantain	
	Poison ivy	

*Synonyms

¹A second or follow-up application may be required.

²Biotypes of large and smooth crabgrass in California have shown varied response to quinclorac. If control failure occurs following a second (or follow-up) application, do not reapply this product. Change to a herbicide with a different mode of action such as [Gordon's Trimec Plus Post-emergent Grass & Broadleaf Herbicide (EPA Reg. No. 2217-709)].

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide

or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Water-based formula
- Water-based formulation

- Proven performance
- Consistently control of lawn weeds like dandelion, spurge, clover, plantain, ground ivy and [various other listed weeds].
- Controls grassy and broadleaf weeds
- “Three-In-One” or “3-in-1”
- Controls crabgrass
- Crabgrass control
- From the makers of Trimec® herbicides.
- Trimec® is a registered trademark of PBI/Gordon Corporation.
- For information call XXX-XXX-XXXX [contact www.xxx-xxxx.com]
- Post-emergent crabgrass control, plus broadleaf weeds
- For use on: Bluegrass, Fescues, Perennial and Annual Ryegrass, and Bermudagrass
- Versatile post-emergent control of both grassy and broadleaf weeds
- Wide window of application for broadleaf weeds
- Foliar application
- Visual response
- Suitable for applications up to 90° [degrees] F
- Easy To Use Water-Based Formulation
- Water based
- Rainfast in six hours
- Rainfast in as little as six hours

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00919.20230526.amend-proposed-clean.doc

3. Reason for Issue: EPA Comments #2 + Registration Review - Quinclorac