


2217-920

01/15/2009

1/14

 <p style="text-align: center;">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p> <p style="text-align: center;">NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	EPA Reg. Number: 2217-920	Date of Issuance: 15 JAN 2009
	Term of Issuance: conditional	
	Name of Pesticide Product: EH-1466 Herbicide	
Name and Address of Registrant (include ZIP Code): PBI/Gordon Corporation 1217 West 12 th Street P.O. Box 014090 Kansas City, Missouri 64101-0090		
<p style="font-size: small;">Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data. 2. Make the following label changes: <ol style="list-style-type: none"> a. Revise the EPA Registration Number to read: EPA Reg. No. 2217-920 b. Under the "User Safety Recommendations" revise the first sentence to read: "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. c. Under the "First Aid" section, in the first sentence, add "for treatment advice" to the end of the first sentence. 3. Submit one year studies for the guidelines 830.6317 (storage stability) and 830.6320 (corrosion characteristics) and must submit the results of the studies to the Agency on completion. It is recommended that the evaluation be made at 0, 3, 6, 9 and 12 months intervals. 		
Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	Date: 15 JAN 2009	

2/14

Page number 2
EPA Reg. No. 2217-920

Submit one copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

3/14

ACCEPTED
with COMMENTS
in EPA Letter Dated:

15 JAN 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EH-1466 HERBICIDE

EPA Reg. No. 2217-XXX

2217-920

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

ACTIVE INGREDIENT:

Triclopyr BEE, butoxyethyl ester	8.40%
Sulfentrazone	0.73%
2,4-D, 2-ethylhexyl ester	31.82%
Dicamba acid	2.43%
INERT INGREDIENTS:	<u>56.62%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.50 lbs 3,5, 6-trichloro-2-pyridinyloxyacetic acid per gallon or 6.04%.
- 0.06 lbs N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.73%
- 1.75 lbs 2,4-dichlorophenoxyacetic acid equivalent per gallon or 21.18%.
- 0.20 lbs 3,6-dichloro-o-anisic acid equivalent per gallon or 2.43%.

Isomer specific by AOAC Methods.
Contains petroleum distillates

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are barrier laminate, nitrile rubber, neoprene rubber, and Viton. If you want more options, follow the instructions for category E on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus



4/14

- chemical-resistant gloves (except for applicators using ground boom equipment) and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations	
<ul style="list-style-type: none"> • Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. 	

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical information.</p>	
<p>NOTE TO PHYSICIAN: Contains petroleum distillates - vomiting may cause aspiration pneumonia.</p>	

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

These chemicals (triclopyr, 2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

5/14

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is :

- coveralls,
- chemical-resistant gloves made of any water-proof material,
- shoes plus socks and
- protective eyewear.

Spray Drift Management

When this product is used in "commercial sod production", the following Best Management Practices for reducing spray drift apply.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

PRODUCT DESCRIPTION:

Designed for turfgrass applications, EH-1466 contains four active ingredients:

- (1) Triclopyr provides broad-spectrum weed control for some of the tough broadleaf weeds such as wild violet, ground ivy, oxalis and wild blackberry.
- (2) Sulfentrazone causes rapid desiccation and yellowing of the plant tissue on emerged, susceptible weeds. Sulfentrazone is in the aryl triazolinone family and inhibits protoporphyrinogen oxidase (Protox), a pivotal enzyme in chlorophyll production. Without this key enzyme, a build-up of peroxide-like compounds occurs, thus causing the plant cell membranes of weeds to rupture. Sulfentrazone provides post emergent weed control for common weed species in turfgrass such as spurge and thistles and suppression of yellow nutsedge.
- (3) 2,4-D is an auxin-type herbicide, which is a class of plant growth regulators. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
- (4) Dicamba is absorbed through the leaves and roots and has multiples modes of actions for hard-to-kill broadleaf weeds.

Combining these herbicides provides a very wide spectrum of weed control for tough and susceptible weeds.

EH 1466 Herbicide controls weeds by affecting multiple sites within the broadleaf weeds. The symptoms of susceptible broadleaf weeds include leaf and stem curl or twisting, and weed yellowing.

EH-1466 Herbicide offers these advantages:

- Excellent postemergent activity with proven performance for some of the toughest broadleaf weeds in turfgrass.
- This product exhibits improved cool-weather performance.
- Sulfentrazone combinations provide rapid and effective weed control for common and troublesome (tough) weed species in turfgrass, including: dandelion, spurge and white clover.
- The speed of action (rate of weed phytotoxicity [yellowing]) and the early weed symptoms are features of sulfentrazone. Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 10 to 14 days.
- Triclopyr combinations broaden the weed control spectrum to include many woody and hard-to-control species.
- This product is generally rainfast in as little as 3 hours.

SPRAY PREPARATION AND TANK MIXTURES:

EH 1466 Herbicide is an emulsifiable concentrate that disperses easily in water and forms a stable emulsion. In certain applications, liquid fertilizer may replace part of the water.

Mixing with water:

Add one-half the required amount of water to the spray tank, then add EH-1466 Broadleaf Herbicide slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying.

When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Do not use tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer the spray solution to alter the pH range as appropriate.

Mixing with liquid fertilizers:

Use suitable sources and rates of fertilizer based upon recommendations of your fertilizer supplier or State Extension Service Specialist.

Verify physical compatibility with a jar test: Always perform a jar test for compatibility before large scale mixing. The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids) or suspensions. Physical compatibility of this product is adequate with liquid nitrogen solutions. Mixing this product with suspensions or N-P-K solutions may not be satisfactory (may be marginal) without pre-mixing this product with water. Premixing this product with 2 parts water will ensure that the emulsifiers are activated enabling the herbicide to be suspended in the fertilizer.

Adjuvants and spray additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then adjuvant(s) combined with EH-1466 Broadleaf Herbicide would not be recommended. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury.

GROUND EQUIPMENT:

Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 10 to 220 gallons per acre with spray pressures adjusted to between 20 to 40 psi. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essential when using this product.
- Over-application or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.

Hand operated sprayers including backpack sprayers, compression sprayers are appropriate for small turfgrass.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

WHERE TO USE:

This product provides broadleaf weed control in the following sites:

- **Ornamental Turfgrass sites:**
 - **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
 - **Ornamental Turf sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
 - **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
- **Non-cropland sites:** include Highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations.
- **Agricultural site:** Commercial sod production

Prohibitions of Sites:

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.

Prohibitions:

- Do not apply this product to St Augustinegrass, bentgrass, greens and tees, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when temperatures are above 85°F, some injury may be expected with spot treatments when air temperatures exceed 85°F.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Injury can occur if this product

9/14

is applied under any of these or other stress conditions. Under any of these stress conditions, any turf damage caused by the use of this product is beyond the control of PBI/Gordon Corporation and all risk is assumed by the buyer and/or user.

- For ground application only; aerial applications are not permitted.

STATE RESTRICTIONS:

- Arizona: The state of Arizona has not approved this product for use on sod farms.

APPLICATION SCHEDULES:

Apply this product to broadleaf weeds that are young and actively growing for the best results. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective. Fall applications provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

For the Listed Residential/domestic sites, Ornamental Turf sites, Institutional sites and Agricultural sites:

Do not apply more than two (2) broadcast treatments of this product per site per year. A second broadcast application or a follow-up application as a spot treatment is recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

For the Listed Non-cropland sites:

Use only one (a single) broadcast treatment of this product per site per year for woody plant infestations, or two broadcast treatments for annual and perennial weeds. It is required to wait 30 days between treatments and a spot treatment may be substituted for the second broadcast treatment if necessary.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to this product.

For Newly Seeded Areas:

Delay application of this product to grass seedlings until after the second or third mowing.

For Newly Sodded, Sprigged, or Plugged Areas:

The application of this product should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Reseeding interval:

Treated areas may be reseeded 3 weeks after application.

Turfgrass tolerance:

- Turfgrass tolerance to this product may vary, and temporary turfgrass yellowing may occur on listed warm-season turfgrass (see Table 1).
- Tolerant turf species listed on this label may exhibit temporary turf injury. The best tolerance occurs under optimal conditions for the turfgrass. Adverse environmental conditions may reduce the selectivity on the turfgrass. Injury may occur under marginal conditions (e.g. low temperatures and drought stress) or under extreme conditions (e.g. high temperatures and high humidity). To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Under any of these stress conditions, to the extent

consistent with applicable law, any turf damage caused by the use of this product is beyond the control of PBI/Gordon Corporation and all risk is assumed by the buyer and/or user.

- Certain spray tank additives (adjuvants, wetting agents, and surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Use adjuvants and spray additives or tank-mix combinations only when your experience indicates that the tank mixture will not result in objectionable turf injury
- Do not broadcast apply this product when temperatures are above 85°F; some injury can also be expected with spot treatments when air temperatures exceed 85°F.

**HOW MUCH TO USE:
USE RATES AND SPRAY VOLUMES:**

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

TABLE 1. USE RATES FOR ORNAMENTAL TURFGRASS, SOD FARMS, AND NONCROPLAND.		
Species	Rate	Recommended Spray Volume
Cool-season Turf		
Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues	3.25 to 4 Pints/Acre (1.2 to 1.5 fl.oz./1,000 sq.ft.)	10 to 220 Gallons/Acre (0.25 to 5.0 Gallons/1,000 sq.ft.)
Warm-season Turf (OPTIONAL)		
(OPTIONAL) Hybrid Bermudagrass, common Bermudagrass, zoysiagrass, and bahiagrass	2 to 2.25 Pints/Acre (0.75 to 0.83 fl.oz./1,000 sq.ft.)	10 to 220 Gallons/Acre (0.25 to 5.0 Gallons/1,000 sq.ft.)
<p>(OPTIONAL) Note: Do not apply to above listed warm-season turfgrass unless turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass. We suggest testing this product on a small area and observe the treated area for 30 days (during normal growing conditions) to determine the acceptability of turf discoloration.</p> <p>Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy.</p> <p>Dormant turf: This product may be applied to fully dormant bermudagrass, fully dormant zoysiagrass and fully dormant bahiagrass.</p>		

TABLE 2 . Limitations on broadcast applications to ornamental turfgrass, sod farms and noncropland.				
Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum seasonal rate
Ornamental turfgrass	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb. triclopyr ae/A) (0.10 lb dicamba ae/A)	2	NA	8.0 pints/A (1.75 lbs 2,4-D ae/A) (0.50 lb. triclopyr ae/A) (0.20 lb dicamba ae/A)
	Sod farms			4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb. triclopyr ae/A) (0.10 lb dicamba ae/A)
Noncropland	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb. triclopyr ae/A) (0.10 lb dicamba ae/A)	1 (woody plants)	NA	8.0 pints/A (1.75 lbs 2,4-D ae/A) (0.50 lb. triclopyr ae/A) (0.20 lb dicamba ae/A)
		2 (annual & perennial weeds)	30 days	

11/14

**SPOT TREATMENT:
WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS AND PUMP-UP TYPE
SPRAYERS):**

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.
- Follow-up applications as spot treatments at a 30 day interval are recommended for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **For cool-season turfgrass listed in Table 1:** Mix 1.2 to 1.5 fl.oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **[OPTIONAL TEXT] For warm-season turfgrass listed in Table 1:** Mix 0.75 to 0.83 fl.oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

CULTURAL TIPS:

Irrigation:

- Do not apply this product through any type of irrigation system.
- Rainfast in as little as 3 hours. Do not apply this product immediately before rainfall or irrigation.
- If possible, do not irrigate or water the turfgrass within 3 to 24 hours after application.
- If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product.

BROADLEAF WEEDS CONTROLLED:

EH-1466 Herbicide will control or suppress the following broadleaf. Apply any time the emerged broadleaf weeds are susceptible.

BROADLEAF WEEDS			
Aster, white heath & white prairie	English Daisy	Lespedeza, common	Ragweed
Bedstraw	False dandelion (*spotted catsear & common catsear)	Mallow, common	Red sorrel (*sheep sorrel)
Beggarweed, creeping		Matchweed	Shepherdspurse
Bindweed	Field bindweed	Mouseear chickweed	Speedwell (Veronica)
Black medic	(*morningglory & creeping jenny)	Mustard	Spurge
Broadleaf plantain		Nettle	Thistle
Buckhorn plantain	Field oxeye-daisy	Nutsedge** (yellow)	Virginia buttonweed
Bull thistle	(*creeping oxeye)	Old world diamond flower	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Burdock, common	Filaree, whitestem & redstem	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Wild carrot
Buttercup, creeping		Parsley-piert	Wild garlic
Carpetweed	Florida betony	Pennsylvania smartweed	Wild geranium
Catnip	Florida pusley	Pepperweed	Wild lettuce
Chickweed	Ground ivy	Pigweed	Wild mustard
Chicory	Groundsel	Pineappleweed	Wild onion
Cinquefoil	Hawkweed	Plantain	Wild strawberry
Clover	Healall	Poison ivy	Wild violet***
Curly dock	Henbit	Poison oak	Yarrow
Dandelion	Innocence (Blue-eyed Mary)	Prickly lettuce (*compass plant)	Yellow rocket
Dayflower		Puncturevine	
Deadnettle	Knotweed	Purple cudweed	
Dock	Lambsquarters	Purslane	and many more broadleaf weeds
Dogfennel	Lawn burweed		

* Synonyms
 ** Suppression only when nutsedge is young and actively growing.
 *** For best results, apply in the spring when wild violets are blooming or apply a late fall application followed by a spring application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. For Plastic Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under

13/14

abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Tough Weed Control
- Tough Weed LOGO
- Targets [tough], [triclopyr] weeds
- Contains: triclopyr + sulfentrazone + 2,4-D + dicamba
- To control some of your toughest [triclopyr] weeds
- Consistently fast control on tough weeds – wild violet, ground ivy, oxalis, thistles, dandelion, spurge, clover, plantain and [various other listed weeds].
- Oil-based
- Oil-based formula
- Oil-based formulation
- Rainfast in as little as 3 hours
- Rainfast [Rainproof] in 3 hours
- [weed name inserted] control in as little as one week
- Proven performance

- Cool-weather weed control option
- From the makers of Trimec® herbicides.
- Trimec® is a registered trademark of PBI/Gordon Corporation.
- For information call XXX-XXX-XXXX [contact www.xxx-xxxx.com]
- ProForm® logo presented on the containers



- For use on: Bluegrass, Fescues, Perennial Ryegrass, [Bermudagrass, Zoysiagrass]
- Fast visual response
- Weed control — FAST!
- Suitable for applications up to 85°F
- Foliar absorption
- Reduces call backs
- Low odor
- Low-odor formulation
- Designed for residential, commercial turf [and golf] applications and many other sites.

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00XXX.20081223.EH1466proposedcorrected.doc
2. **Reason for Issue:** Corrected ingredient statement