

2217-930

8/9/2012

1/16



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

James L Kunstman Ph D
PBI/Gordon Corporation
1217 12th Street
P O Box 014090
Kansas City MO 64101

AUG - 9 2012

Subject Label Notification per PRN 98 10
EH 1471 Herbicide
EPA Registration Number 2217 930
Application Dated July 27 2012

Dear Dr Kunstman

The Agency is in receipt of your application for Pesticide Notification under Pesticide Registration Notice (PRN) 98 10 for EH 1471 Herbicide (EPA Reg No 2217 930) dated July 27 2012 The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98 10 and that the action requested falls within the scope of PRN 98 10 The label submitted with the application has been stamped Notification and will be placed in our records

If you have any questions please contact Emily Hartman via email at hartman.emily@epa.gov or by phone at (703) 347 0189

Sincerely

A handwritten signature in black ink, appearing to read "Kathryn V Montague".


Kathryn V Montague Project Manager 23
Herbicide Division
Registration Division
Office of Pesticide Programs

Handwritten initials "for" in black ink, positioned to the right of the typed name.

2/16

Please read instructions on reverse before completing form

Form Approved 616 B No. 2070 0060 Approval Expires 2 28 95

	United States Environmental Protection Agency Washington DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	--	---	-----------------------

Application for Pesticide Section I

1 Company/Product Number <p style="text-align: center;">2217 930</p>	2 EPA Product Manager <p style="text-align: center;">Kathryn V Montague</p>	3 Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4 Company/Product (Name) <p style="text-align: center;">EH 1471 Herbicide</p>	PM# <p style="text-align: center;">Product Manager—Team 23</p>	
5 Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City Missouri 64101 <input type="checkbox"/> Check if this is a new address	6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(i) my product is similar or identical in composition and labeling to EPA Reg No _____ NOTIFICATION Product Name _____ AUG— 9 2012	

Section – II

<input type="checkbox"/> Amendment Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

Explanation Use additional page(s) if necessary (For section I and Section II)

Labeling notification per PRN 98 10 Notifications Non notifications and Minor Formulation Amendments Section II (N) Other revisions

e mail to jkunstman@pbigordon.com FAX 816-421 2731

Section – III

1 Material This Product Will Be Packaged In				2 Type of Container	
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)		
Certification must be submitted		If Yes No per Unit Packaging wgt container	If Yes No per Package wgt Container		
3 Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4 Size(s) Retail Container 1 and 2.5 gallons		5 Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6 Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section – IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)					
Name James L. Kunstman Ph D		Title Director of Regulatory Services		Telephone No (Include Area Code) 816 460 6202	
Certification I certify that the statements I have made on this form and all attachments thereto are true accurate and complete I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law					6 Date Applicant Received (Stamped)
2 Signature 			3 Title Director of Regulatory Services		
4 Typed Name James L. Kunstman Ph D			5 Date July 27 2012		

3/16



1217 W 12th STREET P O BOX 014090
KANSAS CITY MISSOURI 64101 0090
816 421 4070 1 800 821 7925
FAX 816 474 0462

July 27 2012

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U S Environmental Protection Agency
Room S 4900 One Potomac Yard
2777 South Crystal Drive
Arlington VA 22202 4501
Attn Kathryn V Montague PM 23

Dear Ms Montague

Subject Labeling notification per PRN 98 10 Notifications Non notifications and Minor Formulation Amendments Section II (N) Other revisions

EH 1471 Herbicide (EPA Reg No 2217 930)

***I* Labeling notification per PRN 98 10**

We ask to modify an existing advertising claim Please refer to page 13 of the labeling

***II* Certification**

This notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152 46 and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to the EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA

Enclosures

- 1 Application for Pesticide Amendment (EPA Form 8570 1)
- 2 One (1) annotated version of the draft labeling and
- 3 One (1) version of the draft labeling without notations
- 4 One (1) CD with certification

If you have any questions please call me at 816-460 6292 or contact me at jkunstman@pbigordon.com

Sincerely

James L. Kunstman Ph D
Director of Regulatory Services

4/16

EH-1471 HERBICIDE

NOTIFICATION

AUG - 9 2012

EPA Reg No 2217-930

Alternate Brand Name Q4 Plus Turf Herbicide for Grassy & Broadleaf Weeds

ACTIVE INGREDIENT

Quinclorac	8.43%
Sulfentrazone	0.69%
2,4-D dimethylamine salt	11.81%
Dicamba dimethylamine salt	1.49%

INERT INGREDIENTS

TOTAL	<u>77.58%</u>
	100.00%

THIS PRODUCT CONTAINS

- 0.75 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 8.43%
- 0.06 lb N-[2,4-dichloro-5-(4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl)phenyl]methanesulfonamide per gallon or 0.69%
- 0.88 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 9.81%
- 0.10 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 1.24%

Isomer Specific By AOAC Methods

CAS Registry Numbers Quinclorac (84087-01-4) Sulfentrazone (122836-35-5) Dicamba dimethylamine salt (2300-66-5) 2,4-D dimethylamine salt (2008-39-1)

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si Usted no entiende la etiqueta busque a alguien para que se la explique a Usted en detalle (If you do not understand the label find some one to explain it to you in detail)

STOP! READ THE ENTIRE LABEL FIRST OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION Causes moderate eye irritation Avoid contact with eyes or clothing Harmful if swallowed

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are natural rubber natural rubber blends and laminates If you want more options follow the instructions for category A on an EPA chemical resistance category selection chart

All mixers loaders applicators and other handlers must wear

- long sleeved shirt and long pants
- protective eyewear (such as goggles face shield or safety glasses)
- shoes and socks
- chemical resistant gloves (except for applicators using ground boom equipment) and



chemical resistant apron when mixing or loading cleaning up spills or equipment or otherwise exposed to the concentrate

User Safety Requirements

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

User Safety Recommendations	
Users should wash hands thoroughly with soap and water before eating drinking chewing gum using tobacco or using the toilet	
<ul style="list-style-type: none"> • Users should remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing If pesticide gets on skin wash immediately with soap and water 	
Users should remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing	

First Aid	
If in eyes	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice
If swallowed	Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by a poison control center or doctor Do not give anything by mouth to an unconscious person
If on skin or on clothing	Take off contaminated clothing Rinse skin immediately with plenty of water for 15 20 minutes Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact 1 888 800 5556 for emergency medical treatment information	

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas Do not contaminate water when disposing of equipment wash waters or rinsate

This chemical has properties and characteristics associated with chemicals detected in groundwater The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination Application around a cistern or well may result in contamination of drinking water or groundwater

Physical and Chemical Hazards

Combustible Do not use or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application

For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170

This standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- coveralls worn over short sleeved shirt and short pants
- chemical resistant footwear plus socks
- chemical resistant gloves made of any water proof material and
- protective eyewear

Non Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Reentry Statement Do not enter or allow people (or pets) to enter the treated area until sprays have dried

1 Product Description

4 in 1 Herbicide

EH 1471 Herbicide contains 4 active ingredients including quinclorac and sulfentrazone that broaden the spectrum of weed control EH 1471 Herbicide is a versatile herbicide that will control many troublesome grassy and broadleaf weeds in established turfgrass when used as directed

4 in 1 is better than 3 in 1

- Versatile post emergent herbicide for turfgrass to control both grassy and broadleaf weeds
- Quinclorac is absorbed by foliage and roots and translocated throughout the plant where it uses multiple modes of action to control both grassy and broadleaf weeds One mode of action involves an auxin type herbicidal activity which disrupts plant growth in broadleaf weeds while another mode of action works as an enzyme/cell wall disruptor to control crabgrass and other grassy weeds
- The effects on grassy weeds include stunting gradual reddening yellowing and followed by tissue death Symptoms typically appear 7 to 14 days after application Sulfentrazone provides post emergent weed control for common weed species in turfgrasses such as spurge Foliar contact of sulfentrazone on emerged susceptible weeds results in rapid desiccation and plant tissue death
- Wide window of application for broadleaf weeds beginning at emergence and active weed growth
- The symptoms of susceptible broadleaf weeds resemble those of 2 4 D and include leaf and stem curl or twisting and tissue death

2 Where To Use

This product provides broadleaf control in warm season and cool season turfgrass in 5 use sites. See tables 2 and 3 below for turfgrass species that can be treated with EH 1471 Herbicide

- **Residential/domestic sites** are defined as areas associated with the household or home life including but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes
- **Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations including but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings

Non cropland sites include Highway rights of way (principal interstate, county, private, and unpaved roads), roadsides, roadside ditches, road shoulders, road embankments, dividers and medians, municipal, state and federal lands, airports and military installations

Agricultural site Commercial sod production

Prohibitions of Sites

Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays)

- Do not apply to wetlands (swamps, bogs, potholes, or marshes)
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals. Do not apply to agricultural drainage water or on agricultural ditchbanks

- **DO NOT** apply EH 1471 Herbicide to the following

- Bahiagrass
- Bentgrass
- Carpetgrass
- Centipedegrass
- Dichondra
- St. Augustinegrass
- Greens and collars at golf courses
- Lawns with desirable clovers or legumes
- Ornamentals (flowers, trees, groundcovers, landscape beds and shrubs)

Turfgrass tolerance

- Turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur on fine fescues [hybrid] Bermudagrass, and zoysiagrass (see specific Bermudagrass and zoysiagrass sections). Apply only to fine fescue when it is part of a mixed stand of cool season turf/cool season blends

- Tolerant turf species listed on this label may exhibit temporary turf injury. The best tolerance occurs under optimal conditions for the turfgrass.

Adverse environmental conditions may reduce the selectivity on the turfgrass. Injury may occur under marginal conditions (e.g. low temperatures and drought stress) or under extreme conditions (e.g. high temperatures and high humidity). To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Under any of these stress conditions, any turf damage caused by the use of this product is beyond the control of PBI/Gordon Corporation and all risk is assumed by the buyer and/or user.

- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Use adjuvants and spray additives or tank mix combinations only when your experience indicates that the tank mixture will not result in objectionable turf injury. [See SPRAY PREPARATION Additions of adjuvants (and/or Tank Mixtures) to improve Grassy Weed Control section of this label.]

8/16

- Do not broadcast apply this product when temperatures are above 90 F some injury can also be expected with spot treatments when air temperatures exceed 90 F

3 Cultural Tips

For newly seeded areas

Delay application of this product to grass seedlings until after the second or third mowing or 28 days after emergence

For newly sodded sprigged or plugged areas

The application of this product to newly sodded sprigged or plugged grasses should be delayed until 3 to 4 weeks after the sodding sprigging or plugging operations

Seeding

Turf species listed on this label can be seeded into the treated areas at [4 3 2 1] weeks after the application of this product

Irrigation

Do not apply this product immediately before rainfall or irrigation For best results do not irrigate or water the turfgrass within 24 hours after application If dry conditions exist a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended If rainfall does not occur in 2 to 7 days after application irrigation of at least one half inch is recommended

Mowing

Delay mowing 2 days before and until 2 days after the application of this product Additional stress from low mowing heights may increase the possibility of turf injury Clippings from the first three mowings should be left on the treated area Do not use these clippings as mulch or compost around flowers ornamentals trees or in vegetable gardens

4 Spray Preparation

Mixing with Water

Add one half the required amount of water to the spray tank then add EH 1471 Herbicide slowly with agitation and complete filling the tank with water Mix thoroughly and continue agitation while spraying

When this product is left standing for extended periods of time re agitate to assure uniformity of the spray mixture

Mixing with Liquid fertilizers

In certain applications liquid fertilizer may replace part of the water as a diluent

ALWAYS PREMIX EH 1471 Herbicide WITH WATER BEFORE ADDING TO FLUID FERTILIZERS For liquid nitrogen solutions such as UAN or urea solutions use a premix of 1 part of this product with 4 parts of water or use a premix with a 1 4 ratio of product to water For other fluid fertilizers such as suspensions use a premix of 1 part of this product with 50 to 60 parts of water

Use suitable sources and rates of fertilizer based upon local recommendations Refer to the mixing directions on the labels of the liquid fertilizers Always perform a jar test for compatibility before large scale mixing

The jar test can be conducted by mixing all components in a small container in proportionate quantities If the mixture separates after standing and can be mixed readily by shaking then the mixture can be used and applied with spray equipment providing continuous agitation If large flakes sludge gels or other precipitates form or if a separate oily layer or oil globules appear then the herbicide and the liquid fertilizer should not be prepared as a tank mixture

[Begin optional text]

Additions of Adjuvants (and/or Tank Mixtures) to Improve Grassy Weed Control

Methylated seed oil is a suitable adjuvant for post emergent applications and may cause slight yellowing to desirable turf. The use of nitrogen fertilizer or chelated iron (such as FeRRAMEC® Plus MICROS) can reduce the slight yellowing that may occur on some turfgrass species. Methylated seed oil used must meet the following criteria:

- Be non phytotoxic
- Contain only EPA approved ingredients
- Provide good mixing quality in the jar test and
- Be used only when your experience indicates that this tank mixture will not result in objectionable turfgrass injury

Do not include additives when tank mixing with oil based products or EC (Emulsifiable Concentrate) products as this may cause phytotoxicity (yellowing) of desirable turfgrass. Other additives or adjuvants may be used however certain additives may reduce the selectivity on the turfgrass. Under some environmental conditions some of these products may cause phytotoxicity (yellowing) of desirable turfgrass. Use these adjuvants, spray additives or tank mix combinations only when your experience indicates that the tank mixture will not result in objectionable turf injury.

[End optional text]

5 Spray Equipment

Ground equipment Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate flat fan nozzles, tips and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Hand operated sprayers including backpack sprayers and compression sprayers are appropriate for small turfgrass areas. Calibration and proper application are essential when using this product.

Cleaning spray equipment Clean sprayer before and after using this product. Use soap, household ammonia, detergent and water, or an approved spray tank cleaner and rinse thoroughly. Cross contamination may cause physical incompatibility (mixing problems) or result in turf injury.

Spray distribution

- The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator.
- Uniform applications are essential when using this product. Over application, excessive overlaps or rates above those specified on this label can cause turf injury.
- Avoid spray overlaps with hand held equipment. Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Spray wands fitted with flat fan nozzles should not be waved in a back and forth motion, or in a side to side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side to side motion results in uneven coverage. To avoid excessive spray pattern overlaps, a spray colorant may be used. This product may cause injury to susceptible/non target plants at the use site by contacting the foliage, stems, or roots. To prevent injury to susceptible crops and other desirable broadleaf plants including but not limited to cotton, legumes, tobacco, tomatoes, garden/vegetable crops and ornamentals (flowers, trees, and shrubs) avoid contact with the spray solution, spray droplets, and spray mist (fine droplets).

Chemigation Do not apply this product through any type of irrigation system.

Aerial application Do not apply as an aerial application.

6 Spray Drift Management

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including but not limited to bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use, or consumption. Susceptible crops include but are not limited to cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Additional requirements for ground boom application: Do not apply with a nozzle height greater than 4 feet above the turf canopy.

7 Post Emergent Control of Grassy Weeds

EH 1471 Herbicide can provide post emergent control and suppression of the grassy weeds listed in Table 1. Apply to weeds during the growth stages as shown in Table 1. Applications under adequate soil moisture conditions are preferred. Early summer treatments are generally more effective. Applications in the summer (approximately July 15 to August 15) to older, drought stressed grassy weeds are less effective. Late summer applications (after August 15) to mature crabgrass can be very effective. Weed control is affected by the spray volume, timing, and the weed growth stages (see Tables 1, 2, and 3).

TABLE 1 Leaf stages of grassy weeds and yellow nutsedge for optimum control

Weed species	1 to 3 leaf	4 to 5 leaf (1 tiller)	6 leaf (2 tillers)	7 leaf (3 tillers)	8 leaf (4 tillers)	Mature (late season)
Crabgrass (large and smooth) Barnyardgrass Foxtail (green yellow and giant) Signalgrass (broadleaf)	X	X	X	Footnote 1	Footnote 1 (reduced control)	X
Nutsedge (yellow)	X	Footnote 1	Footnote 1	Footnote 1	Footnote 1 (reduced control)	X

¹ Second or follow up applications on ornamental turfgrass may be required. Early summer treatments are generally more effective. Applications in the summer (approximately July 15 to August 15) to older drought stressed grassy weeds are less effective. Late summer applications (after August 15) to mature crabgrass can be very effective.

8 Post Emergent Control of Broadleaf Weeds

Apply this product to broadleaf weeds that are young and actively growing for the best results. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older drought stressed weeds are less effective. Fall applications provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

9 Applications

The maximum number of broadcast applications is limited to 2 per year. Spot treatments during the spring and summer are suitable for sparse infestations or as a follow up treatment to a broadcast application on an as needed basis. Second or follow up applications as either broadcast or spot treatments should be made after the initial application on ornamental turfgrass and are recommended for more mature weeds, for dense infestations and for adverse environmental conditions.

Other situations that may need two broadcast or follow up treatments include the following:

- Under certain conditions, applications of this product at the 3 to 4 tiller stage of the annual grasses may not provide complete control.
- All weed grasses do not germinate at the same time. The period of germination for crabgrass and annual grasses can extend into the summer after the initial application of this product and results may be poor and erratic.
- Dense infestations of weeds may prevent thorough spray coverage of the target weeds. Biotypes of large and smooth crabgrass in California have shown varied response to quinclorac. If control failure occurs following a second application, do not reapply this product. Change to a herbicide with a different mode of action [such as [product name]].

Extremes in environmental conditions, i.e. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. Under dry conditions, the expression of herbicide symptoms is generally delayed, and weeds hardened off by drought may be less susceptible to this product.

If objectionable turf injury occurs with the first application, then avoid making the second application of this product until the turfgrass recovery is complete.

Do not broadcast apply this product when temperatures are above 90 F. Temporary turfgrass discoloration can also be expected with spot treatments when air temperatures exceed 90 F.

State Restrictions

Arizona Do not use this product on sod farms in Arizona.

California Make broadcast applications only between March 1 and September 1. If troublesome weeds appear during other times of the year, a spot application can be made. While irrigation is necessary and important for plant growth, apply irrigation water efficiently so that no more than 125% of the net irrigation requirement is applied for any irrigation event. Apply efficient irrigations for six months following application of sulfentrazone-containing products. Do not apply product to bare ground.

New York For use in New York State by spot treatment only. Not for sale, distribution, or use in Nassau or Suffolk Counties in New York State.

TABLE 2 BROADCAST TREATMENTS FOR COOL SEASON TURFGRASS	
Application Site	Use Rate per Application
Kentucky Bluegrass Perennial ryegrass Fescues Annual Bluegrass (<i>Poa annua</i>) Rough Bluegrass (<i>Poa trivialis</i>) Annual Ryegrass	7 to 8 pints/A (2.6 to 3.0 fl oz /1000 sq ft)
Spray Volume For Conventional Spray Equipment Use 50 to 220 gal/A (1.2 to 5.0 gal/1000 sq ft)	
For Low Volume Spray Equipment [such as PermaGreen Equipment and backpack sprayers] Equipment should be calibrated to apply at least 20 gallons per acre (0.45 gallons/1,000 sq ft). Use this lower spray volume (0.45 gallons/1,000 sq ft) only when your experience indicates that this volume provides effective weed coverage, adequate weed control, acceptable turf safety/tolerance, and will not result in objectionable turf injury.	
Note Use the higher spray volumes (more than 50 gpa) for dense weed populations.	

TABLE 3 BROADCAST TREATMENTS FOR BERMUDAGRASS AND ZOYSIAGRASS	
Application Site	Use Rate per Application
Bermudagrass (common and hybrid) Zoysiagrass	5 to 7 pints/A (1.8 to 2.6 fl oz /1000 sq ft)
Spray Volume For Conventional Spray Equipment Use 50 to 220 gal/A (1.2 to 5.0 gal/1000 sq ft)	
For Low Volume Spray Equipment [such as PermaGreen Equipment and backpack sprayers] Equipment should be calibrated to apply at least 20 gallons per acre (0.45 gallons/1,000 sq ft). Use this lower spray volume (0.45 gallons/1,000 sq ft) only when your experience indicates that this volume provides effective weed coverage, adequate weed control, acceptable turf safety/tolerance, and will not result in objectionable turf injury.	
<p>Apply only when Bermudagrass and zoysiagrass is actively growing</p> <ul style="list-style-type: none"> • Expect temporary discoloration • Some Bermudagrass hybrids and zoysiagrass cultivars are moderately tolerant to this product and may be more susceptible to discoloration <p>For Bermudagrass hybrids, use lower rates until tolerance to injury can be determined.</p> <p>It is impossible to test all environmental conditions and all Bermudagrass hybrids or zoysiagrass cultivars. We suggest testing this product on a small area and observe the treated area for 30 days (during normal growing conditions) to determine the acceptability of turf discoloration.</p> <ul style="list-style-type: none"> • Some stunting of the Bermudagrass or zoysiagrass should be expected and turf generally recovers in 7 to 21 days <p>Do not apply in the fall during fall to winter transition period Do not apply in the spring during winter to spring transition period</p> <ul style="list-style-type: none"> • To avoid turf injury, use only on Bermudagrass and zoysiagrass that is not under stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides • For optimum results <ul style="list-style-type: none"> • Irrigate 24 hours before and 24 hours after application with 1/2 inch of water • The addition of nitrogen fertilizer or chelated iron [such as FerROMECS® Plus MICROS] may reduce some turf discoloration • Spray in the morning hours and avoid application during extreme hot or dry conditions • Equipment calibration is essential and avoid spray overlaps • Do not use higher pressure equipment; spray pressure should be 40 psi or lower 	

SPOT TREATMENTS

Calibration and proper application are essential when using this product. Spray coverage should be uniform and complete.

Over applications can result in turfgrass injury.

Cool Season Turf listed in Table 2 Mix 2.6 to 3.0 fl oz of this product with 1.0 gallons of water for treatment of approximately 1,000 sq ft of turfgrass. Apply to weeds during the growth stages as shown in Table 1.

Warm Season See specific Bermudagrass and zoysiagrass directions in Table 3. Mix 1.84 to 2.57 fl oz of this product with 1.0 gallons of water for treatment of approximately 1,000 sq ft of turfgrass. Apply to weeds during the growth stages as shown in Table 1.

Limitations for use on Ornamental Turf and Non Cropland areas

The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum application rate is 8 pints of product per acre per application [0.88 lb 2,4-D ae and 0.10 lb dicamba ae/A/application]. The maximum seasonal rate is 16 pints of product per acre per year [1.76 lb 2,4-D ae and 0.20 lb dicamba ae/A/year].

Limitation for use on Sod Farms

The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum application rate is 8 pints of product per acre per application [0.88 lb 2,4-D ae and 0.10 lb dicamba ae/A/application]. The maximum seasonal rate is 16 pints of product per acre per year [1.76 lb 2,4-D ae and 0.20 lb dicamba ae/A/year].

10 Weeds Controlled

EH 1471 Herbicide will control or suppress the following. Apply when weeds are young and actively growing.

WEEDS			
Aster white heath & white prairie	Dollarweed (pennywort) [Doveweed]	Mallow common	Shepherdspurse
Barnyardgrass	English Daisy ¹	Matchweed	Signalgrass ¹
Bedstraw	False dandelion (spotted catsear & common catsear)	Morningglory spp	(Broadleaf)
Beggarweed creeping	Field bindweed	Mouseear chickweed	Speedwell Veronica
Bindweed	(morningglory & creeping jenny)	Nutsedge yellow	(Common Slender and Thymeleaf)
Black medic	Field oxeye daisy (creeping oxeye)	Old world diamond flower	Spurge
Broadleaf plantain	Filaree whitestem & redstem	Oxalis (yellow woodsorrel & creeping woodsorrel)	Thistles
Buckhorn plantain	Florida betony	Parsley piert	Virginia buttonweed ¹
Bull thistle	Florida pusley	Pennsylvania smartweed	White clover (Dutch clover honeysuckle clover white trefoil & purplewort)
Burdock common	Foxtail ¹	Pepperweed	Wild carrot
Buttercup creeping	(green yellow and giant)	Pigweed	Wild garlic
Carpetweed	Ground ivy	Pineappleweed	Wild geranium
Chickweed common	Groundsel	Plantain	Wild lettuce
Chicory	Geranium (Carolina)	Poison ivy	Wild mustard
Cinquefoil	Hawkweed	Poison oak	Wild onion
Clovers	Healall	Prickly lettuce (compass plant)	Wild strawberry
Crabgrass ^{1 2} (large and smooth)	Henbit	Puncturevine	Wild Violet ¹
Curly dock	Innocence (Blue eyed Mary)	Purple cudweed	Yarrow
Dandelion	Knotweed	Purslane	Yellow rocket
Dayflower	Lambsquarters	Ragweed	and many other weeds
Deadnettle	Lawn burweed	Redweed	
Dock	Lespedeza common	Red sorrel (sheep sorrel)	
Dogfennel			
Synonyms			
¹ Follow up application may be required			
² Biotypes of large and smooth crabgrass in California have shown varied response to quinclorac. If control failure occurs following a sequential (or follow up) application, do not reapply this product. Change to a herbicide with a different mode of action [such as [product name]].			

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

PESTICIDE STORAGE Store in original container in a locked storage area inaccessible to children or pets Keep from freezing

PESTICIDE DISPOSAL Pesticide wastes are acutely hazardous Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons]

CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container Offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times

[Pressure rinse as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds Drain for 10 seconds after the flow begins to drip]

[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons]

CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container Offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times

[Pressure rinse as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds Drain for 10 seconds after the flow begins to drip]

[For Refillable Containers]

CONTAINER HANDLING Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose

Container cleaning Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller

To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product By opening and using this product buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception If the terms are not acceptable return this product unopened immediately to the point of purchase and the purchase price will be refunded in full

It is impossible to eliminate all risks inherently associated with use of this product Damage to the treated article ineffectiveness or other unintended consequences can result from use of the product under abnormal conditions such as weather presence of other materials or the manner or use of application etc Such factors and conditions are beyond the control of the manufacturer and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions warnings and cautions

The manufacturer warrants only that this product conforms to the chemical description given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use subject to the inherent risks described below **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE HEREBY EXPRESSLY DISCLAIMED**

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER FOR ANY AND ALL LOSSES DAMAGES OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT WHETHER OR NOT BASED IN CONTRACT NEGLIGENCE STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED AT THE MANUFACTURER'S OPTION TO REPLACEMENT OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THE PRODUCT The Manufacturer must be promptly notified in writing of any claims whether based in contract tort negligence strict liability or otherwise to be eligible to receive either remedy stated above

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this LIMITED WARRANTY AND DISCLAIMER in any manner

APPENDIX

1 Statements which may appear on different label components depending on packaging configuration

- See next panel for additional Precautionary Statements and First Aid
- Net Contents _____
- EPA Est No _____

2 Advertising claims that may be presented on container labeling advertisements brochures and other marketing/sales promotional materials

- Water based formula
Water based formulation
- Proven performance
Consistently fast control on tough weeds like dandelion spurge clover plantain ground ivy and [various other listed weeds]
- Controls grassy and broadleaf weeds
Grassy and broadleaf weed control
- Four In One or 4 in 1
Controls crabgrass barnyardgrass foxtail and signalgrass
From the makers of Trimec® herbicides
Trimec® is a registered trademark of PBI/Gordon Corporation
For information call XXX XXX XXXX [contact [www xxx xxx.com](http://www.xxx.xxx.com)]
- ProForm(TM) logo presented on the containers



- Post emergent control of crabgrass yellow nutsedge plus broadleaf weeds [as listed] – FAST!
- For use on Bluegrass Fescues Perennial Ryegrass and Annual Ryegrass
- Versatile post emergent control of both grassy and broadleaf weeds
Yellow nutsedge control
- Crabgrass and foxtail control
- Wide window of application for broadleaf weeds
- Foliar application
- Protax inhibitor activity for rapid desiccation and death of broadleaf weeds
- Fast visual response
- Suitable for applications up to 90 [degrees] F
Easy To Use Water Based Formulation
Water based
- Rainfast in six [6] hours
Rainfast in as little as six [6] hours
Extra [Xtra] knockout power on grassy weeds
- Extra [Xtra] Power
- Contains 0.75 lbs/gallon of quinclorac
Contains premium ratios of active ingredients
Maximum crabgrass performance
Reseed in [4 3 2 1] weeks
- U S Patent 6 849 579

DOCUMENT CONTROL INFORMATION

1 Unique Label Identifier 002217 00930 20120727 notif proposed highlighted doc

2 Reason for Issue ad claim