

2217-931

7/29/2010

10817



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

James L. Kunstman, Ph.D.
PBI/Gordon Corporation
PO Box 014090
Kansas City, MO 64101

JUL 29 2010

Subject: Notification - Minor Label Revision (Removal of AZ Sod Farm Restriction per PR Notice 98-10).

EH-1473 Herbicide

EPA Reg. No. 2217-931

Your Application, Dated June 9, 2010

Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product.

The Registration Division (RD) has conducted a review of this request for applicability under PRN 98-10 and finds that the label changes requested fall within the scope of the PR Notice. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at (703) 305-1243 or Susan Stanton of my staff at (703) 305-5218.

Sincerely,

A handwritten signature in cursive script that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

| | | | |
|--|---|--|-----------------------|
|  | United States Environmental Protection Agency Washington, DC 20460 | <input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other | OPP Identifier Number |
|--|---|--|-----------------------|

Application for Pesticide - Section I

| | | |
|---|---|--|
| 1. Company/Product Number <p style="text-align: center;">2217-931</p> | 2. EPA Product Manager <p style="text-align: center;">Kathryn V. Montague</p> | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) <p style="text-align: center;">EH-1473 Herbicide</p> | PM# <p style="text-align: center;">Product Manager—Team 23</p> | |
| 5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ | |

Section - II

| | |
|--|--|
| <input type="checkbox"/> Amendment - Explain below. | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - Explain below. |

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Labeling Notification per Pesticide Registration Notice (PRN) 98-10


Please refer to cover letter for details and certification statement.

e-mail to jkunstman@pbigordon.com FAX: 816-421-2731

Section - III

| | | | | | |
|---|--|---|---|--|--------------------------------|
| 1. Material This Product Will Be Packaged In: | | | | | |
| Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | 2. Type of Container | | |
| * Certification must be submitted If "Yes" Unit Packaging wgt. No. per container If "Yes" Package wgt. No. per Container | | | <input checked="" type="checkbox"/> Metal | <input checked="" type="checkbox"/> Plastic | <input type="checkbox"/> Glass |
| | | | <input type="checkbox"/> Paper | <input type="checkbox"/> Other (Specify) _____ | |
| 3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container | | 4. Size(s) Retail Container 2.5, 15, 30 and 55 gallons | | 5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product | |
| 6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | | | | |

Section - IV

| | | |
|--|---|---|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | |
| Name James L. Kunstman, Ph.D. | Title Director of Regulatory Services | Telephone No. (Include Area Code) 816-460-6292 |
| I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | 6. Date Application Received (Stamped) |
| 2. Signature  | 3. Title Director of Regulatory Services | (Stamp area with grid) |
| 4. Typed Name James L. Kunstman, Ph.D. | 5. Date June 9, 2010 | |

4 of 17

EH-1473 HERBICIDE

NOTIFICATION

EPA Reg. No. 2217-931

JUL 29 2010

Alternate Brand Name: Trimec 1000 Low Odor Broadleaf Herbicide

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.

ACTIVE INGREDIENT:

| | |
|---|---------|
| 2,4-D, diethanolamine salt (DEA) | 34.59% |
| 2,4-D, dimethylamine salt (DMA)..... | 1.22% |
| MCPP-p, dimethylamine salt (DMA) | 8.17% |
| Dicamba, dimethylamine salt (DMA) | 2.68% |
| INERT INGREDIENTS: | 53.34% |
| TOTAL | 100.00% |

THIS PRODUCT CONTAINS:

- 2.37 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 24.45%.
- 0.65 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 6.67%.
- 0.22 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 2.23%.

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

WARNING-AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find some one to explain it to you in detail.)

KEEP FROM FREEZING

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals


WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- protective eyewear (goggles, face shield, or safety glasses),
- long-sleeved shirt and long pants,



- shoes and socks,
- chemical-resistant gloves (such as natural rubber, selection category A)
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations

- Users should wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

| | |
|---------------------|--|
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If swallowed | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. |


Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls worn over short-sleeved shirt and short pants,
- chemical-resistant footwear plus socks,
- chemical-resistant gloves made of any water-proof material,
- chemical-resistant headgear for overhead exposure,
- protective eyewear.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

1. Spray Drift Management


A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use only Medium or coarser spray nozzles according to ASAE (S 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.



Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

2. Product Description

Designed for [turfgrass applications] [non-crop] [and IVM (Industrial Vegetation Management)] applications, this product contains four active ingredients.

- (1) 2,4-D in the form of DEA salt, is an auxin-type herbicide, which is a class of plant growth regulators that is low odor and known to control deep rooted perennials. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
- (2) 2,4-D in the form of DMA salt, is an auxin-type herbicide which is a class of plant growth regulators. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
- (3) MCPP-p (mecoprop-p) is an auxin-type herbicide, which is a class of plant growth regulators. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
- (4) Dicamba is absorbed through the leaves and roots and has multiples modes of actions for hard-to-kill broadleaf weeds.

Combining these herbicides provides a very wide spectrum of weed control for susceptible weeds. EH-1473 Herbicide controls weeds by affecting multiple sites within the broadleaf weeds. The symptoms of susceptible broadleaf weeds include leaf and stem curl or twisting, and weed yellowing.


These combined herbicides provide limited residual activity.

EH-1473 Herbicide offers these advantages:

- Excellent postemergent activity with proven performance.
- Low-odor formulation
- The combinations of these active ingredients provide effective weed control for common and troublesome weed species in turfgrass, including: dandelion, spurge and white clover.
- Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 14 to 21 days.

3. Spray Preparation and Tank Mixes

In certain applications, liquid fertilizer may replace part of the water.



Mixing with water:

Add one-half the required amount of water to the spray tank, then add EH-1473 Broadleaf Herbicide slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying. When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Mixing with liquid fertilizers:

Use suitable sources and rates of fertilizer based upon advice of your fertilizer supplier or State Extension Service Specialist.

Verify physical compatibility with a jar test: Always perform a jar test for compatibility before large scale mixing. The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids) or suspensions. Mixing this product with suspensions or N-P-K solutions may not be satisfactory (may be marginal) without pre-mixing this product with water. Premixing this product with 2 to 4 parts water will ensure that the dispersants enable the herbicide to be suspended in the fertilizer.

Adjuvants and spray additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then adjuvant(s) combined with EH-1473 Broadleaf Herbicide would not be recommended. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, [name of registrant] recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

4. Ground Equipment


Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 5 to 220 gallons per acre with spray pressures adjusted to between 20 to 40 psi. Note: For bentgrass (except golf greens) spray volumes, use 145 to 220 gallons per acre. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essential when using this product.
- Over-application or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.

Hand operated sprayers including backpack sprayers, compression sprayers are appropriate for small turfgrass areas.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.



5. Where to Use

This product provides broadleaf control in the following sites.

• **Ornamental Turfgrass sites:**

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental turf sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
- **Non-cropland sites:** including farmyards, fencerows or fence lines, highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians; Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas; Municipal, state, and federal lands: Airports and military installations; railroad rights-of-ways, railroad yards, railroad crossings and railroad bridge abutments; Utility rights-of-way: telephone, pipeline, electrical powerlines, and communication transmission lines.
- **Agricultural site:** Commercial sod production

Prohibitions of Sites:


- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.
- Do not apply to sites being used for commercial timber or seed production, or to other plants being grown for sale or other commercial use.
- Do not apply or allow this product to come into direct contact with cotton, grapes, tobacco, vegetable crops, flowers, fruit or ornamental trees, or other desirable broadleaf plants; small amounts of spray drift may injure susceptible plants, including ornamental trees or shrubs.

Turfgrass tolerance:

- The turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur. Adverse environmental conditions may reduce the selectivity on the turfgrass. Do not apply this product to stressed turf.
- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, [name of registrant] recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

Prohibitions:

- Do not apply this product to St. Augustinegrass, bentgrass greens, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when temperatures are above 90°F, some injury may be expected with spot treatments when air temperatures exceed 90°F. Where state, county or local governments have more stringent temperature regulations, these regulations must be observed.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from disease, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, nematodes, improper mowing or improper applications of fertilizer and pesticides. Injury can occur if this product is applied under any of these or other stress conditions. Under any of these stress conditions, any turf damage caused



by the use of this product is beyond the control of the registrant and all risk is assumed by the buyer and/or user..

- For ground application only; aerial applications are not permitted.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the dripline of desirable trees and shrubs, since injury may result.

STATE RESTRICTIONS:

- **California, Oregon, Washington:** Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et.al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA web site: <http://www.epa.gov/espp/litstatus/wtc/qs-as.htm>.

6. Application Schedules

Apply this product to broadleaf weeds that are young and actively growing for the best results. Fall applications can provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

For the Listed Residential/domestic sites, Ornamental Turf sites, Institutional sites and Agricultural sites:

Do not apply more than two (2) broadcast treatments of this product per site per year. A second broadcast application or a follow-up application as a spot treatment is advised for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

For the Listed Non-cropland sites:

Use only one (a single) broadcast treatment of this product per site per year for woody plant infestations, or two broadcast treatments for annual and perennial weeds. It is required to wait 30 days between treatments and a spot treatment may be substituted for the second broadcast treatment if necessary.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to this product.

For newly seeded areas:

Delay application of this product to grass seedlings until after the second mowing.

For newly sodded, sprigged, or plugged areas:

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.


Reseeding interval:

Treated areas may be reseeded 3 weeks after application.

7. How Much to Use

USE RATES AND SPRAY VOLUMES:

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.



| TABLE 1. USE RATES FOR ORNAMENTAL TURF SITES, NON-CROPLAND AND SOD FARMS | | |
|--|--|--|
| Species | Rate | Spray Volume |
| Cool-season Turf | | |
| Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues | 3.0 to 4 Pints/Acre (1.1 to 1.5 fl.oz./1,000 sq.ft.) | 5 to 220 Gallons/Acre (0.125 to 5.0 Gallons/1,000 sq.ft.) |
| Creeping Bentgrass (excluding golf greens) Apply preferably in May or mid-August through September. Slight turf yellowing will disappear after about one (1) week. Note: Care should be taken to avoid overdosing bentgrass or injury may result. | 1.8 Pints/Acre (0.66 fl.oz./1,000 sq.ft.) | 145 to 220 Gallons/Acre (3.33 to 5.0 Gallons/1,000 sq.ft.) High spray volumes will provide uniform coverage. |
| Warm-season Turf | | |
| Hybrid bermudagrass, common bermudagrass, zoysiagrass, buffalograss and bahiagrass Slight turf yellowing will disappear after about one (1) week. | 3.0 to 4 Pints/Acre (1.1 to 1.5 fl.oz./1,000 sq.ft.) Sod Farms: 2 to 2.25 Pints/Acre (0.75 to 0.83 fl.oz./1,000 sq.ft.) | 5 to 220 Gallons/Acre (0.125 to 5.0 Gallons/1,000 sq.ft.) |
| <p>Note: Do not apply to above listed warm-season turfgrass unless some turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass.</p> <p>Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy.</p> <p>Dormant turf: This product may be applied to fully dormant bermudagrass, fully dormant Zoysiagrass, fully dormant buffalograss and fully dormant bahiagrass.</p> | | |

| TABLE 2. Restrictions on broadcast applications to ornamental turfgrass sites, sod farms, and non-cropland. | | | | |
|---|---|---|---------------------------------------|---|
| Use Site | Maximum Rate per Application | Maximum Number of Applications per Year | Minimum Interval Between Applications | Maximum seasonal rate |
| Ornamental turfgrass sites | 4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCP-P ae/A) (0.11 lb dicamba ae/A) | 2 | 30 days | 8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCP-P ae/A) (0.22 lb dicamba ae/A) |
| Sod farms | 4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCP-P ae/A) (0.11 lb dicamba ae/A) | 2 | 30 Days | 8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCP-P ae/A) (0.22 lb dicamba ae/A) |
| Non-cropland | 4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCP-P ae/A) (0.11 lb dicamba ae/A) | 2 (annual and perennial weeds) | 30 Days | 8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCP-P ae/A) (0.22 lb dicamba ae/A) |
| Non-cropland | 1.15 gal/A | 1 (woody plants) | NA | 1.15 gal/A (2.73 lb 2,4-D ae/A) (0.75 lb MCP-P ae/A) (0.25 lb dicamba ae/A) |

12/29/17

**SPOT TREATMENT:
WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS AND PUMP-UP TYPE
SPRAYERS):**

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.
- Follow-up applications as spot treatments at a 30 day interval are suitable for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **Limitations on spot treatments to residential turfgrass:** Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 1.5 fl.oz. per 1,000 sq.ft. per application (0.33 lb MCP-P acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.
- **For cool-season turfgrass (except Bentgrass) listed in Table 1:** Mix 1.1 to 1.5 fl.oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **For Bentgrass (excluding golf greens):** Mix 0.66 fl.oz. of this product per four (4.0) gallons of water for treatment of approximately 1,000 sq.ft of turfgrass. This high spray volume will provide uniform coverage. Apply any time the emerged broadleaf weeds are susceptible.
- **For warm-season turfgrass listed in Table 1:** Mix 0.75 to 0.83 fl.oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

CULTURAL TIPS:

Irrigation:

- Do not apply this product through any type of irrigation system.
- Rainfast in as little as 8 hours. Do not apply this product immediately before rainfall or irrigation.
- If possible, do not irrigate or water the turfgrass within 8 to 24 hours after application.
- If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.

Mowing:

- For optimum results, delay mowing 2 days before and until 2 days after the application of this product.

8. Broadleaf Weeds Controlled

EH-1473 Herbicide will control the following broadleaf weeds. Apply any time the emerged broadleaf weeds are susceptible.

| BROADLEAF WEEDS | | | |
|------------------------------------|---|---|------------------------------|
| annual fleabane | dovefoot geranium | lawn burweed | shepherdspurse |
| aster, white heath & white prairie | false dandelion (*spotted catsear & common catsear) | lespedeza, common | spotted spurge |
| bedstraw | field bindweed | mallow, common | spurge |
| beggarticks | (*morningglory & creeping jenny) | matchweed | sunflower |
| beggarweed, creeping | field madder | mouseear chickweed | thistle |
| bindweed | field oxeye-daisy (*creeping oxeye) | mustard | velvetleaf (*buttonweed) |
| birdsfoot trefoil | field pennycress | nettle | Venice mallow |
| black medic | filaree, whitestem & | old world diamond flower | Veronica (*corn speedwell) |
| broadleaf plantain | | Oxalis (*yellow woodsorrel & creeping woodsorrel) | Virginia buttonweed ** |
| buckhorn plantain | | parsley-piert | Virginia creeper |
| bull thistle | | Pennsylvania smartweed | western salsify |
| | | | white clover (*Dutch clover, |

Handwritten marks and scribbles at the bottom right of the page.

| BROADLEAF WEEDS | | | |
|--|----------------------|----------------------------|---------------------------|
| burclover | redstem | pennywort (*dollarweed) | honeysuckle clover, white |
| burdock, common | Florida pusley | pepperweed | trefoil & purplewort) |
| Carolina geranium | ground ivy | pigweed | wild carrot |
| carpetweed | groundsel | pineappleweed | wild garlic |
| chickweed, common | hairy bittercross | plantain | wild geranium |
| chicory | hawkweed | poison ivy | wild lettuce |
| cinquefoil | healall | poison oak | wild mustard |
| clover | henbit | prostrate knotweed | wild onion |
| cocklebur | horsenettle | (*knotweed) | wild strawberry |
| compassplant | horseweed | puncturevine | yarrow |
| curly dock | innocence (Blue-eyed | purple cudweed | yellow rocket |
| dandelion | Mary) | purslane | and many more broadleaf |
| dayflower | jimsonweed | ragweed | weeds |
| deadnettle | kochia | redweed | |
| Dock | knotweed | red sorrel (*sheep sorrel) | |
| dogfennel | lambquarters | roundleaf greenbriar | |
| * Synonyms | | | |
| ** A repeat application may be required in 30 days | | | |

9. For Use in Non-Cropland: Brush and Weed Control

High volume foliar applications:

Apply up to 1.15 gallons of product per 100 gallons of water or apply a 1.0 to 1.15% vol/vol spray solution as a full cover spray with high volume equipment. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g. drought).

Spray broadleaf weeds, woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars. The minimum total volume of spray solution required for adequate coverage of solid stands of mixed brush is 25 gallons of spray solution per treated acre. The spray preparation chart for applications on a spray-to-wet basis is shown below in Table 3.

Table 3. Instructions for preparing 25 to 100 gallons of spray solution at 1.0 to 1.15% spray concentration with water for high volume foliar applications.

| Spray solution per acre, Gallons | Amount of Product Needed for Spray Concentration of: | |
|----------------------------------|--|----------|
| | 1.0% | 1.15% |
| 25 | 2.0 pt | 2.3 pt |
| 75 | 6.0 pt | 6.9 pt |
| 100 | 1.0 gal | 1.15 gal |


Equal measures: 1 gallon = 4 quarts = 8 pints = 128 fl.oz.

For Backpack Sprayers and Hand Pump-up Sprayers.

Table 4. Instructions for preparing 1 to 3 gallons of spray solution at 1.0 to 1.15% spray concentration with water for high volume foliar applications.

| Gallons of Water | Amount of Product Needed for Spray Concentration of: | |
|------------------|--|-------------------------------|
| | 1.0% | 1.15% |
| 1 | 1.25 fl.oz. (2.5 Tablespoons) | 1.5 fl.oz. (3 Tablespoons) |
| 2 | 2.5 fl.oz. | 3.0 fl.oz. |
| 3 | 3.8 fl.oz. | 4.5 fl.oz. |

Equal measures: 1 fl.oz. = 2 tablespoons (Tbs.) = 6 teaspoons (tsp.)



| BRUSH [WOODY PLANTS]: | | | |
|------------------------------|--------------|-----------------|-----------------------------|
| Ash | Cedars | Multiflora rose | Sumac |
| Aspen | Elms | Oak | Willow |
| Birch | Gooseberry | Poison ivy | And many more brush species |
| Black cherry | Honey locust | Poison oak | |
| Brambles | Kudzu | Shortleaf pine | |

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]


[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

[Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing



APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

Advertising claims which may be presented on the container labels of supplemental registrants.


- Water-based formula
- Water-based formulation
- Rain-fast in as little as 8 hours
- Rain-fast [Rainproof] in 8 hours
- Designed for woody brush control
- Proven performance
- Consistently controls dandelion, spurge, clover, plantain, ground ivy, oxalis and [various other listed weeds].
- Controls all major lawn [turf] weeds
- Effective post-emergent control at a wide range of temperatures
- From the makers of Trimec® herbicides.
- Trimec® is a registered trademark of PBI/Gordon Corporation.
- A Trimec Complex [LOGO]



- Genuine Trimec Herbicide [LOGO]



- [For] fast dandelion knock-down
- Extra 2,4-D for fast knock-down of dandelions
- For information call XXX-XXX-XXXX [contact www.xxx-xxxx.com]
- For use on: Bluegrass, Fescues, Perennial Ryegrass, [Bermudagrass, Zoysiagrass (other warm-season species listed)]
- For use on cool- and warm-season turf[grass]
- Economical — lower cost water-based formulation
- Easy cleanup
- Suitable for applications up to 90°F
- Foliar absorption
- Low-odor [formula]
- Low-odor formulation[!]
- Outstanding broadleaf weed control
- Four (4) ingredients provides a broad spectrum of weed control



- Four (4) ingredients controls a broad spectrum of weeds
- Mixed amine formulation [DEA and DMA]
- No lingering odor
- Exclusive formulation
- Unlike any other formulation
- Treats up to 2.67 acres
- For use on sod farms
- For use on [sites listed]
- Controls Dandelion, Clover, Knotweed, Henbit, Chickweed, Plantain, Spurge and many other broadleaf weeds.
- Mixed Amine Formula

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00931.20100609.notif-proposed-clean.doc
2. **Reason for Issue:** delete AZ restriction

