- 2217-939

08-19-20110

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

Mr. James Kunstman Director of Regulatory Services PBI/Gordon Corporation 1217 West 12th Street P.O. Box 014090 Kansas City, MO 64101-0090

AUG 19 201

Subject: Label Amendment – Adding an "If on Skin" Statement to the First Aid Box Product Name: EH-1480 Herbicide EPA Registration Number: 2217-939 Decision Number: 447529 Submission Date: March 31, 2011

Dear Mr. Kunstman:

The label amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable.

While no additional data is being requested at this time, any marketing claims made on the pesticide label must be substantiated by data maintained in your files. If data supporting marketing claims made on the product label is not available then those claims must be removed.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

<u>Products shipped after 18 months from the date on this notice or the next printing of the</u> <u>master label whichever occurs first, must bear the new label language</u>. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of this product constitutes acceptance of these conditions. <u>You must submit</u> <u>one copy of the final printed label before the product is released for shipment</u>. If you have any questions please contact Michael Walsh by phone at (703) 308-2972 or via email at walsh.michael@epa.gov".

SEE NEXT PAGE

Page 2 of 2 Label Amendment – Adding to First Aid Section Product Name: EH-1480 Herbicide EPA Registration Number: 2217-939 Decision Number: 447529

(

Sincerely,

Ć

Jahnyn V. Ment

Kathryn V. Montague Product Manager (23) Herbicide Branch Registration Division (7505P)

Enclosure

FH-1480 HERBICIDE

EPA Reg. No. 2217-939

AUG 19 2011 Under the Faderal Insecticide. Pungicida, and Sodernicide Act. as amended, for the pesticide registered under

cecc c

cccc

LLLL cicc

LILLE

ccrect

ACCEPTED

ACTIVE INGREDIENT:	A MOS. 190 2017-1211	
2,4-D, dimethylamine salt [CAS No. 2008-39-1]		
Quinclorac [CAS No. 84087-01-4]		
Dicamba, dimethylamine salt [CAS No. 2300-66-5]		
Dithiopyr [CAS No. 97886-45-8]		
INERT INGREDIENTS:		
TOTAL		

THIS PRODUCT CONTAINS:

0.393 lb 2.4-dichlorophenoxyacetic acid equivalent per gallon or 4.53%

0.157 lb 3.7-dichloro-8-quinolinecarboxylic acid per gallon or 1.81%

0.036 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.42%

- 0.075 lb S,S-dimethyl -2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)- 3,5pyridinedicarbothioate per gallon or 0.86%
- Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if cocc swallowed. Wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves. CECCEC с

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves ever before removing. As soon as possible, wash thoroughly and change into clean clothing.

nobroo/icja **Corporation** An Employee-Owned Company 1217 West 12th Street Kansas City, Missouri 64101 0002217-00939.20110331.amend-proposed-clean.doc

1 of 11

0000

c ccc

 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- [In Optional Icon] Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING

PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. It will also act as a preemergent herbicide to prevent crabgrass and other susceptible weeds from germinating. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)

When to Use: Apply when daytime temperatures are below 90°F.

Questions & Comments: (icon phone) (icon computer) Call 1-800-xxx-xxxx or Visit our Website at <u>www.companyname.com</u>

HOW IT WORKS:

EH-1480 Herbicide contains four proven weed killers that target lawn weeds and crabgrass. This herbicide enters the lawn weeds through their leaves and moves throughout the plant to provide control. It will also form a barrier in the soil to prevent crabgrass and other susceptible weeds from germinating. And lawns listed on this label will not be harmed when used as directed.

WHERE TO USE:

EH-1480 Herbicide can be applied to residential lawns with the following turfgrass species:

Cool Season Turfgrass	Warm Season Turfgrass
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue	Bermudagrass ¹ , common and hybrid Zoysiagrass Buffalograss
Mixtures of cool season grasses containing fine fescues (red, hard and chewings)	

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

DO NOT apply EH-1480 Herbicide to the following: Bahiagrass

Bentgrass, colonial and creeping Carpetgrass Centipedegrass Dichondra Gardens including vegetables, fruit trees, vines and berries Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs) Seashore paspalum St. Augustinegrass Turfgrass species that are not listed on this label. State Restrictions: This product is not intended to be distributed, sold, or used in New York State.

GRASS AND BROADLEAF WEEDS CONTROLLED:

A partial list is presented below:

GRASS WEEDS			
Barnyardgrass Bluegrass, annual Brome Crabgrass, large and smooth	Crowfootgrass Dallisgrass Foxtail, green, yellow and giant Goosegrass	Kikuyugrass Ryegrass, annual and perennial	Sandbur Signalgrass, broadleaf Smutgrass

BROADLEAF WEEDS			2200
Annual yellow sweetclover	Cocklebur	Mugwort	Speedwell
Aster	Common mullein	Musk thistle	Spiny amaranth
Austrian fieldcress	Creeping Jenny (Creeping	Mustard	Spiny cocklebur
Bedstraw	Charlie)	Narrowleaf plantain	Spiny sowthistle
Beggarticks	Cudweed	Mustard	Spotted carsear
Betony, Florida	Curly dock	Narrowleaf vetch	Spotted source
Bindweed, field	Daisy, English	Nettle	Spurweed
Bird vetch	Daisy fleabane	Orange hawkweed	Stinging nettle
Bitter wintercress	Daisy, oxeye	Oriental cocklebur	Strawberry, India mock
Bittercress, hairy	Dandelion	Oxalis	Tall nottle
Bitterweed	Dichondra	Parsley-piert	Tall vervain
Black-eyed Susan	Dogbane	Parsnip	Tansy ragwort
Black medic	Dogfennel	Pearlwort	Tansy mustard
Black mustard	Dollarweed	Pennycress	Tanweed
Blackseed plantain	Elderberry	Peppergrass	Thistle
Blessed thistle	False dandelion	Pepperweed	Trailing crownvetch

BROADLEAF WEEDS			
Bloodflower milkweed	Falseflax	Pigweed	Tumble mustard
Blue lettuce	False sunflower	Pineapple weed	Tumble pigweed
Blue vervain	Fiddleneck	Pineywoods bedstraw	Velvetleaf
Bracted plantain	Florida pusley	Plains coreopsis (tickseed)	Venice mallow
Brassbuttons	Frenchweed	Plantain	Virginia buttonweed
Bristly oxtongue	Galinsoga	Poison ivy	Virginia creeper
Broadleaf dock	Geranium	Poison oak	Virginia pepperweed
Broadleaf plantain	Goathead	Pokeweed	Wavyleaf builthistle
Broomweed	Goldenrod	Poorjoe	Western clematis
Buckhorn	Ground ivy	Prairie sunflower	Western salsify
Buckhorn plantain	Gumweed	Prickly lettuce	White mustard
Bulbous buttercup	Hairy fleabane	Prickly sida	Wild aster
Bull thistle	Hawkweed	Prostrate knotweed	Wild buckwheat
Bullnettle	Healall	Prostrate pigweed	Wild carrot
Burclover	Heartleaf drymary	Prostrate spurge	Wild four-o'clock
Burdock	Heathaster	Prostrate vervain	Wild garlic
Burning nettle	Hedge bindweed	Puncturevine	Wild geranium
Burweed	Hedge mustard	Purslane, common	Wild lettuce
Buttercup	Hemp	Ragweed	Wild marigold
Buttonweed	Henbit	Red sorrel	Wild mustard
Canada thistle	Hoary cress	Redroot pigweed	Wild onion
Carolina geranium	Hoary plantain	Redstem filaree	Wild parsnip
Carpetweed	Hoary vervain	Rough cinquefoil	Wild radish
Catchweed bedstraw	Horsenettle	Rough fleabane	Wild rape
Catnip	Jimsonweed	Russian pigweed	Wild strawberry
Catsear	Knawel	Russian thistle	Wild sweet potato
Chickweed, common	Knotweed	Scarlet pimpernel	Wild vetch
Chickweed, mouseear	Kochia	Scotch thistle	Wild violet
Chicory	Lambsquarters	Sheep sorrel	Woodsorrel
Cinquefoil	Lespedeza	Shepherdspurse	Woolly croton
Clover, crimson	Mallow	Slender plantain	Woolly morningglory
Clover, hop	Marestail	Smallflower galinsoga	Woolly plantain
Clover, red	Matchweed	Smooth dock	Wormseed
Clover, strawberry	Mexicanweed	Smooth pigweed	Yarrow
Clover, sweet	Milk vetch	Sorrel	Yellow rocket
Clover, white	Morningglory	Sowthistle	Yellowflower pepperweed
Cockle	Mouseear hawkweed	Spanishneedles	and other broadleaf weeds

C

6

WHEN TO USE:

Apply this product when weeds are small and actively growing.

(

Weed species	Season	Timing	Maximum number of broadcast applications ¹	Application interval, days	Comments
Crabgrass (large and smooth):	Spring (Pre Emergent)	This product may be applied 1 to 2 weeks before the expected germination of crabgrass.	2	14	When used as a pre emergent this product. will provent crabgrass cseeds from coord germinating and will coord coord of existing corrabgrass plants.
Crabgrass (large and smooth):	Spring (Post Emergent)	From crabgrass emergence until the crabgrass is 3 to 4 inches in height	2	14	When used as a post emergent this product will control existing crabgrass and prevent crabgrass seeds from germinating.

Weed species	Season	Timing	Maximum number of broadcast applications ¹	Application interval, days	Comments
Foxtail (green, giant and yellow):	Spring and early summer	During the early growth stages or when foxtail is 1 to 4 inches in height.	2	14	Re-treat at the same rate at 14 days later for dense populations.
Broadleaf signalgrass	Spring and early summer	During the early growth stages or during the 3 to 7 leaf stage.	2	14	Re-treat at the same rate at 14 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Spring	When broadleaf weeds are small and actively growing.	2	14 to 28	Apply when broadleaf weeds are young and actively growing with adequate soil moisture. If needed, retreat at the same rate 14 to 28 days later.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Fall	When broadleaf weeds are young and actively growing	2	14 to 28	Apply when broadleaf weeds are young and actively growing with adequate soil moisture. If needed, retreat at the same rate 14 to 28 days later.

quinclorac, 0.09 lb dicamba ae, and 0.19 lb dithiopyr per acre per application). The maximum number of broadcast applications is limited to 2 per year. Do not apply more than 2 broadcast applications per year. The maximum seasonal rate is 15.0 fl.oz. of product per 1,000 sq.ft. (2.0 lb 2,4-D ae, 0.80 lb quinclorac, 0.18 lb dicamba ae, and 0.38 dithiopyr per acre per year).

HOW MUCH TO USE: I. ENTIRE LAWN TREATMENTS WITH DIAL TYPE HOSE-END SPRAYERS

Three Easy Steps To Mix And Apply:

1. FILL

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- The application rate for this product is 7.5 fl.oz. of product per 1,000 sg.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 7.5 fl.oz. cf product for every 1,000 sq.ft. of lawn area. Over-application or rates above those recommended on this label can cause turf injury. occo 0000
- Do not add water to the sprayer jar.

2. SET THE DIAL

For hose-end sprayers: Set the dial to the rate per gallon and keep the dial set to this setting while spraying. The dial setting or rate per gallon for this product is 3.0 oz. of product per gallon of watel.

Types of hose-end sprayers	Dial setting cccc
Ortho® Dial'N Spray® Hose-End Sprayer	3.0 oz.
Hudson® Hose End Liquid Concentrate Sprayer	3.0 fl.oz.
Gilmour® Dial-A-Mix® Lawn and Garden Sprayer	18 Teaspoons or 6 tablespoons or 3.0 oz.

3. SPRAY - Apply evenly over the lawn. Avoid skips and excess overlaps during application.

II. ENTIRE LAWN TREATMENTS WITH READY-TO-USE HOSE-END SPRAYERS :

This product may be packaged as a ready-to-use product with no mixing. The retail plastic container is fitted with a sprayer body that delivers spray droplets at a fixed rate.

For READY-TO-USE Hose End Sprayers:

• This retail sprayer bottle will cover the treated areas described below when this product is applied at 7.5 fl.oz. of product per 1,000 sq.ft. Over-application or rates above those recommended on this label can cause turf injury.

		Amount of prod	luct to treat appro	ximate area, sq. f	t.
Species	fl.oz./1,000 sq.ft.	1.0 pt. (16 fl.oz.) sprayer jar or retail container	1.0 qt. (32 fl.oz.) sprayer jar or retail container	2.0 qt. (64 fl.oz.) retail container	1 gallon (128 fl.oz.) retail container
Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season species Bermudagrass ¹ common and hybrid Zoysiagrass and Buffalograss ²	7.5 fl.oz. /1,000 sq.ft.	2,100 sq.ft.	4,200 sq.ft.	8,500 sq.ft.	17,000 sq.ft.

Approximate treated areas for four retail container sizes:

Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F. Do not apply to Tifgreen (328) hybrid Bermudagrass as it may cause undesirable turf injury.

² Do not use on seedling buffalograss during the spring months of the initial year of establishment until the turf is achieved full green up and has an healthy root system.

Instructions:

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
- **CONNECT**: Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the "OFF" position.
- SPRAY: Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet.

Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl.oz. container will contrainer to guide coverage. This xx fl.oz. container to guide coverage. This xx fl.oz. coverage. This xx fl.oz.

• FINISH: To stop spraying, turn [rotate] spray nozzle lever to the "OFF" position. Eurn off water accorrection faucet. To relieve pressure, turn [rotate] spray nozzle to the "ON" position and point prayer away correction any desirable plants before disconnecting the sprayer nozzle from the garden hose.

ccecce

ι. ι ιιιιιί

III. HAND-OPERATED SPRAYERS:

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up):

- Apply to actively growing grass and broadleaf weeds during the early growth stages.
- Measure and mix the proper amount of product with water as described below. Mix 3.0 fl.oz. of product with 1 gallon of water to treat 400 sq.ft. of lawn. See below for additional spray preparations
- Spray evenly over the lawn. Spray coverage should be uniform and complete. Over-application or rates above those recommended on this label can cause turf injury.

Amount of Water in Sprayer, Gallons	Approximate Treated Area, Square feet
1 gallon	400 sq.ft.
1.5 gallons	600 sq.ft.
2 gallons	800 sq.ft.
3 gallons	1,200 sq.ft.
-	Gallons 1 gallon 1.5 gallons 2 gallons

OR

Measure indicated amount of product and pour into tank. Dilute with water according to the following chart. Adjust the nozzle to deliver coarse spray droplets and treat the area indicated.

Amount of Product to Use, Fluid Ounces	Amount of Water in Sprayer, Gallons	Area To Treat, Square feet
3 fl.oz.	1 gallon	400 sq.ft.
6 fl.oz.	2 gallons	800 sq.ft.
9 fl.oz.	3 gallons	1,200 sq.ft.

IV. ENTIRE LAWN TREATEMENTS WITH THE ORTHO READY-SPRAY® SYSTEM:

HOW TO USE THE READY-SPRAY® SYSTEM

Connect:

- 1. Shake well before using.
- 2. Connect sprayer to hose.
- 3. Turn on water.
- 4. Extend hose to farthest area of the lawn to be treated

Spray:

- 1. To begin spraying, point nozzle in the direction you want to spray.
- 2. Bend small plastic tab back and turn knob clockwise to ON position.
- 3. Spray evenly over measured area.

Finish:

- 1. To STOP spraying, rotate the knob counterclockwise to the OFF position.
- 2. Turn off water.
- 3. Relieve water pressure by bending plastic tab back and turning knob to ON position until water slows to a drip. Then, turn knob back to OFF position.
- 4. Disconnect sprayer from hose. εειε ccic **V. FOR TANK SPRAYER:** с ددرد Add 3.0 fl.oz. per gallon of water for each 400 sg.ft. ιιεί ιιια çς For broadcast application: Ortho Dial 'N Spray® Applicator ε. 1. Add 16 fl.oz. of concentrate directly into sprayer jar. DO NOT ADD WATER. LEELE ε 2. Set dial to 3.0 oz. 3. Spray evenly over 2,100 sq.ft. 4. Any unused portion should be poured back into its original container. [Over-application or rates above those recommended on this label can cause turf injury]

VI. FOR [PRODUCT NAME]:

For (Use a) Tank Sprayer:

- Add entire contents (of 1 bottle) (5 oz.) to an empty 2-gallon tank sprayer. Slowly fill with 2 gallons of clean tap water.
 - For Broadcast Application Ortho® Dial 'N Spray® Applicator
 - 1. Set dial to 3.0 oz.
 - 2. Add entire contents of 1 bottle (5 oz) directly into sprayer jar. DO NOT ADD WATER.
 - 3. Spray evenly over 675 sq.ft.

[Over-application or rates above those recommended on this label can cause turf injury]

CULTURAL TIP	S AND NOTES:	
When to Apply:	 For established lawns For newly seeded areas For newly sodded, sprigged, or plugged areas 	 Apply when daytime temperatures are below 90°F. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing. Apply this product 3 to 4 weeks after the sodding, sprigging, or plugging operations.
Reseeding, Overseeding, or Sprigging :		 Reseeding, overseeding, or sprigging of treated areas within 3 months after application of this product, may inhibit the establishment of desirable turfgrasses.
Irrigation:	Non-stressed turfgrass Stress from high temperatures, drought, insects or diseases.	 Preemergence: Adequate soil moisture and favorable growing conditions enhance the performance of this product. This product is effective for crabgrass control when activated by 0.5 inch of rainfall or irrigation. Postemergence: Avoid applications of this product when turfgrasses are under stress since injury may result. If dry conditions exist, schedule irrigations (watering) before and after the application. For best results, delay the irrigation until approximately 24 hours after application of this product.
Mowing:		 Delay mowing 2 days before and until 2 days after the application of this product. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
Nontarget plants:		 Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants. Do not spray exposed roots of ornamentals and trees.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do hot refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor brees outdoor drain.

cicc

LLLC

eccc

ιζει

CCCCCC L C C

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES,

INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

[OPTIONAL WARRANTY STATEMENT]

The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- · See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on the retail containers of supplemental registrants.

- One quart covers up to 4,200 square feet.
- 1 quart treats [up to] 4,200 sq.ft.
- Covers [XXXX] sq.ft.
- Keep from freezing.
- Controls crabgrass, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills even the tough weeds...roots and all!
- Controls even the tough weeds...roots and all!
- Kills the root.
- Prevents crabgrass.
- Controls existing broadleaf weeds and crabgrass.
- Water based.
- Won't harm lawn grass when used according to label directions.

 Kills lawn weeds plus crabgrass. 		
 Kills [Controls] 200+ broadleaf weeds. 		ζζζζ
 Season-Long crabgrass control 		6 C C C C
 Kills the weeds not the lawn 		ccce
 Attach to [garden] hose and spray 	ιι ιια ί ι ι	ο οςις
Simple to use	c, C	
Easy to use	<u>ςςςιζι</u> ι α ιι	
Easy to mix	(((((ε ει
 Proven performance 	َنْ بِ (ب زير د و ب	c
 For home lawn care 		ί ι ιιιιι
 Won't harm lawn grass when used as directed. 		
 Controls (Number) weeds! 		ι ύ (ιί ι ι
 Mixes with Water - Concentrate 		C
Kill and anona 9 fam: listed ward(a)] to al		

• Kill crabgrass & [any listed weed(s)] too!

C C C C C

ιιιι

• CONTROLS

SENSITIVE WEEDS: Crabgrass, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.

HARD-TO-KILL WEEDS: Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and other broadleaf weeds.

- Kills Lawn Weeds plus Crabgrass in One Step
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grass Weeds
- [Also] Kills [Dandelion], [Crabgrass], [Clover], plus 200 Other [Listed] Weeds [As Listed]
- Four-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Kills both Crabgrass & Broadleaf weeds
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- Effectively Kills Both Broadleaf and Certain Weed Grasses
- [XX] [Pint, Quart, etc] Treats Up To [YYYY] Sq.Ft.
- Local Solutions for Local Problems
- For more Lawn and Garden Products Visit [www.website.com]

For any of the ORTHO Products:

- Makes more than XX Gallon!
- Kills Even Tough Weeds Roots & All
- Lawn Weed & Crabgrass Killer
- Kills Dandelions, Clover, and Other Broadleaf Weeds! Plus Kills Crabgrass and Other grass Weeds!
- Kills the Toughest Weeds in these Lawns (any lawn listed on label)
- Open Resealable Label for Directions & Precautions (Press to Reseal)
- People or pets may reenter treatment area after spray has dried.
- Ready-Spray
- Do Not Use on St. Augustine Lawns
 - [PRODUCT NAME]
 - Kills Weeds, Not Lawns(TM)
 - For [Use on] Lawns
 - ORTHO & Your Environment Your yard is a place for your family and pets to enjoy. That's why ORTHO products are designed with care to provide effective solutions to weed problems. For best results please follow instructions for appropriate usage, storage and disposal.
- Kills Over 200 Lawn Weeds plus [including] Crabgrass [in one easy step.
- Guaranteed [Results]
- See Ortho.com for complete Spanish Translation.
- DO NOT use on St. Augustine, including Floratam species.
- NOTICE: Buyer assumes all risks of use, storage, or handling of this product not in accordance with directions
- Rainfast in 1 Hour
- Kills Major [Lawn] Broadleaf Weeds

For [PRODUCT NAME]:

- NET 5 FL.OZ.
- Pre-Measured Concentrate Bottles
- Treats 675 sq.ft.
- No Measuring, No Mess

- [PRODUCT NAME] makes using concentrates easier than ever. No more measuring and no more mess. Mix (contents) (each pre-measured bottle) with two gallons of water to deliver consistent results. And its' revolutionary formula won't harm your lawn Guaranteed.
- Designed for tank sprayers
- 1 bottle makes 2 gallons

3. Trademark acknowledgements

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

DOCUMENT CONTROL INFORMATION

- 1. Unique Label Identifier: 002217-00939.20110331.amend-proposed-clean.doc
- 2. Reason for Issue: First Aid and corrected ad claim