


2217-954

9/15/2011

10/11

| | | |
|---|--|---|
|  <p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460</p> | EPA Reg. Number: 2217-954 | Date of Issuance: SEP 15 2011 |
| | Term of Issuance: unconditional | |
| | Name of Pesticide Product: EH-1457 Herbicide | |
| NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended) | | |
| Name and Address of Registrant (include ZIP Code): PBI/Gordon Corporation 1217 West 12th Street Kansas City, MO 64101 | | |
| <p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> | | |
| <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>The basic and alternate #1 formulations, dated 8/18/2011, are acceptable.</p> <p>This product is registered in accordance with FIFRA section 3(c)(5) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data. 2. Make the following label revision: <ol style="list-style-type: none"> a. Revise "EPA REG. NO.2217-xxx" to "EPA REG. NO. 2217-954." Assure that the establishment number and net contents are also added to the final printed label. b. Revise "ACTIVE INGREDIENT" to "ACTIVE INGREDIENTS" in the ingredient statement. 3. Within one year of the date on this registration notice, Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies must be submitted. It is recommended that the observations be made at 0, 3, 6, 9, and 12 month intervals. 4. Submit one (1) copy of the revised final printed label for the record. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.</p> <p>If you have any questions regarding this notice, please contact Beth Benbow of my staff at 703-347-8072.</p> | | |
| Signature of Approving Official: Kathryn V. Montague Project Manager 23 Herbicide Branch Registration Division (7505P) | Date: SEP 15 2011 | |

EH-1457 HERBICIDE

ACCEPTED
with COMMENTS
In EPA Letter Dated:
SEP 15 2011

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

EPA Reg. No. 2217-XXX

FOR OUTDOOR RESIDENTIAL USE ONLY

2217-954

ACTIVE INGREDIENT:

| | |
|-----------------------------------|----------------|
| MCPA, dimethylamine salt | 4.78% |
| Quinclorac | 1.56% |
| Dicamba, dimethylamine salt | 0.43% |
| OTHER INGREDIENTS: | 93.23% |
| TOTAL | 100.00% |

THIS PRODUCT CONTAINS:

- 0.338 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 3.90%
- 0.135 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 1.56%
- 0.031 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.36%

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

| First Aid | |
|--------------------------------|--|
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on skin or clothing: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |



| | |
|---|---|
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person. |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice. | |

Environmental Hazards

This pesticide is toxic to fish, aquatic invertebrates, and aquatic plants. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Precautions And Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

SHAKE WELL BEFORE USING

| |
|---|
| <p>PRODUCT FACTS:</p> <p>What it Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass, foxtail, and yellow nutsedge. (See inside for complete list)</p> <p>Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)</p> <p>When to Use: Apply when daytime temperatures are between 50° and 90°F (except on Bermudagrass - see below).</p> <p>Questions & Comments: (icon phone) (icon computer) Call 1-800-xxx-xxxx or Visit our Website at www.companyname.com</p> |
|---|

2. How It Works

EH-1457 Herbicide contains three proven weed killers that target lawn weeds and crabgrass. The herbicides enter the lawn weeds through their leaves and move inside the weeds, even down into the roots. And lawns listed on this label will not be harmed when used as directed.

3. Where To Use

| | |
|---|--|
| EH-1457 Herbicide can be applied to the following established turfgrass species: | |
| Cool season turfgrass | Warm season turfgrass |
| Kentucky bluegrass, Ryegrass, perennial and annual, Tall fescue, Bentgrass, Mixtures of cool season grasses containing fine fescues (red, hard and chewings) | Bermudagrass ¹ , common and hybrid Zoysiagrass, Buffalograss, |
| ¹ Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F. | |

| DO NOT apply EH-1457 Herbicide to the following: | |
|---|---|
| Warm season turfgrass | Other areas |
| St. Augustinegrass, Bahigrass, Centipedegrass, Carpetgrass, Seashore paspalum | Lawns with desirable clovers or legumes, Dichondra, Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Turfgrass species that are not listed on this label, Gardens including vegetables, fruit trees, vines and berries. |

4. Weeds Controlled

GRASS WEEDS:

Crabgrass, large and smooth, foxtail, (green, yellow and giant), kikuyugrass, and signalgrass (broadleaf).

| BROADLEAF WEEDS (A partial list is presented below) | | | |
|---|-----------------------------------|-----------------------------|------------------------|
| Annual yellow sweetclover | Cocklebur | Mustard | Spiny cocklebur |
| Aster | Common mullein | Narrowleaf plantain | Spiny sowthistle |
| Austrian fieldcress | Creeping Jenny (Creeping Charlie) | Narrowleaf vetch | Spotted catsear |
| Bedstraw | Cudweed | Nettle | Spotted spurge |
| Beggarticks | Curly dock | Orange hawkweed | Spurweed |
| Betony, Florida | Daisy, English | Oriental cocklebur | Stinging nettle |
| Bindweed, field | Daisy fleabane | Oxalis | Strawberry, India mock |
| Bird vetch | Daisy, oxeye | Parsley-piert | Tall nettle |
| Bitter wintercress | Dandelion | Parsnip | Tall vervain |
| Bittercress, hairy | Dichondra | Pearlwort | Tansy ragwort |
| Bitterweed | Dogbane | Pennycress | Tansy mustard |
| Black-eyed Susan | Dogfennel | Pennywort | Tanweed |
| Black medic | Dollarweed | Peppergrass | Thistle |
| Black mustard | Elderberry | Pepperweed | Trailing crownvetch |
| Blackseed plantain | False dandelion | Pigweed | Tumble mustard |
| Blessed thistle | Falseflax | Pineywoods bedstraw | Tumble pigweed |
| Bloodflower milkweed | False sunflower | Plains coreopsis (tickseed) | Velvetleaf |
| Blue lettuce | Fiddleneck | Plantain | Venice mallow |
| Blue vervain | Florida pusley | Poison ivy | Virginia buttonweed |
| Bracted plantain | Frenchweed | Poison oak | Virginia creeper |
| Brassbuttons | Galinsoga | Pokeweed | Virginia pepperweed |
| Bristly oxtongue | Goathead | Poorjoe | Wavyleaf bullthistle |
| Broadleaf dock | Goldenrod | Prairie sunflower | Western clematis |
| Broadleaf plantain | Ground ivy | Prickly lettuce | Western salsify |
| Broomweed | Gumweed | Prickly sida | White mustard |
| Buckhorn | Hairy fleabane | Prostrate knotweed | Wild aster |
| Buckhorn plantain | Hawkweed | Prostrate pigweed | Wild buckwheat |
| Bulbous buttercup | Healall | Prostrate spurge | Wild carrot |
| Bull thistle | Heartleaf drymary | Prostrate vervain | Wild four-o'clock |
| Bullnettle | Heathaster | Puncturevine | Wild garlic |
| Burclover | Hedge bindweed | Purslane, common | Wild geranium |
| Burdock | Hedge mustard | Ragweed | Wild lettuce |
| Burning nettle | Hemp | Red sorrel | Wild marigold |
| Burweed | Henbit | Redroot pigweed | Wild mustard |
| Buttercup | Hoary cress | Redstem filaree | Wild onion |
| Buttonweed | Hoary plantain | Rough cinquefoil | Wild parsnip |
| Canada thistle | Hoary vervain | Rough fleabane | Wild radish |
| Carolina geranium | Horsenettle | Russian pigweed | Wild rape |
| Carpetweed | Jimsonweed | Russian thistle | Wild strawberry |
| Catchweed bedstraw | Knawel | Scarlet pimpernel | Wild sweet potato |
| Catnip | Knotweed | Scotch thistle | Wild vetch |
| Catsear | Kochia | Sheep sorrel | Wild violet |
| Chickweed, common | Lambsquarters | Shepherdspurse | Woodsorrel |
| Chickweed, mouseear | Lespedeza | Slender plantain | Woolly croton |
| Chicory | | | Woolly morningglory |

| BROADLEAF WEEDS (A partial list is presented below) | | | |
|--|-------------------|-----------------------|-------------------------|
| Cinquefoil | Mallow | Smallflower galinsoga | Woolly plantain |
| Clover, crimson | Matchweed | Smooth dock | Wormseed |
| Clover, hop | Mexicanweed | Smooth pigweed | Yarrow |
| Clover, red | Milk vetch | Sorrel | Yellow rocket |
| Clover, strawberry | Morningglory | Sowthistle | Yellowflower pepperweed |
| Clover, sweet | Mouseear hawkweed | Spanishneedles | and other broadleaf |
| Clover, white | Mugwort | Speedwell | weeds |
| Cockle | Musk thistle | Spiny amaranth | |

5. When To Use

Apply this product when weeds are small and actively growing.

CRABGRASS (large and smooth):

- **Spring:** Apply this product from crabgrass emergence until the crabgrass is 3 to 4 inches in height. This period begins the first time that you can see crabgrass in your lawn. Re-treat at the same rate at 21 days later for dense populations.

FOXTAIL (GREEN GIANT AND YELLOW):

- **Spring and early summer:** Apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height. Re-treat at the same rate at 21 days later for dense populations.

BROADLEAF WEEDS:

- **Spring:** Apply when **dandelion, plantain, buttercup, speedwell, knotweed** and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.
- **Fall:** Apply when **henbit, chickweed, white clover, ground ivy, lawn burweed** and other broadleaf weeds are young and actively growing with adequate soil moisture for best results.

6. How Much To Use

Limitations on broadcast applications to residential turfgrass

The maximum application rate is 8 fl.oz. of product per 1,000 sq.ft. per application (0.92 lb MCPA acid equivalent and 0.08 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year, with a minimum of 21 days between applications. The maximum seasonal rate is 16 fl.oz. of product per 1,000 sq.ft. (1.84 lb MCPA ae and 0.17 lb dicamba ae per acre acid equivalent per acre).

6.1 Entire Lawn Treatments

THREE EASY STEPS TO MIX AND APPLY:

1. Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.
2. **For hose-end sprayers:** Set the dial to the RATE PER GALLON and keep the dial set to this setting while spraying. The RATE PER GALLON for this product is 4 fl.oz. of product per gallon. See instructions below.

For hand-operated sprayers (backpack sprayers, compression (pump-up) sprayers, knapsack sprayers or tank type): The RATE PER 1,000 SQ.FT. for this product is 8 fl.oz. of product per 1,000 sq.ft. of lawn. Measure and mix the proper amount of product with water as described below.

3. Spray evenly over the lawn.

For Ortho® Dial'N Spray® Hose-End Sprayer, Hudson® Hose End Liquid Concentrate Sprayer or Gilmour® Dial-a-Mix® Lawn and Garden Sprayer:

A. Set the dial

For Ortho® Dial'N Spray® Hose-End Sprayer: Set the dial on 4 oz as the RATE PER GALLON and keep the dial set to this setting while spraying.

Hudson® Hose End Liquid Concentrate Sprayer: Set the dial on 4 oz as the RATE PER GALLON and keep the dial set to this setting while spraying.

Gilmour® Dial-A-Mix® Lawn and Garden Sprayer: Set the dial on 4 oz or 8 TBS or 24 TSP as the RATE PER GALLON and keep the dial set to this setting while spraying.

B. Fill

Determine the total lawn area to be sprayed. The application rate for this product is 8 fl.oz. of product per 1,000 sq.ft. of lawn. Add an adequate amount of herbicide to the spray bottle to treat the total lawn area, or add 8 fl.oz. of product for every 1,000 sq.ft. of lawn area. Do not add water to the sprayer jar.

C. Spray

Spray evenly over the lawn.

Example:

Set the dial: Set the dial on 4 oz.

Fill: For a lawn area of 3,000 sq.ft, pour or measure 24 fl.oz of product into sprayer jar. Do not add water.

Spray: Spray uniformly on 3,000 sq.ft. of lawn.

For READY-TO-SPRAY Hose End Sprayers:

This retail sprayer bottle will cover the treated areas described below when this product is applied at 8 fl.oz.of product per 1,000 sq.ft.

Approximate treated areas for four retail container sizes:

| Species | Amount of product (to treat approximate area, sq.ft.) | | | | |
|---|---|---|---|--------------------------------------|--|
| | fl.oz./1,000 sq.ft. | 1.0 pt. (16 fl.oz.) sprayer jar or retail container | 1.0 qt. (32 fl.oz.) sprayer jar or retail container | 2.0 qt. (64 fl.oz.) retail container | 1 gallon (128 fl.oz.) retail container |
| Kentucky bluegrass Ryegrass, perennial and annual Tall fescue Mixtures of cool season species Bermudagrass ¹ , common and hybrid Zoysiagrass and Buffalograss, Bentgrass | 8 fl.oz./1,000 sq.ft. | 2,000 sq.ft. | 4,000 sq.ft. | 8,000 sq.ft. | 16,000 sq.ft. |

¹Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.

Instructions:

- Determine (Measure) the total lawn area to be sprayed. For rectangular areas, multiply the length by the width to obtain total square feet.

- **CONNECT:** Remove bottle cap and screw the spray nozzle onto the bottle. Attach [connect] garden hose to spray nozzle. Turn [rotate] the spray nozzle lever to the "OFF" position.
- **SPRAY:** Turn on water at faucet. Extend hose to the farthest area of lawn [turf] to be treated. To begin spraying, turn [rotate] the spray nozzle lever to the "SPRAY" position. The herbicide mixes automatically with the water. Walk at a steady pace while spraying evenly and working towards the faucet. Refer to view strip [graduated scale] on side of container to guide coverage. This xx fl.oz. container will treat up to xxxx sq.ft. of turfgrass.
- **FINISH:** To stop spraying, turn [rotate] spray nozzle lever to the "OFF" position. Turn off water at faucet. To relieve pressure, turn [rotate] spray nozzle to the "ON" position and point sprayer away from any desirable plants before disconnecting the sprayer nozzle from the garden hose.

[OR]

CONNECT

1. Shake well before use.
2. Connect sprayer to garden hose.
3. Turn on water.

SPRAY

1. Extend hose to farthest area to be treated, point nozzle in direction to be sprayed and turn on sprayer. Work back toward the faucet to avoid walking over treated grass.
2. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas.



[Operating the Sprayer Top


Water control knob (closest to hose). [Graphic] Twist counterclockwise [clockwise] for 'On' & clockwise [counter-clockwise] for 'Off'.

Weed killer control knob. [Graphic] Twist knob until flat side is next to protruding plastic tab. This will release the safety feature. Push in for 'On' & out for 'Off'.

FINISH

1. Turn off sprayer.
2. Turn off water at faucet.
3. Relieve pressure in hose by turning water back on at sprayer until water slows to a drip. Close sprayer. Disconnect sprayer from hose.

[OR]

| | |
|---|--|
| <p>HOW TO USE</p> <p>Shake Well Before Use</p> | <p>CONNECT</p> <ol style="list-style-type: none"> 1. Shake well before use. 2. Connect sprayer to garden hose. 3. Turn on water. |
| | <p>SPRAY</p> <ol style="list-style-type: none"> 1. Extend hose to farthest area to be treated, point sprayer nozzle in direction to be sprayed. 2. Begin spraying – Push & hold lock tab away from knob, use other hand to turn knob clockwise [counter-clockwise] to ON position. 3. Walk at a steady pace while spraying, using an even sweeping motion, and slightly overlap the treated areas. Work back toward the faucet to avoid walking over wet, treated grass. |
| |  |
| | <p>FINISH</p> <ol style="list-style-type: none"> 1. To stop spraying, push & hold lock tab away from knob and use other hand to turn knob [clockwise] counter-clockwise to OFF position. 2. Turn off water at faucet. 3. Relieve water pressure in hose by turning sprayer back ON until water slows to a drip then turn sprayer OFF. [OR Relieve water pressure in hose by turning sprayer to WATER position until water slows to a drip, then turn sprayer to OFF position.] Disconnect sprayer from hose. |

6.2 Spot Treatments For Small Areas Or Individual Weeds

For hand-operated sprayers (backpack, knapsack, tank-type or pump-up): Mix 4 fl.oz. of product with 1 gallon of water. Note: Approximately one (1) gallon of spray solution will treat 500 sq.ft. See below for additional spray preparations. Shake well before spraying. Apply to actively growing weeds during the early growth stages. Spray coverage should be uniform and complete.

| Amount of Product to Use, Fluid Ounces | Amount of Water in Sprayer, Gallons | Approximate Treated Area, Square feet |
|--|-------------------------------------|---------------------------------------|
| 4 fl.oz. | 1 gallon | 500 sq.ft. |
| 6 fl.oz. | 1.5 gallons | 750 sq.ft. |
| 8 fl.oz. | 2 gallons | 1,000 sq.ft. |
| 12 fl.oz. | 3 gallons | 1,500 sq.ft. |

Equal measures: 1 cup = 8 fl.oz. = 0.5 pint. 4 fl.oz. = 8 Tablespoons

[OR]

Measure indicated amount of product and pour into tank. Dilute with water according to the following chart. Adjust the nozzle to deliver coarse spray droplets and treat the area indicated.

| Amount of Product to Use, Fluid Ounces | Amount of Water in Sprayer, Gallons | Area To Treat, Square feet |
|--|-------------------------------------|----------------------------|
| 4 fl.oz. | 1 gallon | 500 sq.ft. |
| 8 fl.oz. | 2 gallons | 1,000 sq.ft. |
| 12 fl.oz. | 3 gallons | 1,500 sq.ft. |

Equal measures: 1 cup = 8 fl.oz. = 0.5 pint. 4 fl.oz. = 8 Tablespoons

7. Cultural Tips

For newly seeded areas:

- Do not apply within 4 weeks after seedling emergence of Kentucky bluegrass, fine fescue mixtures and perennial ryegrass. Apply this product to newly seeded grasses when well-established (approximately 6 weeks after seedling emergence) or after the third mowing.

For newly sodded, sprigged, or plugged areas:

- The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Seeding:

- Turf species listed on this label can be seeded into the treated areas at four (4) weeks after the application of this product.

Irrigation:

- Do not apply this product immediately before rainfall or irrigation. For best results, do not irrigate or water the turfgrass within 24 hours after application.
- Adequate moisture and mild environmental conditions enhance the performance of EH-1457 Herbicide. If dry conditions exist, schedule irrigations (watering) at 24 hours before and after the application.
- Avoid applications of this product when turfgrasses are under stress since injury may result.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

8. Prohibitions

- Apply when daytime temperatures are between 50° and 90°F. Do not treat when air temperatures exceed 90°F as damage to turfgrass may occur. For bermudagrass lawns do not apply when daytime temperatures exceed 85° F.
- Do not apply at rates above those specified on this label.
- Do not apply this product to food crops (gardens, fruits and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL,

CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on the retail containers of supplemental registrants.

- One quart covers up to 4,000 square feet.
- 1 quart treats [up to] 4,000 sq.ft.
- Keep from freezing.
- Controls crabgrass, yellow nutsedge, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills even the tough weeds...roots and all!
- Controls even the tough weeds...roots and all!
- Kills the root.
- Water based.
- Won't harm lawn grass when used according to label directions.
- Kills lawn weeds plus crabgrass.
- Kills [Controls] 200+ broadleaf weeds.
- Kills the weeds not the lawn
- Attach to [garden] hose and spray
- Simple to use
- Easy to use
- Easy to mix
- Proven performance
- For home lawn care
- Won't harm lawn grass when used as directed.
- Controls (Number) weeds!
- Mixes with Water - Concentrate
- Kill crabgrass, yellow nutsedge & [any listed weed(s)] too!
- **CONTROLS**
 - **SENSITIVE WEEDS:** Crabgrass, yellow nutsedge, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.
 - **HARD-TO-KILL WEEDS:** Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and many other broadleaf weeds.
- Kills Lawn Weeds plus Crabgrass in One Step
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step
- Kills Crabgrass plus Broadleaf Weeds
- Kills Listed Weeds
- Kills Hard-to-Control Weeds
- Systemic [Herbicide] [Activity]
- Kills Both Broadleaf and Grass Weeds

- Kills Dandelion, Crabgrass, Clover, plus 200 Other listed Weeds
- New!
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds
- Kills both Crabgrass & Broadleaf weeds
- Effectively Kills Both Broadleaf and certain weed grasses
- Kills Lawn Weeds plus Crabgrass in One Easy Step
- Won't Harm Lawns* (when used as directed)
- Kills 200+ Weeds (to the root)
- Treats 5,000 sq.ft.
- Starts Working Immediately
- All Weather Formula
- Bayer 100% Satisfaction Guaranteed* (or your money back)
- [optional advertising claims when product is sold in "low-water" hose-end sprayers on the order of 40 to 1 dilution ratio or as determined by testing]
 - Water [Saver] [Saving] Formula
 - Water [Saver][Saving] Applicator
 - Water [Saver] [Saving] Sprayer
 - Uses Up To [xx][50]% Less Water* (with the footnote disclaimer: *vs. leading weed hose end sprayers)

3. Trademark acknowledgements

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00XXX.20110909.new-proposed-EH1457-revised-clean.docx
2. Reason for Issue: new registration, revised per EPA comments