

2217-958

7/24/2012

1 of 14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington D C 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

James L. Kunstman Ph D
PBI/Gordon Corporation
Post Office Box 014090
Kansas City Missouri 64101

JUL 24 2012

Subject Amendment revising the Protective Eye wear of applicators statement
Product name EH-1506 Herbicide
EPA Reg No 2217 958
Application Dated March 6 2012

Dear Mr Kunstman

The labeling referred to above submitted in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA) as amended is acceptable

A stamped copy of your label is enclosed for your records This label supersedes all previously accepted labels You must submit one (1) copy of the final printed label before you release the product for shipment Products shipped after eighteen (18) months from the date of this letter must bear the new revised label If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA §6(e) Your release for shipment of the product constitutes acceptance of these conditions

If you have any questions please contact Grant Rowland at 703 347 0254 or at Rowland Grant@epa.gov

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn Montague".

Kathryn Montague
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2014

ACCEPTED
JUL 24 2012
Under the Federal Insecticide
Fungicide, and Rodenticide Act
is excluded for a pesticide
registration
" 2917-958

EH-1506 HERBICIDE

EPA Reg No 2217-958

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170

ACTIVE INGREDIENT	
Dimethylamine salt of MCPA	31.08%
Dimethylamine salt of MCPP p	8.18%
Dimethylamine salt of dicamba	2.77%
INERT INGREDIENTS	<u>57.97%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS
 2.38 lb 2-methyl-4-chlorophenoxyacetic acid equivalent per gallon or 25.38%
 0.63 lb (+) (R) 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 6.76%
 0.22 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 2.30%
 TRIMEC® is a registered trademark of PBI/Gordon Corporation

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si Usted no entiende la etiqueta busque a alguien para que se la explique a Usted en detalle (If you do not understand the label find some one to explain it to you in detail)

KEEP FROM FREEZING

STOP! READ THE ENTIRE LABEL FIRST OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING Causes substantial but temporary eye injury Harmful if swallowed Avoid contact with skin or clothing Do not get in eyes Remove and wash contaminated clothing before reuse Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

Personal Protective Equipment (PPE)

Some materials that are chemical resistant to this product are natural rubber If you want more options follow the instructions for category A on an EPA chemical resistance category selection chart

- All mixers loaders applicators and other handlers must wear
- protective eyewear (goggles face shield or safety glasses)
 - long sleeved shirt and long pants
 - shoes and socks



- chemical resistant gloves when mixing loading or using any hand held equipment
- See engineering controls for additional requirements

Applicators may choose not to wear protective eyewear with dilution rates greater (higher) than 4:1 or greater (higher) than 4 parts of water to 1 part of product

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6) the handler PPE requirements may be reduced or modified as specified in the WPS

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations	
Users should wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco, and using toilet.	
<ul style="list-style-type: none"> • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 	
Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.	

First Aid	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or on clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage.	

Environmental Hazards

This pesticide may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow.

may result in groundwater contamination Application around a cistern or well may result in contamination of drinking water or groundwater

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170

This standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is
coveralls worn over short sleeved shirt and short pants
chemical resistant footwear plus socks
chemical resistant gloves made of any water proof material
protective eyewear

Non Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries or greenhouses

Reentry Statement Do not enter or allow people (or pets) to enter the treated area until sprays have dried

1 Product Description

Designed for [turfgrass applications] [non crop] [and IVM (Industrial Vegetation Management)] applications this product contains three active ingredients

- (1) MCPA is an auxin type herbicide which is a class of plant growth regulators It is absorbed through the leaves and is translocated to the growing points of the plant causing weed stems to curl and twist leaf cupping and withering and eventual plant death
- (2) MCPP p (mecoprop p) is an auxin type herbicide which is a class of plant growth regulators It is absorbed through the leaves and is translocated to the growing points of the plant causing weed stems curl and twist leaf cupping and withering and eventual plant death
- (3) Dicamba is absorbed through the leaves and roots and has multiples modes of actions for hard to kill broadleaf weeds

Combining these herbicides provides a very wide spectrum of weed control for susceptible weeds. EH 1506 Herbicide controls weeds by affecting multiple sites within the broadleaf weeds. The symptoms of susceptible broadleaf weeds include leaf and stem curl or twisting and weed yellowing. These combined herbicides provide limited residual activity.

2 Where to Use

This product provides broadleaf control in the following sites

Ornamental Turfgrass sites

- **Residential/domestic sites** are defined as areas associated with the household or home life including but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes. **Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
 - **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
 - **Non-cropland sites** including farmyards, fencerows or fence lines, highway rights of way (principal interstate, county, private, and unpaved roads), Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians, Industrial sites, Lumberyards, tank farms, fuel or equipment storage areas, Municipal, state, and federal lands, Airports and military installations, railroad rights of way, railroad yards, railroad crossings and railroad bridge abutments, Utility rights of way, telephone, pipeline, electrical powerlines, and communication transmission lines.
- Agricultural site** Commercial sod production

Prohibitions of Sites

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
 - Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
 - Do not apply to agricultural drainage water or on agricultural ditchbanks.
 - Do not apply or allow this product to come into direct contact with cotton, grapes, tobacco, vegetable crops, flowers, fruit, or ornamental trees, or other desirable broadleaf plants. Small amounts of spray drift may injure susceptible plants, including ornamental trees or shrubs.

Turfgrass tolerance

- The turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur. Adverse environmental conditions may reduce the selectivity on the turfgrass. Do not apply this product to stressed turf.
- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Do not use adjuvants and spray additive tank mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, [name of registrant] recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

Prohibitions

- Do not apply this product to St. Augustinegrass, bentgrass, greens, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when temperatures are above 90 F; some injury may be expected with spot treatments when air temperatures exceed 90 F. Where state, county, or local governments have more stringent temperature regulations, these regulations must be observed. To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought, or excess rainfall/irrigation, shaded areas, nematodes, improper mowing.

or improper applications of fertilizer and pesticides Injury can occur if this product is applied under any of these or other stress conditions Under any of these stress conditions any turf damage caused by the use of this product is beyond the control of the registrant and all risk is assumed by the buyer and/or user

- For ground application only aerial applications are not permitted
- Do not use this product on or near desirable plants including contact of spray on exposed root systems or adventitious shoots within the dripline of desirable trees and shrubs since injury may result

3 Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator The interaction of many equipment and weather related factors determine the potential for spray drift The applicator and the grower are responsible for considering all these factors when making decisions

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles

Wind Speed

Apply only when the wind speed is 2 to 10 mph at the application site Do not apply at wind speeds greater than 10 mph

Temperature Inversions

If applying at wind speeds less than 3 mph the applicator must determine if 1) conditions of temperature inversion exist or 2) stable atmospheric conditions exist at or below nozzle height Do not make applications into areas of temperature inversions or stable atmospheric conditions

Additional requirements for ground boom application

Do not apply with a nozzle height greater than 4 feet above the crop canopy

4 Spray Preparation and Tank Mixes

In certain applications liquid fertilizer may replace part of the water

Mixing with water

Add one half the required amount of water to the spray tank then add EH 1506Herbicide slowly with agitation and complete filling the tank with water Mix thoroughly and continue agitation while spraying When this product is left standing for extended periods of time re agitate to assure uniformity of the spray mixture

Mixing with liquid fertilizers

Use suitable sources and rates of fertilizer based upon advice of your fertilizer supplier or State Extension Service Specialist

Verify physical compatibility with a jar test Always perform a jar test for compatibility before large scale mixing The jar test can be conducted by mixing all components in a small container in proportionate quantities If the mixture separates after standing and can be mixed readily by shaking then the mixture can be used and applied with spray equipment providing continuous agitation If large flakes sludge gels or other precipitates form or if a separate oily layer or oil globules appear then the herbicide and the liquid fertilizer must not be prepared as a tank mixture

Liquid fertilizers are either solutions (true fluids) or suspensions Mixing this product with suspensions or N P K solutions may not be satisfactory (may be marginal) without pre mixing this product with water Premixing this product with 2 to 4 parts water will ensure that the dispersants enable the herbicide to be suspended in the fertilizer

Adjuvants and spray additives

Adjuvants (such as surfactants spreaders spreader stickers spray thickeners foaming agents activators detergents and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable then adjuvant(s) combined with EH 1506 Herbicide would not be recommended. Do not use adjuvants and spray additive tank mix combinations unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product [name of registrant] recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

5 Ground Equipment

Spray distribution The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles tips and screens are suitable for broadcast applications. For best spray distribution and coverage select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 5 to 220 gallons per acre with spray pressures adjusted to between 20 to 40 psi. Note: For bentgrass (except golf greens) spray volumes use 145 to 220 gallons per acre. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1 000 square feet).

Calibration and proper application are essential when using this product.

- Over application or rates above those specified on this label can cause turf injury.

Hand held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back and forth motion or in a side to side motion or in a swinging arm motion. Instead the nozzle should be held stationary at the proper height. Side to side motion results in uneven coverage.

Hand operated sprayers including backpack sprayers compression sprayers are appropriate for small turfgrass areas.

After using this product clean sprayer with soap or detergent and water or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

6 Application Schedules

Apply this product to broadleaf weeds that are young and actively growing for the best results. Fall applications can provide improved control for emerged winter annuals and perennials such as henbit chickweed clover and ground ivy.

For Turf Applications

Do not apply more than two (2) broadcast treatments of this product per site per year with a minimum of 30 days between applications. A second broadcast application or a follow up application as a spot treatment is advised for more mature weeds for dense infestations and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations or as a follow up treatment or any time broadleaf weeds are actively growing.

For the Listed Non-cropland sites

Use only one (a single) treatment of this product per site per year for woody plant infestations or two broadcast treatments for annual and perennial weeds. It is required to wait 30 days between treatments and a spot treatment may be substituted for the second broadcast treatment if necessary.

Extremes in environmental conditions e.g temperature and moisture soil conditions and cultural practices may affect the activity of this product Under warm moist conditions herbicide symptoms may be accelerated While under very dry conditions the expression of herbicide symptoms is delayed and weeds hardened off by drought are less susceptible to this product

For newly seeded areas

Delay application of this product to grass seedlings until after the second mowing

For newly sodded sprigged or plugged areas

The application of this product to newly sodded sprigged or plugged grasses should be delayed until 3 to 4 weeks after the sodding sprigging or plugging operations

Reseeding interval

Treated areas may be reseeded 3 weeks after application

9

7 How Much to Use

USE RATES AND SPRAY VOLUMES

Generally the lower application rates within the specified range will provide satisfactory control of sensitive weed species The higher application rates within the specified range will be required for dense infestations of perennial weeds for adverse/extreme environmental conditions or for weeds hardened off or more mature

TABLE 1 USE RATES FOR ORNAMENTAL TURF SITES NON CROPLAND AND SOD FARMS		
Species	Rate	Spray Volume
Cool season Turf		
Kentucky bluegrass annual bluegrass annual ryegrass perennial ryegrass tall fescue red or fine leaf fescues	3 0 to 4 Pints/Acre (1 1 to 1 5 fl oz /1 000 sq ft)	5 to 220 Gallons/Acre (0 125 to 5 0 Gallons/1 000 sq ft)
Creeping Bentgrass (excluding golf greens) Apply preferably in May or mid August through September Slight turf yellowing will disappear after about one (1) week Note Care should be taken to avoid overdosing bentgrass or injury may result	1 8 Pints/Acre (0 66 fl oz /1 000 sq ft)	145 to 220 Gallons/Acre (3 33 to 5 0 Gallons/1 000 sq ft) High spray volumes will provide uniform coverage
Warm season Turf		
Hybrid bermudagrass common bermudagrass zoysiagrass buffalograss and bahiagrass Slight turf yellowing will disappear after about one (1) week	3 0 to 4 Pints/Acre (1 1 to 1 5 fl oz /1 000 sq ft) Sod Farms 2 to 2 25 Pints/Acre (0 75 to 0 83 fl oz /1 000 sq ft)	5 to 220 Gallons/Acre (0 125 to 5 0 Gallons/1 000 q ft)
Note Do not apply to above listed warm season turfgrass unless some turf injury can be tolerated It is impossible to test all environmental conditions for the listed warm season turfgrass Do not apply this product to warm season turfgrass during spring green up or in the fall during the transition period between active growth and dormancy Dormant turf This product may be applied to fully dormant bermudagrass fully dormant Zoysiagrass fully dormant buffalograss and fully dormant bahiagrass		

TABLE 2 Restrictions on broadcast applications to ornamental turfgrass sites sod farms and non cropland				
Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum seasonal rate
Ornamental turfgrass sites and Sod farms	4.0 pints/A [1.19 lb MCPA ae/A 0.31 lb MCPP p ae/A 0.11 lb dicamba ae/A]	2	30 days	8.0 pints/A [2.38 lb MCPA ae/A 0.63 lb MCPP p ae/A 0.22 lb dicamba ae/A]

SPOT TREATMENT WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS AND PUMP UP TYPE SPRAYERS)

- Apply any time the emerged broadleaf weeds are actively growing
 - Calibration and proper application are essential when using this product
 - Uniform applications are essential when using this product Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury
 - Hand held techniques Wands fitted with flat fan nozzle tips may be used with the appropriate technique Flat fan nozzles should not be waved in a back and forth motion or in a side to side motion or in a swinging arm motion Instead the nozzle should be held stationary at the proper height Side to side motion results in uneven coverage
 - Follow up applications as spot treatments at a 30 day interval are suitable for more mature weeds for dense infestations and for adverse environmental conditions
- For cool season turfgrass (except Bentgrass) listed in Table 1** Mix 1.1 to 1.5 fl oz of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq ft of turfgrass Apply any time the emerged broadleaf weeds are susceptible
- **For Bentgrass (excluding golf greens)** Mix 0.66 fl oz of this product per four (4.0) gallons of water for treatment of approximately 1,000 sq ft of turfgrass This high spray volume will provide uniform coverage Apply any time the emerged broadleaf weeds are susceptible
 - **For warm season turfgrass listed in Table 1** Mix 0.75 to 0.83 fl oz of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq ft of turfgrass Apply any time the emerged broadleaf weeds are susceptible
 - **Limitations on spot treatments to turfgrass** Spot treatment is defined as a treatment area no greater than 1,000 sq ft per acre The maximum application rate is 1.5 fl oz per 1,000 sq ft per application [0.31 lb MCPP p acid equivalent per acre] The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications

CULTURAL TIPS

Irrigation

- Restriction Do not apply this product through any type of irrigation system
- Rainfast in as little as 8 hours Do not apply this product immediately before rainfall or irrigation
- If possible do not irrigate or water the turfgrass within 8 to 24 hours after application
- If dry conditions exist a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended

Mowing

- For optimum results delay mowing 2 days before and until 2 days after the application of this product

8 Broadleaf Weeds Controlled

EH 1506 Herbicide will control the following broadleaf weeds Apply any time the emerged broadleaf weeds are susceptible

BROADLEAF WEEDS			
annual fleabane	dogfennel	knotweed	red sorrel (sheep sorrel)
aster white heath & white prairie	dovefoot geranium	lambsquarters	roundleaf greenbriar
bedstraw	false dandelion (spotted catsear & common catsear)	lawn burweed	shepherdspurse
beggarticks	field bindweed	lespedeza common	spotted spurge
beggarweed creeping	(morningglory & creeping jenny)	mallow common	spurge
bindweed	field madder	matchweed	sunflower
birdsfoot trefoil	field oxeye-daisy (creeping oxeye)	mouseear chickweed	thistle
black medic	field pennycress	mustard	velvetleaf (buttonweed)
broadleaf plantain	filaree whitestem & redstem	nettle	Venice mallow
buckhorn plantain	Florida pusley	old world diamond flower	Veronica (corn speedwell)
bull thistle	ground ivy	Oxalis (yellow woodsorrel)	Virginia buttonweed
burclover	groundsel	parsley piert	Virginia creeper
burdock common	hairy bittercress	Pennsylvania smartweed	western saisify
Carolina geranium	hawkweed	pennywort (dollarweed)	white clover (Dutch clover
carpetweed	healall	pepperweed	honeysuckle clover white
chickweed common	henbit	pigweed	trefoil & purplewort)
chicory	horsenettle	pineappleweed	wild carrot
cinquefoil	horseweed	plantain	wild garlic
clover	innocence (Blue eyed Mary)	poison ivy	wild geranium
cocklebur	jimsonweed	poison oak	wild lettuce
compassplant	kochia	prostrate knotweed (knotweed)	wild mustard
curly dock		puncturevine	wild onion
dandelion		purple cudweed	wild strawberry
dayflower		purslane	yarrow
deadnettle		ragweed	yellow rocket
Dock		redweed	and many more broadleaf weeds
Synonyms			
A repeat application may be required in 30 days			

9 Spot Treatment in Non Cropland Brush and Weed Control

High volume foliar applications

Apply up to 1 15 gallons of product per 100 gallons of water or apply a 1 0 to 1 15% vol/vol spray solution as a full cover spray with high volume equipment Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard to control species for mature plants during the late summer or under adverse environmental conditions (e g drought)

Spray broadleaf weeds woody plants or mixed brush uniformly and thoroughly by wetting all leaves stems bark and root collars The minimum total volume of spray solution required for adequate coverage of solid stands of mixed brush is 25 gallons of spray solution per treated acre The spray preparation chart for applications on a spray to wet basis is shown below in Table 3

Table 3 Instructions for preparing 25 to 100 gallons of spray solution at 1 0 to 1 15 / spray concentration with water for high volume foliar applications		
Spray solution per acre Gallons	Amount of Product Needed for Spray Concentration of	
	1 0 /	1 15 /
25	2 0 pt	2 3 pt
75	6 0 pt	6 9 pt
100	1 0 gal	1 15 gal
Equal measures 1 gallon = 4 quarts = 8 pints = 128 fl oz		

For Backpack Sprayers and Hand Pump up Sprayers

Table 4 Instructions for preparing 1 to 3 gallons of spray solution at 1 0 to 1 15/ spray concentration with water for high volume foliar applications

Gallons of Water	Amount of Product Needed for Spray Concentration of	
	1 0 /	1 15 /
1	1 25 fl oz (2 5 Tablespoons)	1 5 fl oz (3 Tablespoons)
2	2 5 fl oz	3 0 fl oz
3	3 8 fl oz	4 5 fl oz

Equal measures 1 fl oz = 2 tablespoons (Tbs) = 6 teaspoons (tsp)

Limitations on spot treatments to non cropland Do not apply more than one (1) spot treatment per season Do not apply more than 1 15 gallons of product per acre per year [2 73 lb MCPA acid equivalent 0 72 lb MCPP p acid equivalent and 0 25 lb dicamba acid equivalent per acre per year]

BRUSH [WOODY PLANTS]

Ash	Cedars	Multiflora rose	Sumac
Aspen	Elms	Oak	Willow
Birch	Gooseberry	Poison ivy	And many more brush species
Black cherry	Honey locust	Poison oak	
Brambles	Kudzu	Shortleaf pine	

STORAGE AND DISPOSAL
Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Store in original container in a locked storage area inaccessible to children or pets Keep from freezing

PESTICIDE DISPOSAL Pesticide wastes are acutely hazardous Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law If these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons]
CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying Offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times

[OR

Pressure rinse as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds Drain for 10 seconds after the flow begins to drip]

[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons]

CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying Offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container 1/4 full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Turn the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times

[OR

Pressure rinse as follows Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds Drain for 10 seconds after the flow begins to drip]

[For Refillable Containers]

CONTAINER HANDLING Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose

Container cleaning Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller

To clean the container before final disposal empty the remaining contents from this container into application equipment or a mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times

Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or by other procedures allow by state and local authorities

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product By opening and using this product buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception If the terms are not acceptable return this product unopened immediately to the point of purchase and the purchase price will be refunded in full

It is impossible to eliminate all risks inherently associated with use of this product Damage to the treated article ineffectiveness or other unintended consequences can result from use of the product under abnormal conditions such as weather presence of other materials or the manner of use or application etc Such factors and conditions are beyond the control of the manufacturer and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions warnings and cautions

The manufacturer warrants only that this product conforms to the chemical description given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use subject to the inherent risks described below **TO THE EXTENT CONSISTENT WITH APPLICABLE**

LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE HEREBY EXPRESSLY DISCLAIMED

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER FOR ANY AND ALL LOSSES DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER OR NOT BASED IN CONTRACT NEGLIGENCE STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED AT THE MANUFACTURER S OPTION TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT The Manufacturer must be promptly notified in writing of any claims whether based in contract tort negligence strict liability or otherwise to be eligible to receive either remedy stated above

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner

APPENDIX

1 Statements which may appear on different label components depending on packaging configuration

- See next panel for additional Precautionary Statements and First Aid
 Net Contents _____
 EPA Est. No. _____

2 Advertising claims that may be presented on container labeling advertisements brochures and other marketing/sales promotional materials

- 3 Way Herbicide
 Contains TRIMEC® Herbicide
 Peel here to open [instructions for fix a form label]
- Consumer Access Numbers
 Controls Dandelion Clover Knotweed Henbit Chickweed Plantain Spurge and many other [listed] broadleaf weeds [as listed]
- One Gallon Covers Up To [2.5] Acres
- Water based formula
- Water based formulation
- Rain fast in as little as 8 hours
 Rain fast [Rainproof] in 8 hours
 Consistently controls dandelion spurge clover plantain ground ivy oxalis and [various other listed weeds]
 Controls major [listed] lawn [turf] weeds [as listed]
 Effective post emergent control at a wide range of temperatures (50 to 85 F)
 From the makers of Trimec® herbicides
 Trimec® is a registered trademark of PBI/Gordon Corporation
 A Trimec Complex [LOGO]
- Genuine Trimec Herbicide [LOGO]
 [For] fast dandelion knock down
- For use on Bluegrass Fescues Perennial Ryegrass [Bermudagrass Zoysiagrass (other warm season species listed)]
- For use on cool and warm season turf[grass]

- Economical — lower cost water based formulation
- Economical control of a wide spectrum of broadleaf weeds
- Easy cleanup
- Suitable for applications up to 90 F
- Foliar absorption
- Outstanding broadleaf weed control
 - For use on sod farms
- Labeled for sod farm use
- For use on [sites listed]
- Controls Dandelion Clover Knotweed Henbit Chickweed Plantain Spurge and many other [listed] broadleaf weeds [as listed]
- Lawn applicator formula [Trimec complex]
- Registered for use on [in] cool and warm season grasses
- EH 1506 Herbicide offers these advantages
 - Excellent postemergent activity with proven performance
 - The combinations of these active ingredients provide effective weed control for common and troublesome weed species in turfgrass including dandelion spurge and white clover
 - Rain fast in as little as 8 hours
 - The weed injury symptoms can be noticed within hours of the application and plant death can occur within 14 to 21 days

DOCUMENT CONTROL INFORMATION

1 Unique Label Identifier 002217 00958 20120306 amend proposed highlighted docx

2 Reason for Issue add eyewear for diluted mix