



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 9, 2023

Eric D. Smith
Director
Regulatory Affairs
PBI/Gorgon Corporation
Post Office Box 014090
Kansas City, Missouri 64101

Subject: Registration Review Label Mitigation for Quinclorac
Product Name: EH-1513 ORP
EPA Registration Number: 2217-964
Application Date: January 31, 2019
Decision Number: 592118

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Quinclorac Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

Page 2 of 2
EPA Reg. No. 2217-964
Decision No. 592118

approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a long horizontal flourish extending to the right.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

2,4-D QUINCLORAC DICAMBA	GROUP	4	HERBICIDE
--------------------------------	-------	---	-----------

EH-1513 ORP


EPA Reg. No. 2217-964

ACTIVE INGREDIENT:

2,4-D	0.675%
Dicamba	0.060%
Quinclorac	0.346%
OTHER INGREDIENTS:	<u>98.919%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 13.50 lb 2,4-dichlorophenoxyacetic acid equivalent per ton (CAS 94-75-7)
 - 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton (CAS 1918-00-9)
 - 6.92 lb quinclorac per ton (CAS 84087-01-4)
- Isomer Specific By AOAC Methods.



KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet [inside pages] for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.

OR

See back panel for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.

Guaranteed Analysis: See appendix for example.
FL Fertilizer No. ____

Net Contents: ____
EPA Est. No. ____

Company Name and Address:
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks, and
- waterproof gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

These chemicals (2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

[Product name] is formulated [designed] to feed your lawn and control most annual grasses and broadleaf weeds commonly found in home lawns. The product contains a lawn fertilizer and three different herbicides.

- The three herbicides combine to kill many different kinds of weeds.
- Grass weeds: Postemergent control of crabgrass, foxtails, and other annual grass weeds..
- Broadleaf weeds: Postemergent control of [over] [XXX] broadleaf weeds including dandelion, chickweed, thistle and clover. [A partial list of broadleaf weeds is presented on this label.]

2. Weed Resistance Management

For resistance management, this product contains Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 877-800-5556.

3. Precautions and Restrictions:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.
- EH-1513 ORP is not sold, distributed, or registered within the state of New York. Do not use in the state of New York.

4. Where To Use

[Product name] may be used on the following established turf species:

Established Cool Season Turfgrass: Kentucky bluegrass, rough bluegrass (*Poa trivialis*), perennial ryegrass, fine fescues, tall fescue

Established Warm Season Turfgrass: Bermudagrass, buffalograss, seashore paspalum, zoysiagrass

[Product name] may NOT be used on: annual bluegrass (*Poa annua*), bahiagrass, bentgrass, carpetgrass, centipedegrass, dichondra, St. Augustinegrass. Do not use this product on turfgrass where desirable clovers are present.

[Product name] may be used on warm-season and cool-season turfgrass in 4 use sites.

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs only), and office buildings.
- **Non-cropland sites** include highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations.

Guidelines to follow: The use of [product name] on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

WEEDS CONTROLLED

When applied as directed, this product will control and/or suppress* the following weeds:

Annual Grass Weeds [7]:

Barnyardgrass*	Crabgrass (smooth, southern, large)	Foxtail (green, yellow, giant)
----------------	-------------------------------------	--------------------------------

Broadleaf Weeds [52]: [see appendix for complete list or include here]

Bedstraw	False dandelion	Poison ivy
Beggarticks	Ground ivy	Poison oak
Black medic	Healall	Pokeweed
Broadleaf plantain	Henbit	Purslane, common
Buckhorn plantain	Knotweed	Ragweed
Burclover	Lambsquarters	Red sorrel
Burdock	Lespedeza	Redroot pigweed
Carpetweed	Mallow	Sheep sorrel
Chickweed, common	Matchweed	Shepherdspurse
Chickweed, mouseear	Mustard	Slender plantain
Chicory	Narrowleaf plantain	Smooth dock
Clover, hop	Oxalis	Smooth pigweed
Clover, strawberry	Parsley-piert	Spurge
Clover, white	Pearlwort	Thistle
Curly dock	Pennycress	Woodsorrel
Daisy, oxeye	Pepperweed	Yarrow
Dandelion	Pigweed	
Dichondra	Plantain	

* for acceptable control of barnyardgrass two applications made at a 6 to 10 week interval are needed

5. When To [Apply] [Use]

This product may be applied in the spring, early summer, or fall. Do not apply more than 2 applications per year [season] with at least 30 days between applications.

[Optional additional text for more specific timing for grass weeds]

CRABGRASS (large and smooth):

- **Spring:** For the best results, apply this product from crabgrass emergence until the 6 leaf stage (2 tillers) or before the crabgrass is 3 to 4 inches in height. Generally, this period begins or corresponds to the first time that you can see crabgrass in your lawn.

FOXTAIL (GREEN, GIANT AND YELLOW):

- **Spring and early summer:** For the best results, apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height.

BROADLEAF WEEDS:

- **Spring:** Apply when **dandelion, plantain, buttercup, speedwell, knotweed** and other broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F) for best results.
- **Fall:** Apply when **henbit, chickweed, white clover, ground ivy, lawn burweed** and other broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F) for best results.

[End optional text]

6. How To Apply

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain.
- Poor results will occur if applied to dry lawns. When dry conditions exist, irrigate [or water] the turfgrass 24 hours before application.
- For best results, irrigate after the application. However do not irrigate [or water] sooner than 24 hours after application.
- Apply this product when the wind is calm.
- Do not spread granules onto flowers, vegetables, and ornamental shrubbery.
- Turf species listed on this label can be seeded, sodded, sprigged, or plugged into the treated areas 4 weeks after the application of this product.
- When treating newly planted areas, wait approximately 6 weeks after seedling emergence before applying this product.
- Delay mowing 2 days before and until 3 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

7. How Much To Use

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

[Apply 2.4 to 4.5 pounds of product per 1,000 sq.ft. per application to established turfgrass.] [Apply [2.4] [3.6] [4.5] pounds of product per 1,000 sq. ft. per application to established turfgrass.]

[For products using the 4.5 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 4.5 pounds of product per 1,000 sq.ft. per application [1.32 lbs 2,4-D ae, 0.12 lb dicamba ae, and 0.68 lb quinclorac per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a

minimum of 30 days between applications. The maximum annual (seasonal) rate is 9 pounds of product per 1,000 sq.ft. [2.64 lbs 2,4-D ae, 0.24 lb dicamba ae, and 1.36 lb quinclorac per acre].

[For products using the 3.6 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application [1.06 lbs 2,4-D ae, 0.10 lb dicamba ae, and 0.54 lb quinclorac per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum annual (seasonal) rate is 7.2 pounds of product per 1,000 sq.ft. [2.12 lbs 2,4-D ae, 0.20 lb dicamba ae, and 1.08 lb quinclorac per acre].

[For products using the 2.4 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 2.4 pounds of product per 1,000 sq.ft. per application [0.71 lbs 2,4-D ae, 0.06 lb dicamba ae, and 0.36 lb quinclorac per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum annual (seasonal) rate is 4.8 pounds of product per 1,000 sq.ft. [1.41 lbs 2,4-D ae, 0.13 lb dicamba ae, and 0.72 lb quinclorac per acre].

8. [Spreader Settings] [Coverage] [Distribution]

[Note to Supplemental Registrants: Choose one or more of these application or distribution patterns]

Use the rate listed on this bag. This XX pound bag should be used on XXXX square feet of lawn. A uniform broadcast distribution of product on the treated area is essential. The listed spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage. Never apply by hand.

[Optional: An extended spreader setting listing can be found at [xxx.com]]

Spreader Settings	
Spreader	Setting
Spreader 1	XXX
Spreader 2	XXX

SPREADERS AND CALIBRATION [or SPREADER CALIBRATIONS AND PATTERNS]: Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

- For 4.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 220 square feet and adjust to apply at the rate of 1 pound of product per 220 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet.

SPREADERS AND CALIBRATION:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

Choose one of these application or distribution patterns:

1. **Single Pass** - Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.
[OR Single Pass - Always apply EH1352 Weed & Feed with a properly calibrated rotary or drop type fertilizer spreader. NEVER APPLY BY HAND. To avoid gaps or streaks, first spread border of lawn. Fill in center with uniform, parallel passes. Spreader settings are approximate. Be sure to check spreader before beginning, adjusting if necessary.] To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.]
2. **Double Pass** - Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the

lawn two (2) times. Close the spreader openings when turning to change directions or when stopping.

3. **Cross Hatching** - Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks or the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and cold.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. To avoid wastes, use the entire contents in this bag by application according to the label instructions. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container (bag). Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

[Condition of Sale and Limitation of Warranty and Liability [alternate warranty for 10404]

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of [company name] including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of [company name]. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold [company name] harmless for any claims relating to such factors. [Company name] warrants that this product conforms to the chemical description

on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of [company name], and Buyer and/or User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, [COMPANY NAME] MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOW ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF [COMPANY NAME] FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF [COMPANY NAME] REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL [COMPANY NAME] BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

[Company Name] offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of [company name].]

APPENDIX

1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Use [Apply] where [anytime] weeds are actively growing.
- Kills [Controls] existing crabgrass
- Kills existing weeds, [and feeds your lawn] in one step
- Apply Spring and Fall
- Controls [kills] desired weed species [from attached listing may be inserted here] and other grass and broadleaf weeds [and many other listed broadleaf weeds].
- Control of grass and broadleaf weeds [as listed]!
- 2 in 1: Kills existing grass and broadleaf weeds and fertilizes [in one step]
- Kills [Controls] over XXX grass and broadleaf weeds
- Kills the weeds present in your lawn
- Kills two types of weeds with one product
- Kills all major weeds [as listed]
- Kills major lawn weeds
- Kills dandelion, clover and crabgrass!
- Kills existing weeds
- Kills dandelion, crabgrass and more
- Northern and Southern Grasses, Bluegrass, Fescue, Zoysia, Bermuda, Perennial Ryegrass
- Treats Northern and Southern Grasses
- Kills [Over] [XXX] [+] Weeds [Weed Types], Even [Including] the Roots.
- Especially effective on hard-to-kill weeds
- Kills broadleaf weeds in lawns
- Kills major broadleaf weeds, even the roots.
- Kills dandelion and many other broadleaf weeds.
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion.
- Tough weed killer
- Kills [Destroys] [even] the tough[est] [hard-to-kill] broadleaf weeds [like clover, dandelion, oxalis]
- Kills [major] broadleaf weeds [like clover, dandelion, oxalis]

- Provides broadleaf control
- Controls [Kills] clover, dandelion, plantain, chickweed and [other broadleaf,weeds] [others as listed]
- Clobbers clover
- Kills the root
- Kills weed roots!
- Kills tough lawn weeds
- Consistently fast on: tough weeds like clover, plantain, ground ivy and spurge [various other weeds listed]
- For home lawn care
- Targets weeds, leaves [grasses][the lawn] unharmed
 - Controls Dandelion, White Clover, Plantain, Chickweed, Spurge and Many Other Listed Weeds
 - Kills Weeds While Feeding Your Lawn
 - Kills major broadleaf weeds including roots.
 - Use as directed to kill major broadleaf weeds in your lawn.
 - For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, hybrid bermudagrass
 - Kills Crabgrass and Fertilizes
- Covers [X,XXX] [XX,XXX] square feet [sq ft] [sq. ft.].
- Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding
- Information concerning the raw materials composing this product can be obtained by writing to: [company address], referring to the item number found on this bag.
- Poly Plus is comprised of Polymer Coated Urea
- Feed for 3 months guaranteed!
- Starts working on contact
- Quick greening and extended feeding
- Phosphorus-Free
- Kills over 250 weeds as listed
- Controls a wide spectrum of broadleaf weeds
- Controls crabgrass
- Kills weeds down to the roots
- Kills dandelions and feeds grass
- Builds strong, deep roots
- Kills dandelions and other major lawn weeds
- Kills the weeds not the lawn
- Lawn fertilizer plus weed control
- Promotes root growth for a thicker, greener lawn
- Controls crabgrass escapes
- Contains LESCO® Poly Plus® OPTITM Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.
- Contains LESCO® NOS™ (Nitrogen Optimizing System) for increased nitrogen use efficiency and extended greening.
- Contains LESCO® NOS™ Plus (Nitrogen Optimizing System) for increased nitrogen use efficiency and extended greening.
- Contains LESCO® SulfRX™ Polymer Coated Sulfur Coated Urea to provide a uniform growth with extended nitrogen feeding.

2. Trademark acknowledgement statements:

- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

3. Fertilizer Labeling Requirements in States:

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N)	x.x%
x.x% Ammoniacal Nitrogen	
x.x% Water Insoluble Nitrogen	
x.x% Urea Nitrogen	
x.x% Other Water Soluble Nitrogen	
Available Phosphate (P2O5)	x.x%
Soluble Potash (K2O)	x.x%
Iron (Fe)	x.x%
Nutrient Sources: [Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer.]	

- Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>.
- Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing ____ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.
- If the use rate applies less than 1 pound nitrogen per 1,000 sq.ft. per application, the following statement may appear: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq.ft.
- Advisory for Fertilizers Containing Iron: Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

Broadleaf Weeds: [Controls][Kills][250][more than ###] weeds! (List of weeds may include the following):

Amaranth, purple	Brassbuttons	Cinquefoil
Amaranth, slender	Bristly oxtongue	Clover, crimson
Aster, heath	Broomweed	Clover, alyce
Aster, white heath	Buckhorn	Clover, hop
Aster, purple	Bullnettle	Clover, rabbit foot
Aster, white prairie	Burclover	Clover, red
Artichoke	Burdock	Clover, strawberry
Austrian fieldcress	Burweed, lawn (spurweed)	Clover, yellow sweet
Bedstraw, Catchweed	Buttercup, bulbous	Clover, white sweet
Bedstraw, smooth	Buttercup, creeping	Clover, white
Bedstraw, Northern	Buttonweed, common	Cockle
Bedstraw, pineywoods	(poorjoe)	Cocklebur
Beggarticks, hairy	Buttonweed, virginia	Cocklebur, oriental
Betony, Florida	Carolina geranium (wild)	Cocklebur, spiny
Bindweed, Field	Carpetweed	Common mullein
Bindweed, Hedge	Catnip	Corn Spurry
Bitter Wintercress	Catsear, spotted (false	Creeping Beggarweed
Bittercress, hairy	dandelion)	Creeping Jenny (creeping
Bitterweed	Chamber bitter	charlie)
Black-eyed Susan	Chickweed, common	Cupid shaving Brush
Black Medic	Chickweed, mouseear	Cudweed
Blood flower milkweed	Chickweed, sticky	Daisy, English
Blue lettuce	Chicory	Daisy, oxeye

Dandelion, common
Dandelion, false
Day flower
Deadnettle
Dichondra
Dock, broadleaf
Dock, curly
Dogbane
Dogfennel
Dollarweed (Pennywort)
Dove weed
Elderberry
Falseflax
False sunflower
Fiddleneck
Filaree, redstem
Filaree, whitestem
Fleabane, daisy (annual)
Fleabane, rough
Galinsoga, hairy
Galinsoga, smallflower
Goldenrod
Ground ivy
Groundsel
Gumweed
Hairy fleabane
Hawkweed, meadow
Hawkweed, mouseear
Hawkweed, orange
Hawkweed, yellow
Healall
Heartleaf drymary
Hemp
Henbit
Hoary cress
Horsenettle
Horseweed
Jimsonweed
Johnny-Jumpup violet
Knawel
Knotweed, prostrate
Kochia
Lambsquarters
Lespedeza
Mallow, alkali
Mallow, bristly
Mallow, common
Mallow, dwarf
Mallow, venice
Marcela
Matchweed
Mexicanweed
Moneywort
Morningglory, bigroot
Morningglory, ivyleaf
Morningglory, mexican

Morningglory, red
Morningglory, tall
Morningglory, wolly
Mustard, ball
Mustard, black
Mustard, blue
Mustard, hedge
Mustard, indian,
Mustard, tansy
Mustard, tumble
Mustard, white
Mustard, wormseed
Nettle, stinging (burning)
Nettle, tall
Nettle, wood
Parsley-piert
Parsnip
Pearlwort
Pennycress (frenchweed)
Mugwort
Pepperweed, perennial
Pepperweed, virginia
(Peppergrass)
Pigweed, prostate
Pigweed, redroot
Pigweed, russian
Pigweed, smooth
Pigweed, tumble
Pineapple weed
Plains coreopsis (tickseed)
Plantain, blackseed
Plantain, bracted
Plantain, broadleaf
Plantain, buckhorn
Plantain, hoary
Plantain, narrowleaf
Plantain, slender
Plantain, woolly
Poison ivy
Poison oak
Pokeweed
Prairie Sunflower
Prickly lettuce (compass
plant)
Prickly sida
Prostrate knotweed
Puncturevine
Purslane, common
Pusley, brazil
Pusley, florida
Pusley, large flower
Ragweed, bur
Ragweed, common
Ragweed, lancleaf
Ragweed, western
Redstem filaree

Rough cinquefoil
Scarlet pimpernel
Shepardspurse
Smartweed, ladysthumb
Smartweed, pale
Smartweed, pennsylvania
Smooth chaff-flower
Smoothdock
Sorrel, red (sheep)
Sowthistle, annual
Sowthistle, spiny
Spanish needles
Speedwell, bi-lobed
Speedwell, common
Speedwell, corn
Speedwell, ivyleaf
Speedwell, persian
Speedwell, purslane
Speedwell, slender
Speedwell, snow
Speedwell, thymeleaf
Spiny amaranth
Spurge, spotted
Spurge, prostrate
Spurweed
Star of Bethlehem
Strawberry, India mock
Tansy ragwort
Tanweed
Thistle, blessed
Thistle, bull
Thistle, canada
Thistle, flodman (prairie)
Thistle, musk
Thistle, platte
Thistle, plumeless
Thistle, russian
Thistle, scotch
Thistle, wavyleaf
Thistle, yellow (horrible)
Trailing crownvetch
Velvetleaf
Vervain, blue
Vervain, hoary
Vervain, prostrate
Vervain, tall
Vetch, bird
Vetch, hairy
Vetch, milk
Vetch, narrowleaf
Vetch, wild
Virginia creeper
Western clematis
Western salsify
Wild aster
Wild buckweat

Wild carrot
Wild four-o'clock
Wild lettuce
Wild marigold
Wild mustard (yellow rocket)
Wild parsnip

Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Woodsorrel, creeping
Woodsorrel, violet

Woodsorrel, yellow (Oxalis)
Woolly croton
Wormseed
Yarrow, common
Yarrow, western
Yarrow, yellow

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00964.20230525.amend-proposed-clean.doc

2. Reason for Issue: Registration Review – Quinclorac + Comments & 7/11/19 FTA merger