

US ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave NW Washington D C 20460 EPA Reg Number

Date of Issuance

2217 970

AUG 23 2012

NOTICE OF PESTICIDE

x Registration

__ Reregistration

(under FIFRA as amended)

Term of Issuance unconditional

Name of Pesticide Product

EH 1502 Weed & Feed

Name and Address of Registrant (include ZIP Code)

PBI/Gordon Corporation 1217 West 12th St

Kansas City, MO 64101

Note Changes in labeling differing in substance from that accepted in commercial commercial commercial by the Registration Division prior to use of the labelin commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant the above named pesticide is hereby registered/reregistered under the Federal Insecticide Fungicide and Rodenticide Act Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

The basic and alternate #1 and #2 formulations (dated 2/1/2012) are acceptable and will be added to your file

This product is registered in accordance with FIFRA section 3(c)(5) provided that you

- Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data
- 2 Make the following label revision(s)
 - a Revise EPA Reg No 2217 XXX to EPA Reg No 2217 970
 - b Assure that the EPA establishment number and net contents are also added to the final printed label
 - c Revise ACTIVE INGREDIENT to ACTIVE INGREDIENTS in the ingredient statement
 - d Per the MCPA MCPP p and Dicamba REDs revise the first sentence of the Environmental Hazards section to read This pesticide may be toxic to fish aquatic invertebrates aquatic plants and may adversely affect non target plants
 - e Revise the section heading PRECAUTIONS AND RESTRICTIONS to USE RESTRICTIONS In addition move the text Do not apply directly to or near water when rain is expected that day From the OTHER USE PRECAUTIONS section to the USE RESTRICTIONS section since this is listed as a use restriction in the MCPP p RED
- 3 Within one year of the date on this registration notice the Storage Stability (830 6317) and Corrosion Characteristics (830 6320) studies must be submitted. They must be conducted in the commercial containers of this product and observations should be made at 0 3 6 9 and 12 month intervals.
- 4 Per 40 CFR 156 10(6) submit one copy of your final printed labeling before you release the product for shipment

Continued on Page 2

Signature of Approving Official

Kathryn V Montague Project Manager 23

Herbicide Branch (7505P)
Registration Division (7505P)

Date

AUG 23 2012

EPA Form 8570 6

Page 2 of 2 Notice of Registration 2217 970

If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA sec 6(e) Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice please contact Beth Benbow of my staff at benbow bethany@epa gov

EH-1502 WEED & FEED

EPA Reg No 2217-XXX 970

ACTIVE INGREDIENT MCPA 2 ethylhexyl ester Mecoprop p acid Dicamba acid OTHER INGREDIENTS TOTAL

1 193% 0 174% 0 073%

98 560%

100 000%

WILL COMMENTS In EPA Letter Dated AUG 232012 Under the Federal In

Fungicide and Rodenticide Act as amended for the pesticide registered under EPA Reg No

THIS PRODUCT CONTAINS

15 30 lb 2 methyl 4 chlorophenoxyacetic acid equivalent per ton or 0 765% CAS 94 74 6 3 48 lb (+) (R) 2 (2 methyl 4 chlorophenoxy)propionic acid equivalent per ton or 0 174% CAS 16484 77 8

1 46 lb 3 6 dichloro o anisic acid equivalent per ton or 0 073% CAS 1918 00 9 TRIMEC® is a registered trademark of PBI/Gordon Corporation

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST OBSERVE ALL PRECAUTIONS AND FOLLOW **DIRECTIONS CAREFULLY**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION Causes moderate eye irritation Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating drinking chewing gum using tobacco or using the toilet

First Aid				
If in eyes	Hold eye open and rinse slowly and gently with water for 15 20 minutes Remove contact lenses if present after the first 5 minutes then continue rinsing eye Call a poison control center or doctor for treatment advice			
	uct container or label with you when calling a poison control center or doctor or going for may also contact 1 877 800 5556 for emergency medical treatment information			

Environmental Hazards

This pesticide may adversely affect non target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains drainage ditches gutters or surface waters. Applying his product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway sidewalk or street back onto the treated area of the lawn or garden will help to preven run off to water bodies or drainage systems Do not contaminate water when disposing of equipment washwater or rınsate



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet either directly or through drift Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled
- Not for use on turf being grown for sale or other commercial use as sod or for commercial seed production or for research purposes

INSTRUCTIONS

EH 1502 Weed & Feed is a combination of three proven weed killers. MCPA mecoprop p and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed plantain, henbit, spurge and many others commonly found in home lawns. In addition, a [x x x (grade percentage)] fertilizer provides a supply of plant nutrients.

OTHER USE PRECAUTIONS

Do not use on dichondra carpetgrass St Augustinegrass or turfgrass where desirable clovers are present

Do not apply this product to newly seeded grasses until well established or after the second or third mowing

Do not use this product on flowers vegetables groundcovers gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area

This product will not prevent weed seeds from producing new plants

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs since injury may result

Do not apply directly to or near water storm drains gutters sewers or drainage ditches. Do not apply within 25 feet of rivers fish ponds lakes streams reservoirs marshes estuaries bays and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway sidewalk gutter or street back onto the treated area. To prevent product run off do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day

HOW MUCH TO APPLY

Apply 2 5 to 3 0 pounds of product per 1 000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 3 0 pounds of product per 1 000 sq ft per application [1 0 lb MCPA ae 0 227 lb MCPP p ae and 0 095 lb dicamba ae per acre per application] The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 6 0 pounds of product per 1 000 sq ft per year [2 0 lb MCPA ae 0 455 lb MCPP p acid equivalent, and 0 191 lb dicamba ae per acre per year]

WHEN TO APPLY

Apply when broadleaf weeds are actively growing in spring or early fall

Apply when the air is calm (Choose a calm day) to ensure uniform coverage and avoid spreading granules onto flowers vegetables and ornamental shrubbery

5/9

Possible retreatment may be necessary if lawn is heavily overrun by weeds if adverse low moisture conditions prevail or if weeds are in a state of poor growth

WHAT TO EXPECT

- 1 Mow lawn to normal height 1 to 2 days before application
- Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5)
- 3 Apply when broadleaf weeds are young and actively growing preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water.
- 4 Avoid applying when temperatures are consistently above 90 F
- 5 Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

DISTRIBUTION AND SPREADER PATTERNS

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand

Coverage To determine how much you need to apply multiply the length of your lawn by its width. This equals your total square footage. Deduct all non lawn areas (house gardens etc.) One bag covers up to XXX square feet.

[Note to Supplemental Registrants Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also equipment settings of drop spreaders rotary spreaders and other granular applicators may be presented on the container labels.]

Bag Size Lbs	Application Rate Lbs /1 000 Sq Ft	Treated Area for This Bag Size Sq Ft	Bag Size Lbs	Application Rate Lbs /1 000 Sq Ft	Treated Area for This Bag Size Sq Ft
14	2 5 2 75 3 0	5600 5100 4700	19	2 5 2 75 3 0	7600 6900 6300
15	2 5 2 75 3 0	6000 5450 5000	20	2 5 2 75 3 0	8000 7200 6700
16	2 5 2 75 3 0	6400 5800 5300	25	2 5 2 75 3 0	10000 9000 7500
17	2 5 2 75 3 0	6800 6200 5700	30	2 5 2 75 3 0	12000 11000 10000
18	2 5 2 75 3 0	7200 6500 5400			

SINGLE PASS DISTRIBUTION

For best results apply this product evenly and uniformly avoiding streaking skips or overlaps

- 1 For rotary spreaders operate at intervals corresponding to the swath width For drop spreaders overlap the wheel tracks to ensure complete coverage
- 2 First spread the borders of your lawn and header strips for turning around. Then apply this product to the center with straight line passes.
- 3 Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping

CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution calibrate spreader before application

- For 2 5 pounds of product per 1 000 square feet of turfgrass Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet
- For 2 75 pounds of product per 1 000 square feet of turfgrass. Check your spreader on 360 square feet and adjust to apply at the rate of 1 pound of product per 360 square feet.
- For 3 0 pounds of product per 1 000 square feet of turfgrass. Check your spreader on 330 square feet and adjust to apply at the rate of 1 pound of product per 330 square feet.

Spreader	D	Delivers lb/1 000 sq ft	ft
	2 5 lb	2 75 lb	3 0 lb
LESCO® Rotary calibration gauge setting #	X	X	X
Ortho® Drop (5000)	Х	Х	X
Ortho® Rotary (6000)	X	X	X
Precision® Red DevilO Rotary	X	X	Х
Quaker® Rotary	X	X	X
Republic® EZ Drop	X	X	X
RepublicO EZ Grow	X	X	X
Republic® EZ Rotary	Х	X	X
Scotts® Drop (PF3)	X	X	Х
Scotts® Easy Green	X	X	X
Scotts® Speedy Green	X	X	X
True Temper Rotary	X	X	Х

OR

Spreader settings for 1 0 pound of product per 330 square feet		
Scotts EasyGreen® Rotary	26	
Scotts Drop	5	
Scotts SpeedyGreen® Rotary	3.5	
GreenView Drop (Spred II)	4	
Central Drop (#205)	5.5	
Sears Drop (#19810)	4	
Ortho Drop (#5000)	5	
Precision Drop (Red Devil)	4	

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover sweet	Mugwort	Spiny cocklebur
Aster	Clover white	Musk thistle	Spiny sowthistle
Austrian fieldcress	Cockie	Mustard	Spotted catsear
Bedstraw	Cocklebur	Narrowleaf plantain	Spotted spurge
Beggarticks	Common mullein	Narrowleaf vetch	Spurweed
Betony Florida	Creeping Jenny	Nettle	Stinging nettle
Bindweed field	Cudweed	Orange hawkweed	Strawberr, India mock
Bird vetch	Curly dock	Oxalis	Tall nettle
Bitter wintercress	Daisy English	Parsley piert	Tall vervain
Bittercress hairy	Daisy fleabane	Parsnip	Tansy ragwort
Bitterweed	Daisy oxeye	Pearlwort	Tansy mustard
Black eyed Susan	Dandelion	Pennycress	Thistle
Black medic	Dichondra	Pepperweed	Trailing c owr vetch
Black mustard	Dogfennel	Pigweed	Tumble niustard
Blackseed plantain	Dollarweed	Pineywoods bedstraw	Tumble pigweed

BROADLEAF WEEDS			
Blessed thistle	False dandelion	Plains coreopsis	Velvetleaf
Bloodflower milkweed	Falseflax	Plantain	Venice mallow
Blue lettuce	False sunflower	Poison ivy	Virginia buttonweed
Blue vervain	Fiddleneck	Poison oák	Virginia creeper
Bracted plantain	Florida pusley	Pokeweed	Virginia pepperweed
Bristly oxtongue	Frenchweed	Poorjoe	Wavyleaf bullthistle
Broadleaf dock	Galinsoga	Prairie sunflower	Western clematis
Broadleaf plantain	Goldenrod	Prickly lettuce	Western salsify
Broomweed	Ground ivy	Prickly sida	White mustard
Buckhorn	Gumweed	Prostrate knotweed	Wild aster
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	Wild buckwheat
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild carrot
Bull thistle	Healall	Prostrate vervain	Wild four a clock
Bullnettle	Heartleaf drymary	Puncturevine	Wild garlic
Burclover	Heath aster	Purslane common	Wild geranium
Burdock	Hedge bindweed	Ragweed	Wild lettuce
Burning nettle	Hedge mustard	Red sorrel	Wild marigold
Burweed	Henbit	Redroot pigweed	Wild mustard
Buttercup	Hoary cress	Redstem filaree	Wild onion
Buttonweed	Hoary plantain	Rough cinquefoil	Wild parsnip
Canada thistle	Hoary vervain	Rough fleabane	Wild radish
Carolina geranium	Horsenettle	Russian pigweed	Wild rape
Carpetweed	Jimsonweed	Scarlet pimpernel	Wild strawberry
Catchweed bedstraw	Knawel	Sheep sorrel	Wild sweet potato
Catnıp	Knotweed	Shepherdspurse	Wild vetch
Catsear	Kochia	Slender plantain	Woodsorrel
Chickweed common	Lambsquarters	Smailflower galınsoga	Woolly croton
Chickweed mouseear	Lespedeza	Smooth dock	Woolly morningglory
Chicory	Mallow	Smooth pigweed	Woolly plantain
Cinquefoil	Matchweed	Sorrel	Wormseed
Clover crimson	Mexicanweed	Sowthistle	Yarrow
Clover hop	Milk vetch	Spanishneedles	Yellow rocket
Clover red	Morningglory	Speedwell	Yellowflower pepperweed
Clover strawberry	Mouseear hawkweed	Spiny amaranth	and other broadleaf weeds

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals

PESTICIDE DISPOSAL AND CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container **If empty** Place in trash or offer for recycling if available **If partly filled** Call your local solid waste agency or www website com for disposal instructions. Never place unused product down any indoor or outdoor drain

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED THE MANUFACTURER NEITHER MAKES NOR INTENDS AN / EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHAN ABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE HEREBY EXPRESSLY DISCLAIMED TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO CASE SHALL THE MAINJFACTURER BE LIABLE FOR INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT If these terms are not acceptable return this produc unopened immediately to the point of purchase and the purchase price will be refunded in full. The turms

of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere

APPENDIX

1 Statements which may appear on different label components depending on packaging configuration

- See next panel for additional Precautionary Statements and First Aid
- Net Weight ___
- EPA Est No
- [Guaranteed Analysis and other required fertilizer related statements may be presented on the front panel of the container label. A sample Guaranteed Analysis and possible statements are presented in the Appendix on page 9 of this master label.]
- FL 987

2 Advertising claims that may be presented on container labeling advertisements brochures and other marketing/sales promotional materials

- Controls Tough Broadleaf Weeds in Established Lawns
- Kills Over (number) [Listed] Broadleaf Weeds [As Listed]
- [Bag] [Treats] [Covers] [Up To] [XXXX] Square Feet
- · Greens turf in autumn and accelerates spring recovery

3 Guaranteed Analysis

o Guaranteed Analysis	
GUARANTEED ANALYSIS XX X X	
Total Nitrogen (N)	XX%
x x% ammoniacal nitrogen	
x x% urea nitrogen	
Available Phosphate (P2O5)	X /6
Soluble Potash (K20)	X/
Sulfur (S)	X /6
x x % free sulfur (S)	
Derived from polymer coated sulfur coated urea urea ammonium phosphate potassium chloride A portion of the urea has been coated to provide x x% coated slow release nitrogen This product and/or its manufacture is covered by one or more of the following U S patents and intercounterparts thereof US Patent Numbers xxxxxxx	national

Note Fertilizer grades with nutrient contents expressed as the percentages of N P2O5 and K2O may include the following 23 3 7 or x x x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.

4 Fertilizer Labeling Requirements

- Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm
- Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1 000 square feet per growing season.
 When used as directed this product will supplement your regular fertilizer program by providing ______
 pounds of nitrogen per 1 000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.

5 Advisory For Fertilizers Containing Iron

Keep this product off sidewalks patios driveways etc. This fertilizer may contain iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately

before flushing with water If concrete is wet when contact occurs staining may be impossible to prevent Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete

6 Referrals to other products used sequentially

- For best results apply (brand name) products
- Apply using the 4 Step Program
- 4 Step Program

OR

- STEP 1 Crabgrass Preventor Apply March early May Prevents crabgrass foxtail goosegrass poa annua and other annual weed grasses Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening
- STEP 2 Weed & Feed Apply May June Kills dandelion clover plantain and over XXX other broadleaf weeds Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening
- STEP 3 Lawn Food Apply June August Nourishes your lawn throughout the summer months with extended feeding of Nitrogen Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening
- STEP 4 Winterizer & Weed Control Apply late August mid October Kills dandelion clover plantain and over XXX other broadleaf weeds Maximizes root building Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening
- STEP 4 Starter & Winterizer Apply late August November Maximizes root building Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening. Use with new seeding and sodding of turfgrass.

DOCUMENT CONTROL INFORMATION

- 1 Unique Label Identifier 002217 00XXX 20120201 new proposed EH1502 docx
- 2 Reason for Issue New product