


2217-973

07/18/2012

1/9

 <p align="center">U S ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave , NW Washington, D C 20460</p> <p align="center">NOTICE OF PESTICIDE <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA as amended)</p>	EPA Reg Number 2217-973	Date of Issuance JUL 18 2012
	Term of Issuance unconditional	
	Name of Pesticide Product EH-1524 HERBICIDE	

Name and Address of Registrant (include ZIP Code)
PBI/Gordon Corporation
1217 West 12th St
Kansas City, MO 64101


~~Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.~~

On the basis of information furnished by the registrant the above named pesticide is hereby registered/reregistered under the Federal Insecticide Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment the Administrator on his motion may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

The basic and alternate #1 formulations (dated 3/29/2012) are acceptable and will be added to your file

- This product is registered in accordance with FIFRA section 3(c)(5) provided that you
- 1 Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data
 - 2 Make the following label revision(s)
 - a Revise "EPA Reg No 2217-XXX" to "EPA Reg No 2217-973 "
 - b Assure that the EPA establishment number and net contents are also added to the final printed label
 - c Revise "ACTIVE INGREDIENT" to "ACTIVE INGREDIENTS" in the ingredient statement
 - d Remove "GENERAL" from the section heading, "GENERAL PRECAUTIONS AND RESTRICTIONS"
 - 3 Within one year of the date on this registration notice, the Storage Stability (830 6317) and Corrosion Characteristics (830 6320) studies must be submitted. They must be conducted in the commercial containers of this product and observations should be made at 0, 3, 6, 9, and 12 month intervals
 - 4 Per 40 CFR 156 10(6), submit one copy of your final printed labeling before you release the product for shipment

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. If you have any questions regarding this notice, please contact Beth Benbow of my staff at benbow bethany@epa.gov

Signature of Approving Official Kathryn V Montague Project Manager 23 Herbicide Branch Registration Division (7505P) 	Date JUL 18 2012
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------

EH-1524 HERBICIDE

EPA Reg. No. 2217-XXX

ACTIVE INGREDIENT

2,4-D	0.675%
Dicamba	0.060%
Dithiopyr	0.121%
Quinclorac	0.346%
OTHER INGREDIENTS	<u>98.798%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS

- 13.50 lb 2,4-dichlorophenoxyacetic acid equivalent per ton (CAS 1928-43-4)
 - 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton (CAS 1918-00-9)
 - 2.42 lb dithiopyr per ton (CAS 97886-45-8)
 - 6.92 lb quinclorac per ton (CAS 84087-01-4)
- Isomer Specific By AOAC Methods

**ACCEPTED
with COMMENTS
In EPA Letter Dated.
JUL 18 2012**

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

2217-973

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long sleeved shirt and long pants, shoes plus socks and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by a poison control center or doctor • Do not give anything by mouth to an unconscious person

If on skin or on clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency call toll free 1-877-800-5556</p>	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent run-off to water bodies or drainage systems.

These chemicals (2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1 General Precautions and Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod or for commercial seed or for research purposes.
- EH 1524 Herbicide is not sold, distributed, or registered within the state of New York. Do not use in the state of New York.

Product Description

[Product name] is formulated [designed] to control most annual grasses and broadleaf weeds commonly found in home lawns. The product contains four different herbicides:

- The four herbicides combine to kill many different kinds of weeds at different stages of development.
- Grass weeds: Preemergent control of crabgrass, foxtails, and other annual grass weeds; Postemergent control of crabgrass.
- Broadleaf weeds: Preemergent control of [chickweed, henbit, knotweed, shepherdspurge, spurge, woodsorel, and other] broadleaf weeds; Postemergent control of [over] [XXX] broadleaf weeds including dandelion, chickweed, thistle, and clover. [A partial list of broadleaf weeds is presented on this label.]

2 Where To Use

[Product name] may be used on the following established turf species:

- Established Cool Season Turfgrass:** Kentucky bluegrass, rough bluegrass (*Poa trivialis*), perennial ryegrass, fine fescues, tall fescue, seashore paspalum.
- Established Warm Season Turfgrass:** Bermudagrass, buffalograss, zoysiagrass.

[Product name] may NOT be used on annual bluegrass (*Poa annua*) bahiagrass bentgrass carpetgrass centipedegrass dichondra St Augustinegrass Do not use this product on turfgrass where desirable clovers are present

Guidelines to follow The use of [product name] on lawns and ornamental turfgrasses that are not well established or have been weakened by stress due to mechanical operations chemical injury disease insects pests excessive temperatures or lack of moisture may increase the chances for turfgrass injury Allow turfgrass weakened by these stresses to recover before application of this product

WEEDS CONTROLLED			
When applied as directed this product will control and/or suppress* the following weeds			
Annual Grass Weeds			
Annual bluegrass (<i>Poa annua</i>) Annual ryegrass	Barnyardgrass* Crabgrass (smooth southern large)	Foxtail (green yellow giant)	Goosegrass* (crowfootgrass)
A partial listing of established broadleaf weeds [see appendix for complete list or include here]			
Bedstraw Beggarticks Black medic Broadleaf plantain Buckhorn plantain Burclover Burdock Carpetweed Chickweed common Chickweed mouseear Chicory Clover hop Clover strawberry Clover white	Curly dock Daisy oxeye Dandelion Dichondra False dandelion Ground ivy Healall Henbit Knotweed Lambsquarters Lespedeza Mallow Matchweed Morningglory	Mustard Narrowleaf plantain Oxalis Parsley piert Pearlwort Pennycress Pepperweed Pigweed Plantain Poison ivy Poison oak Pokeweed Purslane common Ragweed	Red sorrel Redroot pigweed Sheep sorrel Shepherdspurse Slender plantain Smooth dock Smooth pigweed Spurge Thistle Woodsorrel Yarrow and other broadleaf weeds
* for acceptable control of barnyardgrass and goosegrass two applications made at a 6 to 10 week interval are needed			

3 When To Apply

This product may be applied as a single application or as split applications in the spring early summer or fall

For annual grasses such as crabgrass (smooth, southern and large) and foxtail (green, yellow, and giant) Apply in the spring and early summer For best results apply this product from 2 weeks before weed germination until the weed is 4 inches tall For dense populations make a second application no sooner than 30 days after the first application

For broadleaf weed control Spring applications for [choose weeds from broadleaf weed list] Fall applications for [choose weeds from broadleaf weed list] For best results apply when broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F)

4 How To Apply

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain
- Poor results will occur if applied to dry lawns When dry conditions exist irrigate [or water] the turfgrass 24 hours before application
- For best results irrigate after the application However do not irrigate [or water] sooner than 24 hours after application
- Apply this product when the wind is calm
- Do not spread granules onto flowers vegetables and ornamental shrubbery

- Turf species listed on this label can be seeded sodded sprigged or plugged into the treated areas 4 weeks after the application of this product
- When treating newly planted areas wait approximately 6 weeks after seedling emergence before applying this product
- Delay mowing 2 days before and until 3 days after the application of this product Additional stress from low mowing heights may increase the possibility of turf injury
- Clippings from the first three mowings should be left on the treated area Do not use clippings from the treated areas as mulch or compost around flowers ornamentals trees or in vegetable gardens

5 How Much To Use

Use a drop or rotary spreader designed to apply granular herbicides Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result

[Apply 2.4 to 4.5 pounds of product per 1 000 sq ft per application to established turfgrass] [Apply [2.4] [3.6] [4.5] pounds of product per 1 000 sq ft per application to established turfgrass]

[For products using the 4.5 lb/1000 sq ft rate] The maximum broadcast application rate is 4.5 pounds of product per 1 000 sq ft per application [1.32 lb 2.4 D ae 0.12 lb dicamba ae 0.24 lb dithiopyr and 0.68 lb quinclorac per acre per application] The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications The maximum annual (seasonal) rate is 9.2 pounds of product per 1 000 sq ft [2.65 lb 2.4-D ae 0.24 lb dicamba ae 0.48 lb dithiopyr and 1.36 lb quinclorac per acre]

[For products using the 3.6 lb/1000 sq ft rate] The maximum broadcast application rate is 3.6 pounds of product per 1 000 sq ft per application [1.06 lb 2.4 D ae 0.09 lb dicamba ae 0.19 lb dithiopyr and 0.54 lb quinclorac per acre per application] The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications The maximum annual (seasonal) rate is 7.2 pounds of product per 1 000 sq ft [2.12 lb 2.4-D ae 0.19 lb dicamba ae 0.38 lb dithiopyr and 1.08 lb quinclorac per acre]

[For products using the 2.4 lb/1000 sq ft rate] The maximum broadcast application rate is 2.4 pounds of product per 1 000 sq ft per application [0.71 lb 2.4 D ae 0.06 lb dicamba ae 0.13 lb dithiopyr and 0.36 lb quinclorac per acre per application] The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications The maximum annual (seasonal) rate is 4.8 pounds of product per 1 000 sq ft [1.42 lb 2.4-D ae 0.12 lb dicamba ae 0.26 lb dithiopyr and 0.72 lb quinclorac per acre]

SUGGESTED SPREADER SETTINGS

[Note to reader A listing of spreaders and suggested settings for the application rate of 2.4 to 4.5 lb per 1 000 sq ft will be presented on the container labels]

The above spreader settings are approximate The age and condition of spreader speed of operation and evenness of terrain may require slightly different settings for desired coverage

CALIBRATION INSTRUCTIONS [examples]

- For 4.5 pounds of product per 1 000 square feet of turfgrass Check your spreader on 220 square feet and adjust to apply at the rate of 1 pound of product per 220 square feet
- For 3.6 pounds of product per 1 000 square feet of turfgrass Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet
- For 2.4 pounds of product per 1 000 square feet of turfgrass Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage and disposal

PESTICIDE STORAGE Keep in original container in a cool dry area away from extreme heat and cold

PESTICIDE DISPOSAL AND CONTAINER HANDLING Nonrefillable container Do not reuse or refill this container (bag) Completely empty bag into application equipment **If empty** Place bag in trash or offer empty bag for recycling if available **If partly filled** Call your local solid waste agency [or [www website com](http://www.website.com)] for disposal instructions Never place unused product down any indoor or outdoor drain

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WHICH ARE HEREBY EXPRESSLY DISCLAIMED TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT If these terms are not acceptable return this product unopened immediately to the point of purchase and the purchase price will be refunded in full The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere

APPENDIX

1 Statements which may appear on different label components depending on packaging configuration

- See next panel for additional Precautionary Statements and First Aid
- Net Weight _____
- EPA Est No _____

2 Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials

- Use [Apply] where [anytime] weeds are actively growing
- Use [Apply] anytime [when] crabgrass [weed] seeds are germinating
- Prevents [Stops] crabgrass and certain broadleaf weeds [weed seed] [from germinating] [for up to XXX months]
- Kills [Controls] existing crabgrass and prevents crabgrass seed germination
- Prevents [crabgrass] [weed seed germination] [and] kills existing weeds in one step
- Apply Spring and Fall
- Controls [kills] desired weed species [from attached listing may be inserted here] and other grass and broadleaf weeds [and many other listed broadleaf weeds]
- Controls major [listed] weeds [as listed] for XX [weeks] [months]
- [Complete] [Total] weed control for XX [weeks] [months]
- Eliminates lawn weeds as listed pre and post emergent control of [crabgrass] [grass and broadleaf weeds]
- Control of grass and broadleaf weeds [as listed]
- Kills [Controls] over XXX grass and broadleaf weeds
- Kills the weeds present in your lawn & prevents more from growing

- Kills two types of weeds with one product
- Kills [listed] major weeds [as listed]
- Kills and prevents major lawn weeds
- Kills and prevents dandelion clover and crabgrass!
- Kills existing weeds & prevents new ones from growing
- Kills dandelion crabgrass and more
- Northern and Southern Grasses Bluegrass Fescue Zoysia Bermuda Perennial Ryegrass
- Treats Northern and Southern Grasses
- Kills [Over] [XXX] [+] Weeds [Weed Types] Even [Including] the Roots
- Especially effective on hard to kill weeds
- Kills broadleaf weeds in lawns
- Kills major broadleaf weeds even the roots And [brand name] is formulated to kill only the weeds not the lawn Use as directed to kill major broadleaf weeds in your lawn
- Kills dandelion and many other broadleaf weeds
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion
- Tough weed killer
- Kills [Destroys] [even] the tough[est] [hard-to kill] broadleaf weeds [like clover dandelion oxalis]
- Kills [major] broadleaf weeds [like clover dandelion oxalis]
- Provides broadleaf control
- Controls [Kills] clover dandelion plantain chickweed and [other broadleaf weeds] [others as listed]
- Clobbers clover
- Kills the root
- Kills weed roots!
- Kills tough lawn weeds
- Consistently fast on tough weeds like clover plantain ground ivy and spurge [various other weeds listed]
- For home lawn care
- For a beautiful lawn
- Kills the weeds not the lawn
- Kills weeds not lawn grasses
- Kills [down to] the root [not the lawn]
- For use on lawns
- Targets weeds leaves [grasses][the lawn] unharmed
- Controls Dandelion White Clover Plantain Chickweed Spurge and Many Other Listed Weeds
- Especially Effective on Hard to-Kill Weeds
- Kills major broadleaf weeds including roots And (brand name) is formulated to kill only the weeds not the lawn Use as directed to kill major broadleaf weeds in your lawn
- For use on Kentucky bluegrass fescue spp perennial ryegrass common bermudagrass zoysiagrass hybrid bermudagrass
- [brand name] Turf Note – Control Those Pesky Weeds
 - Many broadleaf weeds [chickweed henbit] germinate in the fall but are not visible in your lawn until spring Applying (product name) in the fall can provide effective results If needed you can make a follow up application in the spring
- Project Guide – For a thicker greener more beautiful lawn follow (Product Program) It s easy to follow and can be started at anytime by simply applying the product that corresponds with the current season!
- Proper watering and mowing practices along with regular feedings will help promote a beautiful green lawn
- One application does it all - crabgrass dandelion viney weeds even clover
- Covers [up to] [XXXX] sq ft
- Controls [names of weeds from weed list] [and other broadleaf weeds]
- One Application in Spring Provides Control Well into Summer
- Covers [X XXX] [XX XXX] square feet [sq ft] [sq ft]
- Guaranteed [Results] or Your Money Back*

- If you are not satisfied with this product for any reason send your written request for a refund to the address at left together with the original dated store receipt showing the price you paid Please allow 6 to 8 weeks for processing

3 Trademark acknowledgement statements

- DIMENSION® is a registered trademark of Dow AgroSciences L L C
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

Broadleaf Weeds [Controls][Kills][253][more than ###] weeds! (List of weeds may include the following)			
Amaranth purple	Cupid shaving Brush	Mustard ball	Speedwell common
Amaranth slender	Cudweed	Mustard black	Speedwell corn
Aster heath	Daisy English	Mustard blue	Speedwell ivyleaf
Aster white heath	Daisy oxeye	Mustard hedge	Speedwell persian
Aster purple	Dandelion common	Mustard indian	Speedwell purslane
Aster white prairie	Dandelion false	Mustard tansy	Speedwell slender
Artichoke	Day flower	Mustard tumble	Speedwell snow
Austrian fieldcress	Deadnettle	Mustard white	Speedwell thymeleaf
Bedstraw Catchweed	Dichondra	Mustard wormseed	Spiny amaranth
Bedstraw smooth	Dock broadleaf	Nettle stinging (burning)	Spurge spotted
Bedstraw Northern	Dock curly	Nettle tall	Spurge prostrate
Bedstraw pineywoods	Dogbane	Nettle wood	Spurweed
Beggarticks hairy	Dogfennel	Parsley-piert	Star of Bethlehem
Betony Florida	Dollarweed	Parsnip	Strawberry India mock
Bindweed Field	(Pennywort)	Pearlwort	Tansy ragwort
Bindweed Hedge	Dove weed	Pennycress (frenchweed)	Tanweed
Bitter Wintercress	Elderberry	Mugwort	Thistle blessed
Bittercress hairy	Falseflax	Pepperweed perennial	Thistle bull
Bitterweed	False sunflower	Pepperweed virginia	Thistle canada
Black eyed Susan	Fiddleneck	(Peppergrass)	Thistle flodman (prairie)
Black Medic	Filaree redstem	Pigweed prostate	Thistle musk
Blood flower milkweed	Filaree whitestem	Pigweed redroot	Thistle platte
Blue lettuce	Fleabane daisy	Pigweed russian	Thistle plumeless
Brassbuttons	(annual)	Pigweed smooth	Thistle russian
Bristly oxtongue	Fleabane rough	Pigweed tumble	Thistle scotch
Broomweed	Galinsoga hairy	Pineapple weed	Thistle wavyleaf
Buckhorn	Galinsoga smallflower	Plains coreopsis (tickseed)	Thistle yellow (horrible)
Bullnettle	Goldenrod	Plantain blackseed	Trailing crownvetch
Burclover	Ground ivy	Plantain bracted	Velvetleaf
Burdock	Groundsel	Plantain broadleaf	Vervain blue
Burweed lawn (spurweed)	Gumweed	Plantain buckhorn	Vervain hoary
Buttercup bulbous	Hairy fleabane	Plantain hoary	Vervain prostrate
Buttercup creeping	Hawkweed meadow	Plantain narrowleaf	Vervain tall
Buttonweed common	Hawkweed mouseear	Plantain slender	Vetch bird
(poorjoe)	Hawkweed orange	Plantain woolly	Vetch hairy
Buttonweed virginia	Hawkweed yellow	Poison ivy	Vetch milk
Carolina geranium (wild)	Healall	Poison oak	Vetch narrowleaf
Carpetweed	Heartleaf drymary	Pokeweed	Vetch wild
Catnip	Hemp	Prairie Sunflower	Virginia creeper
Catsear spotted (false dandelion)	Henbit	Prickly lettuce (compass plant)	Western clematis
Chamber bitter	Hoary cress	Prickly sida	Western salsify
Chickweed common	Horsenettle	Prostrate knotweed	Wild aster
Chickweed mouseear	Horseweed	Puncturevine	Wild buckweat
Chickweed sticky	Jimsonweed	Purslane common	Wild carrot
Chicory	Johnny Jumpup violet	Pusley brazil	Wild four o'clock
Cinquefoil	Knawel	Pusley florida	Wild garlic
Clover crimson	Knotweed prostrate	Pusley large flower	Wild lettuce
Clover alyce	Kochia	Ragweed bur	Wild marjgold
Clover hop	Lambsquarters	Ragweed common	Wild mustard (yellow rocket)
Clover rabbit foot	Lespedeza	Ragweed lanceleaf	Wild on cr
Clover red	Mallow alkali	Ragweed western	Wild parsnip
	Mallow bristly	Redstem filaree	Wild radish

Broadleaf Weeds [Controls][Kills][253][more than ###] weeds! (List of weeds may include the following)			
Clover strawberry	Mallow common	Rough cinquefoil	Wild rape
Clover yellow sweet	Mallow dwarf	Scarlet pimpernel	Wild strawberry
Clover white sweet	Mallow venice	Shepardspurse	Wild sweet potato
Clover white	Marcela	Smartweed ladysthumb	Wild violet
Cockle	Matchweed	Smartweed pale	Woodsorrel creeping
Cocklebur	Mexicanweed	Smartweed pennsylvania	Woodsorrel violet
Cocklebur oriental	Moneywort	Smooth chaff flower	Woodsorrel yellow (Oxalis)
Cocklebur spiny	Morningglory bigroot	Smoothdock	Woolly croton
Common mullein	Morningglory ivyleaf	Sorrel red (sheep)	Wormseed
Corn Spurry	Morningglory mexican	Sowthistle annual	Yarrow common
Creeping Beggarweed	Morningglory red	Sowthistle spiny	Yarrow western
Creeping Jenny (creeping charlie)	Morningglory tall	Spanish needles	Yarrow yellow
	Morningglory wolly	Speedwell bi lobed	

DOCUMENT CONTROL INFORMATION

1 Unique Label Identifier 002217 00XXX 20120329 new proposed-EH1524 doc

2 Reason for Issue New Registration