

2217-992

2/27/2014

1/10



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

2217-992

Date of Issuance:

FEB 27 2014

NOTICE OF PESTICIDE:

x Registration
Reregistration
(under FIFRA, as amended)

Term of Issuance: unconditional

Name of Pesticide Product:

EH-1523 ORP

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation
1217 West 12th St.
Kansas City, MO 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

- 1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label revision(s):
a. Revise "EPA Reg. No.2217-XXX" to "EPA Reg. No. 2217-992."
b. Assure that the EPA establishment number and net contents are also added to the final printed label.
3. Within one year of the date on this registration notice, the Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies must be submitted. They must be conducted in the commercial containers of this product and observations should be made at 0, 3, 6, 9, and 12 month intervals.
4. Per 40 CFR 156.10(6), submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec.6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. The basic and alternate #1 formulations (dated 9/27/2013) are acceptable and will be added to your file. If you have any questions regarding this notice, please contact Beth Benbow of my staff at benbow.bethany@epa.gov.

Signature of Approving Official:

Kathryn V. Montague
Project Manager 23
Herbicide Branch
Registration Division (7505P)

[Handwritten signature of Kathryn V. Montague]

Date:

FEB 27 2014

EH-1523 ORP

EPA Reg. No. 2217-XXX

ACCEPTED

FEB 27 2014

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registration number 2217-992

ACTIVE INGREDIENTS:

2,4-D acid.....	0.675%
Dicamba acid.....	0.060%
Quinclorac.....	0.294%
OTHER INGREDIENTS:	98.971%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 13.50 lb 2,4-dichlorophenoxyacetic acid equivalent per ton (CAS 1928-43-4)
- 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton (CAS 1918-00-9)
- 5.88 lb quinclorac per ton (CAS 84087-01-4)

Isomer Specific By AOAC Methods.

GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N)	x.x%
x.x% Ammoniacal Nitrogen	
x.x% Water Insoluble Nitrogen	
x.x% Urea Nitrogen	
x.x% Other Water Soluble Nitrogen	
Available Phosphate (P2O5)	x.x%
Soluble Potash (K2O)	x.x%
Iron (Fe)	x.x%
Nutrient Sources: [Guaranteed analysis and nutrient sources to be supplied based on fertilizer analysis used by manufacturer.]	

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.



All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks, and
- chemical- resistant gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
--------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

These chemicals (2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

[Product name] is formulated [designed] to feed your lawn and control most annual grasses and broadleaf weeds commonly found in home lawns. The product contains a lawn fertilizer and three different herbicides.

- The three herbicides combine to kill many different kinds of weeds.
- Grass weeds: Postemergent control of crabgrass, foxtails, and other annual grass weeds.
- Broadleaf weeds: Postemergent control of [over] [XXX] broadleaf weeds including dandelion, chickweed, thistle and clover. [A partial list of broadleaf weeds is presented on this label.]

2. Precautions and Restrictions:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.
- [EH-1523 ORP] is not sold, distributed, or registered within the state of New York. Do not use in the state of New York.
- Do not apply more than 4.5 lbs product/1,000 sq.ft. [1.31 lb 2,4-D ae, 0.12 lb dicamba ae, and 0.57 lb quinclorac per acre] per application
- Do not apply more than 9.0 lbs product/1,000 sq.ft. [2.62 lb 2,4-D ae, 0.24 lb dicamba ae, and 1.14 lb quinclorac per acre] per year.

3. Where To Use

[EH-1523 ORP] may be used on the following established turf species:

Established Cool Season Turfgrass: Kentucky bluegrass, annual bluegrass (*Poa annua*), rough bluegrass (*Poa trivialis*), perennial ryegrass, fine fescues, tall fescue

Established Warm Season Turfgrass: Bermudagrass, buffalograss, seashore paspalum, zoysiagrass

[EH-1523 ORP] may NOT be used on: bahiagrass, bentgrass, carpetgrass, centipedegrass, dichondra, St. Augustinegrass. Do not use this product on turfgrass where desirable clovers are present.

[EH-1523 ORP] may be used on warm-season and cool-season turfgrass in 4 use sites.

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs only), and office buildings.
- **Non-cropland sites** include highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations.

Guidelines to follow: The use of [product name] on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

WEEDS CONTROLLED			
When applied as directed, this product will control and/or suppress* the following weeds:			
Annual Grass Weeds:			
Barnyardgrass*	Crabgrass (smooth, southern, large)	Foxtail (green, yellow, giant)	
A partial listing of established broadleaf weeds: [see appendix for complete list or include here]			
Bedstraw	Curly dock	Mustard	Red sorrel
Beggarticks	Daisy, oxeye	Narrowleaf plantain	Redroot pigweed
Black medic	Dandelion	Oxalis	Sheep sorrel
Broadleaf plantain	Dichondra	Parsley-piert	Shepherdspurse
Buckhorn plantain	False dandelion	Pearlwort	Slender plantain
Burclover	Ground ivy	Pennycress	Smooth dock
Burdock	Healall	Pepperweed	Smooth pigweed
Carpetweed	Henbit	Pigweed	Spurge

Chickweed, common	Knotweed	Plantain	Thistle
Chickweed, mouseear	Lambsquarters	Poison ivy	Woodsorrel
Chicory	Lespedeza	Poison oak	Yarrow
Clover, hop	Mallow	Pokeweed	and other broadleaf
Clover, strawberry	Matchweed	Purslane, common	weeds
Clover, white	Morningglory	Ragweed	
* for acceptable control of barnyardgrass two applications made at a 6 to 10 week interval are needed			

4. When To [Apply] [Use]

This product may be applied in the spring, early summer, or fall. Do not apply more than 2 applications per year [season] with at least 21 days between applications.

[Optional additional text for more specific timing for grass weeds]

CRABGRASS (large and smooth):

- **Spring:** For the best results, apply this product from crabgrass emergence until the 6 leaf stage (2 tillers) or before the crabgrass is 3 to 4 inches in height. Generally, this period begins or corresponds to the first time that you can see crabgrass in your lawn.

FOXTAIL (GREEN, GIANT AND YELLOW):

- **Spring and early summer:** For the best results, apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height.

BROADLEAF WEEDS:

- **Spring:** Apply when **dandelion, plantain, buttercup, speedwell, knotweed** and other broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F) for best results.
- **Fall:** Apply when **henbit, chickweed, white clover, ground ivy, lawn burweed** and other broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F) for best results.

[End optional text]

5. How To Apply

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain.
- Poor results will occur if applied to dry lawns. When dry conditions exist, irrigate [or water] the turfgrass 24 hours before application.
- For best results, irrigate after the application. However do not irrigate [or water] sooner than 24 hours after application.
- Apply this product when the wind is calm.
- Do not spread granules onto flowers, vegetables, and ornamental shrubbery.
- Turf species listed on this label can be seeded, sodded, sprigged, or plugged into the treated areas 4 weeks after the application of this product.
- When treating newly planted areas, wait approximately 6 weeks after seedling emergence before applying this product.
- Delay mowing 2 days before and until 3 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

6. How Much To Use

6/10

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

[Apply 2.4 to 4.5 pounds of product per 1,000 sq.ft. per application to established turfgrass.] [Apply [2.4] [3.6] [4.5] pounds of product per 1,000 sq. ft. per application to established turfgrass.]

[For products using the 4.5 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 4.5 pounds of product per 1,000 sq.ft. per application [1.31 lb 2,4-D ae, 0.12 lb dicamba ae, and 0.57 lb quinclorac per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum annual (seasonal) rate is 9 pounds of product per 1,000 sq.ft. [2.62 lb 2,4-D ae, 0.24 lb dicamba ae, and 1.14 lb quinclorac per acre].

[For products using the 3.6 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application [1.06 lb 2,4-D ae, 0.09 lb dicamba ae, and 0.46 lb quinclorac per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum annual (seasonal) rate is 7.2 pounds of product per 1,000 sq.ft. [2.12 lb 2,4-D ae, 0.18 lb dicamba ae, and 0.92 lb quinclorac per acre].

[For products using the 2.4 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 2.4 pounds of product per 1,000 sq.ft. per application [0.71 lbs 2,4-D ae, 0.06 lb dicamba ae, and 0.31 lb quinclorac per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum annual (seasonal) rate is 4.8 pounds of product per 1,000 sq.ft. [1.41 lbs 2,4-D ae, 0.12 lb dicamba ae, and 0.62 lb quinclorac per acre].

7. [Spreader Settings] [Coverage] [Distribution]

[Note to Supplemental Registrants: Choose one or more of these application or distribution patterns]

Use the rate listed on this bag. This XX pound bag should be used on XXXX square feet of lawn. A uniform broadcast distribution of product on the treated area is essential. The listed spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage. Never apply by hand.

[Optional: An extended spreader setting listing can be found at [xxx.com]]

Spreader Settings	
Spreader	Setting
Spreader 1	XXX
Spreader 2	XXX

SPREADERS AND CALIBRATION [or SPREADER CALIBRATIONS AND PATTERNS]: Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

- For 4.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 220 square feet and adjust to apply at the rate of 1 pound of product per 220 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet.

SPREADERS AND CALIBRATION:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product. Choose one of these application or distribution patterns:

1. **Single Pass** - Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping.
[OR Single Pass - Always apply [EH-1523 ORP] with a properly calibrated rotary or drop type fertilizer spreader. NEVER APPLY BY HAND. To avoid gaps or streaks, first spread border of lawn. Fill in center with uniform, parallel passes. Spreader settings are approximate. Be sure to check spreader before beginning, adjusting if necessary.] To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.]
2. **Double Pass** - Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions or when stopping.
3. **Cross Hatching** - Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks or the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

STORAGE AND DISPOSAL
 Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and cold.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

PAPER AND PLASTIC BAGS
CONTAINER HANDLING STATEMENTS: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight ____

- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Use [Apply] where [anytime] weeds are actively growing.
- Kills [Controls] existing crabgrass
- Kills existing weeds, [and feeds your lawn] in one step
- Apply Spring and Fall
- Controls [kills] desired weed species [from attached listing may be inserted here] and other grass and broadleaf weeds [and many other listed broadleaf weeds].
- Control of grass and broadleaf weeds [as listed]!
- 2 in 1: Kills existing grass and broadleaf weeds and fertilizes [in one step]
- Kills [Controls] over XXX grass and broadleaf weeds
- Kills the weeds present in your lawn
- Kills two types of weeds with one product
- Kills major weeds [as listed]
- Kills major lawn weeds
- Kills dandelion, clover and crabgrass!
- Kills existing weeds
- Kills dandelion, crabgrass and more
- Northern and Southern Grasses, Bluegrass, Fescue, Zoysia, Bermuda, Perennial Ryegrass
- Treats Northern and Southern Grasses
- Kills [Over] [XXX] [+] Weeds [Weed Types], Even [Including] the Roots.
- Especially effective on hard-to-kill weeds
- Kills broadleaf weeds in lawns
- Kills major broadleaf weeds, even the roots.
- Kills dandelion and many other broadleaf weeds.
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion.
- Tough weed killer
- Kills [Destroys] [even] the tough[est] [hard-to-kill] broadleaf weeds [like clover, dandelion, oxalis]
- Kills [major] broadleaf weeds [like clover, dandelion, oxalis]
- Provides broadleaf control
- Controls [Kills] clover, dandelion, plantain, chickweed and [other broadleaf,weeds] [others as listed]
- Clobbers clover
- Kills the root
- Kills weed roots!
- Kills tough lawn weeds
- Consistently fast on: tough weeds like clover, plantain, ground ivy and spurge [various other weeds listed]
- For home lawn care
- Targets weeds, leaves [grasses][the lawn] unharmed
- Controls Dandelion, White Clover, Plantain, Chickweed, Spurge and Many Other Listed Weeds
- Kills Weeds While Feeding Your Lawn
- Kills major broadleaf weeds including roots.
- Use as directed to kill major broadleaf weeds in your lawn.
- For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, hybrid bermudagrass
- Kills Crabgrass and Fertilizes
- Covers [X,XXX] [XX,XXX] square feet [sq ft] [sq. ft.].
- Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding
- Information concerning the raw materials composing this product can be obtained by writing to: [company address], referring to the item number found on this bag.
- Poly Plus is comprised of Polymer Coated Urea

- Feed for 3 months guaranteed!
- Starts working on contact
- Quick greening and extended feeding
- Phosphorus-Free
- Kills over 250 weeds as listed
- Controls a wide spectrum of broadleaf weeds
- Controls crabgrass
- Kills weeds down to the roots
- Kills dandelions and feeds grass
- Builds strong, deep roots
- Kills dandelions and other major lawn weeds
- Kills the weeds not the lawn
- Lawn fertilizer plus weed control
- Promotes root growth for a thicker, greener lawn
- Controls crabgrass escapes

3. Trademark acknowledgement statements:

- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

4. Fertilizer Labeling Requirements in States:

- Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>.
- Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing ___ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.
- If the use rate applies less than 1 pound nitrogen per 1,000 sq.ft. per application, the following statement may appear: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq.ft.
- Advisory for Fertilizers Containing Iron: Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

Broadleaf Weeds: [Controls][Kills][253][more than ###] weeds! (List of weeds may include the following):			
Amaranth, purple	Creeping Jenny (creeping charlie)	Morningglory, red	Sowthistle, spiny
Amaranth, slender	Cupid shaving Brush	Morningglory, tall	Spanish needles
Aster, heath	Cudweed	Morningglory, woolly	Speedwell, bi-lobed
Aster, white heath	Daisy, English	Mustard, ball	Speedwell, common
Aster, purple	Daisy, oxeye	Mustard, black	Speedwell, corn
Aster, white prairie	Dandelion, common	Mustard, blue	Speedwell, ivyleaf
Artichoke	Dandelion, false	Mustard, hedge	Speedwell, persian
Austrian fieldcress	Day flower	Mustard, indian,	Speedwell, purslane
Bedstraw, Catchweed	Deadnettle	Mustard, tansy	Speedwell, slender
Bedstraw, smooth	Dichondra	Mustard, tumble	Speedwell, snow
Bedstraw, Northern	Dock, broadleaf	Mustard, white	Speedwell, thymeleaf
Bedstraw, pineywoods	Dock, curly	Mustard, wormseed	Spiny amaranth
Beggarticks, hairy	Dogbane	Nettle, stinging (burning)	Spurge, spotted
Betony, Florida	Dogfennel	Nettle, tall	Spurge, prostrate
Bindweed, Field	Dollarweed (Pennywort)	Nettle, wood	Spurweed
Bindweed, Hedge	Dove weed	Parsley-piert	Star of Bethlehem
Bitter Wintercress	Elderberry	Parsnip	Strawberry, India mock
Bittercress, hairy		Pearwort	Tansy ragwort
Bitterweed		Pennycress (frenchweed)	Tanweed
Black-eyed Susan		Mugwort	Thistle, blessed

Broadleaf Weeds: [Controls][Kills][253][more than ###] weeds! (List of weeds may include the following):			
Black Medic	Falseflax	Pepperweed, perennial	Thistle, bull
Blood flower milkweed	False sunflower	Pepperweed, virginia	Thistle, canada
Blue lettuce	Fiddleneck	(Peppergrass)	Thistle, flodman (prairie)
Brassbuttons	Filaree, redstem	Pigweed, prostate	Thistle, musk
Bristly oxtongue	Filaree, whitestem	Pigweed, redroot	Thistle, platte
Broomweed	Fleabane, daisy	Pigweed, russian	Thistle, plumeless
Buckhorn	(annual)	Pigweed, smooth	Thistle, russian
Bullnettle	Fleabane, rough	Pigweed, tumble	Thistle, scotch
Burclover	Galinsoga, hairy	Pineapple weed	Thistle, wavyleaf
Burdock	Galinsoga, smallflower	Plains coreopsis (tickseed)	Thistle, yellow (horrible)
Burweed, lawn (spurweed)	Goldenrod	Plantain, blackseed	Trailing crownvetch
Buttercup, bulbous	Ground ivy	Plantain, bracted	Velvetleaf
Buttercup, creeping	Groundsel	Plantain, broadleaf	Vervain, blue
Buttonweed, common	Gumweed	Plantain, buckhorn	Vervain, hoary
(poorjoe)	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Buttonweed, virginia	Hawkweed, meadow	Plantain, narrowleaf	Vervain, tall
Carolina geranium (wild)	Hawkweed, mouseear	Plantain, slender	Vetch, bird
Carpetweed	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Catnip	Hawkweed, yellow	Poison ivy	Vetch, milk
Catsear, spotted (false	Healall	Poison oak	Vetch, narrowleaf
dandelion)	Heartleaf drymary	Pokeweed	Vetch, wild
Chamber bitter	Hemp	Prairie Sunflower	Virginia creeper
Chickweed, common	Henbit	Prickly lettuce (compass plant)	Western clematis
Chickweed, mouseear	Hoary cress	Prickly sida	Western salsify
Chickweed, sticky	Horsenettle	Prostrate knotweed	Wild aster
Chicory	Horseweed	Puncturevine	Wild buckweat
Cinquefoil	Jimsonweed	Purslane, common	Wild carrot
Clover, crimson	Johnny-Jumpup violet	Pusley, brazil	Wild four-o'clock
Clover, alyce	Knawel	Pusley, florida	Wild lettuce
Clover, hop	Knotweed, prostrate	Pusley, large flower	Wild marigold
Clover, rabbit foot	Kochia	Ragweed, bur	Wild mustard (yellow rocket)
Clover, red	Lambsquarters	Ragweed, common	Wild parsnip
Clover, strawberry	Lespedeza	Ragweed, lanceleaf	Wild radish
Clover, yellow sweet	Mallow, alkali	Ragweed, western	Wild rape
Clover, white sweet	Mallow, bristly	Redstem filaree	Wild strawberry
Clover, white	Mallow, common	Rough cinquefoil	Wild sweet potato
Cockle	Mallow, dwarf	Scarlet pimpernel	Woodsorrel, creeping
Cocklebur	Mallow, Venice	Shepherd's purse	Woodsorrel, violet
Cocklebur, oriental	Marcela	Smartweed, ladysthumb	Woodsorrel, yellow (Oxalis)
Cocklebur, spiny	Matchweed	Smartweed, pale	Woolly croton
Common mullein	Mexicanweed	Smartweed, pennsylvania	Wormseed
Corn Spurry	Moneywort	Smooth chaff-flower	Yarrow, common
Creeping Beggarweed	Morningglory, bigroot	Smoothdock	Yarrow, western
	Morningglory, ivyleaf	Sorrel, red (sheep)	Yarrow, yellow
	Morningglory, mexican	Sowthistle, annual	

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00XXX.20140224.new-proposed-clean-EH1523ORP.doc

2. Reason for Issue: EPA Comments #1