



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 11, 2023

Eric D. Smith
Director, Regulatory Affairs
PBI/Gorgon Corporation
P.O. Box 860350
Shawnee, KS 66286

Subject: Registration Review Label Mitigation for Quinclorac
Product Name: EH-1523 ORP
EPA Registration Number: 2217-992
Application Date: January 31, 2019
Decision Number: 592117

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Quinclorac Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently

Page 2 of 2
EPA Reg. No. 2217-992
Decision No. 592117

approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Caleb Carr via email at carr.caleb@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington", with a long horizontal flourish extending to the right.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

2,4-D QUINCLORAC DICAMBA	GROUP	4	HERBICIDE
--------------------------------	-------	---	-----------

EH-1523 ORP


EPA Reg. No. 2217-992

ACTIVE INGREDIENTS:

2,4-D acid.....	0.675%
Dicamba acid.....	0.060%
Quinclorac	0.294%
OTHER INGREDIENTS:	<u>98.971%</u>
TOTAL	<u>100.000%</u>

THIS PRODUCT CONTAINS:

- 13.50 lb 2,4-D acid per ton (CAS 94-75-7)
 - 1.20 lb dicamba acid per ton (CAS 1918-00-9)
 - 5.88 lb quinclorac per ton (CAS 84087-01-4)
- Isomer Specific By AOAC Methods.



KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet [inside pages] for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. [OR]

See back panel for complete [Precautionary Statements, including First Aid, and] Directions for Use, including Storage and Disposal.

Guaranteed Analysis: See appendix for example.

FL Fertilizer No. ____

Net Contents: ____

EPA Est. No. ____

Company Name and Address:

PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks, and
- chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, poly-ethylene, polyvinyl chloride ≥ 14 mils, or Viton ≥ 14 mils.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

2,4-D and dicamba have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

[EH-1523 ORP] is formulated [designed] to feed your lawn and control listed annual grasses and broadleaf weeds commonly found in home lawns. The product contains a lawn fertilizer and three different herbicides.

- The three herbicides combine to kill many different kinds of weeds.
- Grass weeds: Postemergent control of crabgrass, foxtails, and other annual grass weeds.
- Broadleaf weeds: Postemergent control of [over] [XXX] broadleaf weeds including dandelion, chickweed, thistle and clover.

2. Weed Resistance Management

For resistance management, this product contains Group 4 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 877-800-5556.

3. Use Restrictions:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed or for research purposes.
- [EH-1523 ORP] is not sold, distributed, or registered within the state of New York. Do not use in the state of New York.

- Do not apply more than 4.5 lbs product/1,000 sq.ft. (1.31 lb 2,4-D ae, 0.12 lb dicamba ae, and 0.57 lb quinclorac per acre) per application
- Do not apply more than 9.0 lbs product/1,000 sq.ft. (2.62 lb 2,4-D ae, 0.24 lb dicamba ae, and 1.14 lb quinclorac per acre) per year.
- Do not apply more than 2 applications per year with at least 21 days between applications.

4. Where To Use

[EH-1523 ORP] may be used on the following established turf species:

Established Cool Season Turfgrass: Kentucky bluegrass, annual bluegrass (*Poa annua*), rough bluegrass (*Poa trivialis*), perennial ryegrass, fine fescues, tall fescue

Established Warm Season Turfgrass: Bermudagrass, buffalograss, seashore paspalum, zoysiagrass

[EH-1523 ORP] may NOT be used on: bahiagrass, bentgrass, carpetgrass, centipedegrass, dichondra, St. Augustinegrass. Do not use this product on turfgrass where desirable clovers are present.

[EH-1523 ORP] may be used on warm-season and cool-season turfgrass in 4 use sites.

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs only), and office buildings.
- **Non-cropland sites** include highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, roadside ditches, road shoulders, road embankments, dividers and medians; municipal, state and federal lands; airports and military installations.

Guidelines to follow: The use of [EH-1523 ORP] on lawns and ornamental turfgrasses that are not well established, or have been weakened by stress due to mechanical operations, chemical injury, disease, insects, pests, excessive temperatures, or lack of moisture may increase the chances for turfgrass injury. Allow turfgrass weakened by these stresses to recover before application of this product.

WEEDS CONTROLLED			
When applied as directed, this product will control and/or suppress* the following weeds:			
Annual Grass Weeds:			
Barnyardgrass*	Crabgrass (smooth, southern, large)	Foxtail (green, yellow, giant)	
* for acceptable control of barnyardgrass two applications made at a 6 to 10 week interval are needed			
Broadleaf Weeds [248 Weeds]			
Amaranth, purple	Creeping Beggarweed	Morningglory, mexican	Speedwell, bi-lobed
Amaranth, slender	Creeping Jenny	Morningglory, red	Speedwell, common
Amaranth, spiny	(creeping charlie)	Morningglory, tall	Speedwell, corn
Aster, heath	Cupid shaving Brush	Morningglory, woolly	Speedwell, ivyleaf
Aster, white heath	Cudweed	Mugwort	Speedwell, Persian
Aster, purple	Daisy, English	Mustard, ball	Speedwell, purslane
Aster, white prairie	Daisy, oxeye	Mustard, black	Speedwell, slender
Aster, wild	Dandelion, common	Mustard, blue	Speedwell, snow
Artichoke	Day flower	Mustard, hedge	Speedwell, thymeleaf
Austrian fieldcress	Deadnettle	Mustard, Indian,	Spurge, spotted
Bedstraw, Catchweed	Dichondra	Mustard, tansy	Spurge, prostrate
Bedstraw, smooth	Dock, broadleaf	Mustard, tumble	Star of Bethlehem
Bedstraw, Northern	Dock, curly	Mustard, white	Strawberry, India mock

Bedstraw, pineywoods	Dock, smooth	Mustard, wormseed	Tansy ragwort
Beggarticks, hairy	Dogbane	Nettle, stinging (burning)	Tanweed
Betony, Florida	Dogfennel	Nettle, tall	Thistle, blessed
Bindweed, Field	Dollarweed	Nettle, wood	Thistle, bull
Bindweed, Hedge	(Pennywort)	Parsley-piert	Thistle, Canada
Bitter Wintercress	Dove weed	Parsnip	Thistle, flodman (prairie)
Bittercress, hairy	Elderberry	Pearlwort	Thistle, musk
Bitterweed	Falseflax	Pennycress (frenchweed)	Thistle, platte
Black-eyed Susan	False sunflower	Pepperweed, perennial	Thistle, plumeless
Black Medic	Fiddleneck	Pepperweed, Virginia	Thistle, Russian
Blood flower milkweed	Filaree, redstem	(Peppergrass)	Thistle, scotch
Blue lettuce	Filaree, whitestem	Pigweed, prostrate	Thistle, wavyleaf
Brassbuttons	Fleabane, daisy	Pigweed, redroot	Thistle, yellow (horrible)
Bristly oxtongue	(annual)	Pigweed, Russian	Trailing crownvetch
Broomweed	Fleabane, hairy	Pigweed, smooth	Velvetleaf
Bullnettle	Fleabane, rough	Pigweed, tumble	Vervain, blue
Burclover	Galinsoga, hairy	Pineapple weed	Vervain, hoary
Burdock	Galinsoga, smallflower	Plains coreopsis (tickseed)	Vervain, prostrate
Burweed, lawn (spurweed)	Goldenrod	Plantain, blackseed	Vervain, tall
Buttercup, bulbous	Ground ivy	Plantain, bracted	Vetch, bird
Buttercup, creeping	Groundsel	Plantain, broadleaf	Vetch, hairy
Buttonweed, common	Gumweed	Plantain, buckhorn	Vetch, milk
(poorjoe)	Hawkweed, meadow	Plantain, hoary	Vetch, narrowleaf
Buttonweed, Virginia	Hawkweed, mouseear	Plantain, narrowleaf	Vetch, wild
Carolina geranium (wild)	Hawkweed, orange	Plantain, slender	Virginia creeper
Carpetweed	Hawkweed, yellow	Plantain, woolly	Western clematis
Catnip	Healall	Poison ivy	Western salsify
Catsear, spotted (false	Heartleaf drymary	Poison oak	Wild buckweat
dandelion)	Hemp	Pokeweed	Wild carrot
Chamber bitter	Henbit	Prairie Sunflower	Wild four-o'clock
Chickweed, common	Hoary cress	Prickly lettuce (compass plant)	Wild garlic
Chickweed, mouseear	Horsenettle	Prickly sida	Wild lettuce
Chickweed, sticky	Horseweed	Puncturevine	Wild marigold
Chicory	Jimsonweed	Purslane, common	Wild mustard (yellow rocket)
Cinquefoil	Johnny-Jumpup violet	Pusley, Brazil	Wild onion
Cinquefoil, rough	Knawel	Pusley, Florida	Wild parsnip
Clover, alyce	Knotweed, prostrate	Pusley, large flower	Wild radish
Clover, crimson	Kochia	Ragweed, bur	Wild rape
Clover, hop	Lambsquarters	Ragweed, common	Wild strawberry
Clover, rabbit foot	Lespedeza	Ragweed, lanceleaf	Wild sweet potato
Clover, red	Mallow, alkali	Ragweed, western	Wild violet
Clover, strawberry	Mallow, bristly	Scarlet pimpernel	Woodsorrel, creeping
Clover, yellow sweet	Mallow, common	Shepard's purse	Woodsorrel, violet
Clover, white sweet	Mallow, dwarf	Smartweed, ladysthumb	Woodsorrel, yellow (Oxalis)
Clover, white	Mallow, venice	Smartweed, pale	Woolly croton
Cockle	Marcela	Smartweed, Pennsylvania	Wormseed
Cocklebur	Matchweed	Smooth chaff-flower	Yarrow, common
Cocklebur, oriental	Mexicanweed	Sorrel, red (sheep)	Yarrow, western
Cocklebur, spiny	Moneywort	Sowthistle, annual	Yarrow, yellow
Common mullein	Morningglory, bigroot	Sowthistle, spiny	
Corn Spurry	Morningglory, ivyleaf	Spanish needles	

5. When To [Apply] [Use]

This product may be applied in the spring, early summer, or fall. Do not apply more than 2 applications per year with at least 21 days between applications.

[Optional additional text for more specific timing for grass weeds]

CRABGRASS (large and smooth):

- **Spring:** For the best results, apply this product from crabgrass emergence until the 6 leaf stage (2 tillers) or before the crabgrass is 3 to 4 inches in height. Generally, this period begins or corresponds to the first time that you can see crabgrass in your lawn.

FOXTAIL (GREEN, GIANT AND YELLOW):

- **Spring and early summer:** For the best results, apply this product to foxtail during the early growth stages or when foxtail is 1 to 4 inches in height.

BROADLEAF WEEDS:

- **Spring:** Apply when **dandelion, plantain, buttercup, speedwell, knotweed** and other broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F) for best results.
- **Fall:** Apply when **henbit, chickweed, white clover, ground ivy, lawn burweed** and other broadleaf weeds are young and actively growing with adequate soil moisture and favorable temperatures (50° to 80°F) for best results.

[End optional text]

6. How To Apply

- This product needs to stick to the leaves of broadleaf weeds and should be applied when the grass is wet from dew or rain.
- Poor results will occur if applied to dry lawns. When dry conditions exist, irrigate [or water] the turfgrass 24 hours before application.
- For best results, irrigate after the application. However do not irrigate [or water] sooner than 24 hours after application.
- Apply this product when the wind is calm.
- Do not spread granules onto flowers, vegetables, and ornamental shrubbery.
- Turf species listed on this label can be seeded, sodded, sprigged, or plugged into the treated areas 4 weeks after the application of this product.
- When treating newly planted areas, wait approximately 6 weeks after seedling emergence before applying this product.
- Delay mowing 2 days before and until 3 days after the application of this product. Additional stress from low mowing heights may increase the possibility of turf injury.
- Clippings from the first three mowings should be left on the treated area. Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

7. How Much To Use

Use a drop or rotary spreader designed to apply granular fertilizers. Avoid skips or excessive overlaps as poor weed control or turfgrass injury may result.

[Apply 2.4 to 4.5 pounds of product per 1,000 sq.ft. per application to established turfgrass.] [Apply [2.4] [3.6] [4.5] pounds of product per 1,000 sq. ft. per application to established turfgrass.]

[For products using the 4.5 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 4.5 pounds of product per 1,000 sq.ft. per application (1.31 lb 2,4-D ae, 0.12 lb dicamba ae, and 0.57 lb quinclorac per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum annual rate is 9 pounds of product per 1,000 sq.ft. (2.62 lb 2,4-D ae, 0.24 lb dicamba ae, and 1.14 lb quinclorac per acre).

[For products using the 3.6 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 3.6 pounds of product per 1,000 sq.ft. per application (1.06 lb 2,4-D ae, 0.09 lb dicamba ae, and 0.46 lb quinclorac per acre per application). The maximum number of broadcast applications is limited to 2 per year with a

minimum of 21 days between applications. The maximum annual rate is 7.2 pounds of product per 1,000 sq.ft. (2.12 lb 2,4-D ae, 0.18 lb dicamba ae, and 0.92 lb quinclorac per acre).

[For products using the 2.4 lb/1000 sq.ft. rate:] The maximum broadcast application rate is 2.4 pounds of product per 1,000 sq.ft. per application (0.71 lbs 2,4-D ae, 0.06 lb dicamba ae, and 0.31 lb quinclorac per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 21 days between applications. The maximum annual rate is 4.8 pounds of product per 1,000 sq.ft. (1.41 lbs 2,4-D ae, 0.12 lb dicamba ae, and 0.62 lb quinclorac per acre).

8. [Spreader Settings] [Coverage] [Distribution]

[Note to Supplemental Registrants: Choose one or more of these application or distribution patterns]

Use the rate listed on this bag. This XX pound bag should be used on XXXX square feet of lawn. A uniform broadcast distribution of product on the treated area is essential. The listed spreader settings are approximate. The age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage. Never apply by hand.

[Optional: An extended spreader setting listing can be found at [xxx.com]]

Spreader Settings	
Spreader	Setting
Spreader 1	XXX
Spreader 2	XXX

SPREADERS AND CALIBRATION [or SPREADER CALIBRATIONS AND PATTERNS]: Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

- For 4.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 220 square feet and adjust to apply at the rate of 1 pound of product per 220 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 2.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 420 square feet and adjust to apply at the rate of 1 pound of product per 420 square feet.

SPREADERS AND CALIBRATION:

Drop spreaders and rotary spreaders should be calibrated for uniform application of this product.

Choose one of these application or distribution patterns:

1. **Single Pass** - Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions or when stopping. To avoid gaps and streaks, first spread border of lawn and then fill in center with uniform, parallel passes. Walk at a steady pace so as to apply evenly throughout the lawn.
[OR **Single Pass** - Always apply [EH-1523 ORP] with a properly calibrated rotary or drop type fertilizer spreader. NEVER APPLY BY HAND. To avoid gaps or streaks, first spread border of lawn. Fill in center with uniform, parallel passes. Spreader settings are approximate. Be sure to check spreader before beginning, adjusting if necessary.]
2. **Double Pass** - Reduce the application rate by one-half of the specified rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions or when stopping.
3. **Cross Hatching** - Reduce the application rate by one-half of the specified rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks or

the spreader pattern of the first trip. Close the spreader opening when turning to change directions or when stopping.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep in original container in a cool, dry area, away from extreme heat and cold.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

[PAPER AND PLASTIC BAGS]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or other procedures approved by State and local authorities.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Use [Apply] where [anytime] weeds are actively growing.
- Kills [Controls] existing crabgrass
- Kills existing weeds, [and feeds your lawn] in one step
- Apply Spring and Fall
- Controls [kills] desired weed species [from attached listing may be inserted here] and other grass and broadleaf weeds [and many other listed broadleaf weeds].
- Control of grass and broadleaf weeds [as listed]!
- 2 in 1: Kills existing grass and broadleaf weeds and fertilizes [in one step]
- Kills [Controls] over XXX grass and broadleaf weeds
- Kills the weeds present in your lawn
- Kills two types of weeds with one product
- Kills major weeds [as listed]
- Kills major lawn weeds
- Kills dandelion, clover and crabgrass!
- Kills existing weeds
- Kills dandelion, crabgrass and more
- [For use on] Northern and Southern Grasses, Bluegrass, Fescue, Zoysia, Bermuda, Perennial Ryegrass
- Treats Northern and Southern Grasses
- Kills [Over] [XXX] [+] Weeds [Weed Types], Even [Including] the Roots.
- Effective on hard-to-kill weeds
- Kills broadleaf weeds in lawns
- Kills major broadleaf weeds, even the roots.
- Kills dandelion and many other broadleaf weeds.
- Use on Lawns to Kill the Toughest Weeds like Clover and dandelion.
- Tough weed killer
- Kills [Destroys] [even] the tough[est] [hard-to-kill] broadleaf weeds [like clover, dandelion, oxalis]
- Kills [major] broadleaf weeds [like clover, dandelion, oxalis]
- Provides broadleaf control
- Controls [Kills] clover, dandelion, plantain, chickweed and [other broadleaf weeds as listed] [others as listed]
- Clobbers clover
- Kills the root
- Kills weed roots!
- Kills tough lawn weeds
- Consistently fast on: tough weeds like clover, plantain, ground ivy and spurge [various other weeds listed]
- For home lawn care
- Targets weeds, leaves [grasses][the lawn] unharmed
 - Controls Dandelion, White Clover, Plantain, Chickweed, Spurge and Many Other Listed Weeds
 - Kills Weeds While Feeding Your Lawn
 - Kills major broadleaf weeds including roots.
 - Use as directed to kill major broadleaf weeds in your lawn.
 - For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, zoysiagrass, hybrid bermudagrass
 - Kills Crabgrass and Fertilizes
- Covers [X,XXX] [XX,XXX] square feet [sq ft] [sq. ft.].
- Contains LESCO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding

- Information concerning the raw materials composing this product can be obtained by writing to: [company address], referring to the item number found on this bag.
- Poly Plus is comprised of Polymer Coated Urea
- Feed for 3 months guaranteed!
- Starts working on contact
- Quick greening and extended feeding
- Phosphorus-Free
- Kills over 250 weeds as listed
- Controls a wide spectrum of broadleaf weeds
- Controls crabgrass
- Kills weeds down to the roots
- Kills dandelions and feeds grass
- Builds strong, deep roots
- Kills dandelions and other major lawn weeds
- Kills the weeds not the lawn
- Lawn fertilizer plus weed control
- Promotes root growth for a thicker, greener lawn
- Controls crabgrass escapes
- www.ecgrow.com
- Q3® is a registered trademark of PBI/Gordon Corporation.
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

2. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “2.1 State Restrictions”]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] Typically a lawn will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing ___ pounds of nitrogen per 1,000 square feet. Consult your local Extension office for questions regarding a fertilizer program suitable to your geographical region.

- If the use rate applies less than 1 pound nitrogen per 1,000 sq.ft. per application, the following statement may appear: Supplemental applications of lawn fertilizer are needed to achieve a minimum of 1 pound nitrogen per 1,000 sq.ft.
- [Advisory for Fertilizers Containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer [contains iron which] may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- California, Oregon, Washington: “[When applied as directed, this products meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.]Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00992.20230621.amend-accepted-clean.doc

2. Reason for Issue: 12/5/2019 SAL + Reg Review Changes from 6/12/23 SAL. Merging the RD & PRD label versions.