

2217-995

6/10/2014

V/14



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

2217-995

Date of Issuance:

JUN 10 2014

NOTICE OF PESTICIDE:

[x] Registration

[] Reregistration

(under FIFRA, as amended)

Term of Issuance: unconditional

Name of Pesticide Product:

EH 1529 Herbicide

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, MO 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

- 1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following label revisions:
a. Revise "EPA REG. NO.2217-xxx" to "EPA REG. NO. 2217-995."
b. Assure that the establishment number and net contents are also added to the final printed label.
4. Submit one (1) copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

If you have any questions regarding this notice, please contact Beth Benbow of my staff at 703-347-8072.

Signature of Approving Official:

Kathryn V. Montague
Project Manager 23
Herbicide Branch
Registration Division (7505P)

[Handwritten signature of Kathryn V. Montague]

Date:

JUN 10 2014

2/14

ACCEPTED
JUN 10 2014
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2217-995

EH-1529 HERBICIDE

EPA Reg. No. 2217-XXX

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

Not for sale, distribution or use in Nassau or Suffolk Counties in New York State.

ACTIVE INGREDIENTS:

Penoxsulam	0.45%
Triclopyr BEE, butoxyethyl ester	4.07%
Sulfentrazone	0.67%
2,4-D, 2-ethylhexyl ester	29.71%
Dicamba acid	2.25%
OTHER INGREDIENTS:	<u>62.85%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 0.04 lb Penoxsulam per gallon or 0.45%.
 - 0.26 lb 3,5,6-trichloro-2-pyridinyloxyacetic acid equivalent per gallon or 2.93%.
 - 0.06 lb N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.67%.
 - 1.75 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 19.70%.
 - 0.20 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 2.25%.
- Isomer specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals

CAUTION: Harmful if swallowed. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are barrier laminate, nitrile rubber, neoprene rubber, and Viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.



All mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks
- chemical-resistant gloves (except for applicators using ground boom equipment) and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical information.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

These chemicals (triclopyr, 2,4-D and dicamba) have properties and characteristics associated with chemicals detected in groundwater. The use of these chemicals in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

4/14

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is :

- coveralls,
- chemical-resistant gloves made of any water-proof material,
- chemical-resistant footwear,
- protective eyewear, and
- chemical-resistant headgear for overhead exposure.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

1. Spray Drift Management

When this product is used in commercial sod production, the following Best Management Practices for reducing spray drift apply.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Additional requirements for ground boom application: Do not apply with a nozzle height greater than 4 feet above the crop canopy.

2. Product Description

Designed for turfgrass applications, EH-1529 Herbicide is a post-emergent foliar-applied herbicide containing multiple modes of action that provides a broad spectrum of weed control. EH-1529 Herbicide is absorbed via leaves, shoots and roots and begins to work immediately by translocating throughout the entire plant to achieve effective control of annual and perennial broadleaf weeds including clover, ground ivy, Virginia buttonweed, dandelion, plantain, dollarweed, Florida betony, yellow nutsedge, English daisy, wild violet, ground ivy, oxalis, wild blackberry, spurge, thistles, yellow nutsedge and many hard to kill weeds. The unique formulation ceases weed growth within hours after application and provides rapid visual response yellowing and desiccation for complete control in cool-season and warm-season turf types listed.

EH-1529 Herbicide controls weeds by affecting multiple sites within the broadleaf weeds. The symptoms of susceptible broadleaf weeds include leaf and stem curl or twisting, and weed yellowing.

EH-1529 Herbicide offers these advantages:

- Excellent postemergent activity with proven performance for some of the toughest broadleaf weeds in turfgrass.
- This product exhibits improved cool-weather performance.
- High selectivity (turfgrass safety) in established cool-season turfgrass and select warm-season turfgrass.
- EH-1529 Herbicide provides rapid and effective weed control for common and troublesome (tough) weed species in turfgrass, including: dandelion, spurge and white clover.
- The speed of action (rate of weed phytotoxicity [yellowing]) and the early weed symptoms are features of EH-1529 Herbicide. Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 10 to 14 days.
- The unique combination provides effective weed control for common and troublesome weed species in turfgrass, such as dollarweed (pennywort).
- This product is rainfast in as little as 3 hours.

6/14

3. Spray Preparation And Tank Mixtures

EH-1529 Herbicide is an aqueous suspo-emulsion (SE) that can be diluted with water or liquid fertilizer to form a stable emulsion.

Mixing with water:

Add one-half the required amount of water to the spray tank, then add EH-1520 Herbicide slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying.

When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Mixing with liquid fertilizers:

Use suitable sources and rates of fertilizer based upon guidance of your fertilizer supplier or State Extension Service Specialist.

Verify physical compatibility with a jar test: Always perform a jar test for compatibility before large scale mixing. The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids) or suspensions. Physical compatibility of this product is adequate with liquid nitrogen solutions. Mixing this product with suspensions or N-P-K solutions may not be satisfactory (may be marginal) without pre-mixing this product with water. Premixing this product with 2 parts water will ensure that the emulsifiers are activated enabling the herbicide to be suspended in the fertilizer.

Adjuvants and spray additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then adjuvant(s) combined with EH-1520 Herbicide would not be advised. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury.

4. Ground Equipment

Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 10 to 220 gallons per acre with spray pressures adjusted to between 20 to 40 psi. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essential when using this product.
- Over-application or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.

Hand operated sprayers including backpack sprayers, compression sprayers are appropriate for small turfgrass areas.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

5. Where To Use

This product provides broadleaf weed control in the following sites:

• **Ornamental Turfgrass sites:**

- **Residential/domestic sites** are defined as turfgrass established around areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental Turf sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
- **Non-cropland sites:** including farmyards, fencerows or fence lines; highway rights-of-way (principal, interstate, county, private, and unpaved roads); roadsides, road shoulders, road embankments, dividers, and medians; industrial sites, lumberyards, tank farms, fuel or equipment storage areas; municipal, state, and federal lands; airports and military installations; telephone rights-of-way, pipeline rights-of-way.
- **Agricultural site:** Commercial sod production

Prohibitions of Sites:

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (salt water bays). Do not apply to wetlands (swamps, bogs, potholes, and marshes).
- Do not apply to greens and tees established on golf courses.
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks or canals.
- Do not allow livestock to graze on any areas treated with this product.

Prohibitions:

- Do not apply this product to St Augustinegrass, creeping bentgrass, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when ambient temperatures are above 90°F. Some injury may be expected with spot treatments when air temperatures exceed 90°F.
- For ground application only. Aerial applications are not permitted.
- Do not collect grass clippings for use as mulch around plants.
- Do not apply this product through any type of irrigation.
- Do not apply this product to bare ground or paved surfaces.

STATE RESTRICTIONS:

- **Arizona:** The state of Arizona has not approved this product for use on sod farms.
- **New York:** Not for sale, distribution or use in Nassau or Suffolk Counties in New York State.
- **California:** Make broadcast applications only between March 1 and September 1. If troublesome weeds appear during other times of the year, a spot application can be made. While irrigation is necessary and important for plant growth, apply irrigation water efficiently so that no more than 125% of the net irrigation requirement is applied for any irrigation event. Apply efficient irrigations for six months following application of sulfentrazone containing products. Do not apply product to bare ground.

8/14

6. Application Schedules

Apply this product to broadleaf weeds that are young and actively growing for the best results. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective. Fall applications provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

For the Listed Residential/domestic sites, Ornamental Turf sites, Institutional sites and Agricultural sites:

Do not apply more than 2 broadcast treatments of this product per site per year. A second broadcast application or a follow-up application as a spot treatment is suggested for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

For the Listed Non-cropland sites:

For more mature weeds or for adverse environmental conditions, a follow-up application as a spot treatment or additional broadcast application may be required.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

Apply up to 2 broadcast treatments of this product per site per year for annual and perennial weeds. For woody plants, one post-emergent application is permitted per year. Wait 30 days between treatments.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to this product.

For Newly Seeded Areas:

Delay application of this product to grass seedlings until after the second or third mowing.

For Newly Sodded, Sprigged, or Plugged Areas:

The application of this product should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Reseeding interval:

Treated areas may be reseeded 4 weeks after application.

Turfgrass tolerance:

- Turfgrass tolerance to this product may vary, and temporary turfgrass yellowing may occur on listed warm-season turfgrass (see Table 1).
- Tolerant turf species listed on this label may exhibit temporary turf injury. The best tolerance occurs under optimal conditions for the turfgrass. Adverse environmental conditions may reduce the selectivity on the turfgrass. Injury may occur under marginal conditions (e.g. low temperatures and drought stress) or under extreme conditions (e.g. high temperatures and high humidity). To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, low soil pH, nematodes, improper mowing or improper applications of fertilizer and pesticides. Under any of these stress conditions, to the extent consistent with applicable law, any turf damage caused by the use of this product is beyond the control of PBI/Gordon Corporation and all risk is assumed by the buyer and/or user.
- Certain spray tank additives (adjuvants, wetting agents, and surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Use

adjuvants and spray additives or tank-mix combinations only when your experience indicates that the tank mixture will not result in objectionable turf injury

- Do not broadcast apply this product when temperatures are above 90°F; some injury can also be expected with spot treatments when air temperatures exceed 90°F.

7. How Much To Use

USE RATES AND SPRAY VOLUMES:

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

TABLE 1. USE RATES FOR ORNAMENTAL TURFGRASS, SOD FARMS, AND NON-CROPLAND.		
Species	Rate	Spray Volume
Cool-season Turf		
Kentucky bluegrass, annual bluegrass, and annual ryegrass	3.25 to 4 Pints/Acre (1.2 to 1.5 fl.oz./1,000 sq.ft.)	10 to 220 Gallons/Acre (0.25 to 5.0 Gallons/1,000 sq.ft.)
Perennial ryegrass, tall fescue, red or fine leaf fescues	2 pints/Acre (0.75 fl.oz./1,000 sq.ft.)	10 to 220 Gallons/Acre (0.25 to 5.0 Gallons/1,000 sq.ft.)
Warm-season Turf		
Hybrid Bermudagrass, common Bermudagrass, buffalograss, centipedegrass, seashore paspalum, kikuyugrass, zoysiagrass, and bahiagrass	2 to 2.25 Pints/Acre (0.75 to 0.83 fl.oz./1,000 sq.ft.)	10 to 220 Gallons/Acre (0.25 to 5.0 Gallons/1,000 sq.ft.)
Dormant turf: This product may be applied to fully dormant warm-season turfgrass.		
Note: Do not apply to above listed warm-season turfgrass unless turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass. We suggest testing this product on a small area and observe the treated area for 30 days to determine the acceptability of turf discoloration.		
Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy.		
If treating Kentucky bluegrass or annual bluegrass in a mixed stand with perennial ryegrass, tall fescue, or red/fine leaf fescues, use 2 pints/acre unless some injury to the sensitive species can be tolerated.		
EH-1529 Herbicide should only be applied to turfgrass species that are listed in Table 1 unless trial use indicates that the turf species not listed is tolerant to EH-1529 Herbicide.		

Limitations on broadcast treatments for ornamental turfgrass, sod farms, and non-cropland:

The maximum application rate is 4 pints of product per acre per application [0.88 lb 2,4-D ae, 0.13 lb triclopyr ae, 0.10 lb dicamba ae, 0.03 lb sulfentrazone, and 0.02 lb Penoxsulam per acre]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8 pints of product per acre [1.75 lb 2,4-D ae, 0.26 lb triclopyr ae, 0.20 lb dicamba ae, 0.06 lb sulfentrazone, and 0.04 lb penoxsulam per acre].

SPOT TREATMENT WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS AND PUMP-UP TYPE SPRAYERS):

- Apply any time the emerged broadleaf weeds are actively growing. Spot treatments can be applied at the dosage rates equal to the specific dosage rates for broadcast treatments.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury.

10/14

- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.
- Follow-up applications as spot treatments at a 30 day interval are suggested for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **For cool-season turfgrass listed in Table 1:** Mix 1.2 to 1.5 fl.oz. of this product per 1 gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **For warm-season turfgrass listed in Table 1:** Mix 0.75 to 0.83 fl.oz. of this product per 1 gallon of water for treatment of approximately 1,000 sq.ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.

CULTURAL TIPS

Irrigation:

- Do not apply this product through any type of irrigation system.
- Rainfast in as little as 3 hours. Do not apply this product immediately before rainfall or irrigation.
- If possible, do not irrigate or water the turfgrass within 3 to 24 hours after application.
- If dry conditions exist, schedule irrigation or watering 24 hours before and 24 hours after application.

Mowing:

- Delay mowing 2 days before and until 2 days after the application of this product. Do not collect grass clippings for use as mulch around plants.

8. Broadleaf Weeds Controlled

EH-1529 Herbicide will control or suppress the following broadleaf. Apply any time the emerged broadleaf weeds are susceptible.

Broadleaf Weeds			
Aster, white heath & white prairie	Dock	Knotweed	Puncturevine
Bedstraw	Dogfennel	Lambsquarters	Purple cudweed
Beggarweed, creeping	Dollarweed (*pennywort)	Lawn burweed	Purslane
Bindweed	Doveweed	Lespedeza, common	Ragweed
Black medic	English Daisy	Mallow, common	Red sorrel (*sheep sorrel)
Broadleaf plantain	False dandelion (*spotted catsear & common catsear)	Matchweed	Shepherd's purse
Buckhorn plantain	Field bindweed	Mouseear chickweed	Speedwell (Veronica)
Bull thistle	(*morningglory & creeping jenny)	Mustard	Spurge
Burdock, common	Field oxeye-daisy (*creeping oxeye)	Nettle	Thistle
Buttercup, creeping	Filaree, whitestem & redstem	Nutsedge (yellow)	Virginia buttonweed
Carpetweed	Florida betony	Old world diamond flower	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)
Catnip	Florida pusley	Oxalis (*yellow woodsorrel & creeping woodsorrel)	Wild carrot
Chickweed	Ground ivy	Parsley-piert	Wild garlic
Chicory	Groundsel	Pennsylvania smartweed	Wild geranium
Cinquefoil	Hawkweed	Pennycress, field	Wild lettuce
Clover	Healall	Pepperweed	Wild mustard
Compassplant (*prickly lettuce)	Henbit	Pigweed	Wild onion
Cudweed	Innocence (Blue-eyed Mary)	Pineappleweed	Wild strawberry
Curly dock		Plantain	Wild violet**
Dandelion		Poison ivy	Yarrow
Dayflower		Poison oak	Yellow rocket
Deadnettle		Prickly lettuce (*compass plant)	

* Synonyms
 ** For best results, apply in the spring when wild violets are blooming or apply a late fall application followed by a spring application.

11/14

9. For Use In Non-Cropland: Brush And Weed Control

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Spot treatment or individual woody plant treatment in noncropland: Apply to trees and brush when foliage is fully expanded and plants are actively growing. Spray broadleaf weeds, woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars.

Spot treatments can be applied at rates equal to the broadcast rate up to 4 pints of product per acre. The spray volume will depend on the density, height and type of brush (woody plants). For a total spray volume of 100 gallons, mix 4 pints of products with 100 gallons of water or prepare a 0.5% spray solution (vol/vol). Refer to Table 3 for quick-mix instructions for the preparation of spray concentrations up to 2.5%

Table 3. Instructions for preparing 100 to 400 gallons of spray solution up to 2.5% spray concentration with water.

Spray solution per acre, Gallons	Amount of Product Needed for Spray Concentration of:					
	0.13%	0.17%	0.25%	0.50%	1.0%	2.5%
1 gal	---	---	---	0.64 fl.oz.	1.2 fl.oz.	3.2 fl.oz.
10 gal	1.5 fl.oz.	2 fl.oz.	3 fl.oz.	6 fl.oz.	12 fl.oz.	32 fl.oz.
20 gal	3 fl.oz.	4 fl.oz.	6 fl.oz.	12 fl.oz.	25 fl.oz.	4 pints
50gal	8 fl.oz.	10 fl.oz.	16 fl.oz.	32 fl.oz.	4 pints	---
100 gal	16 fl.oz.	20 fl.oz.	32 fl.oz.	4 pints	---	---
200 gal	32 fl.oz.	42 fl.oz.	4 pints	---	---	---
300 gal	48 fl.oz.	4 pints	---	---	---	---
400 gal	4 pints	---	---	---	---	---

Equal measures: 1 gallon = 4 quarts = 8 pints = 128 fl.oz. 1 fl.oz. = 2 tablespoons (Tbs) = 6 teaspoons (tsp)

Use a single application for rights-of-way, including electrical power lines, communication lines, pipelines, highways and railroads that intersect wooded areas or stands of trees, brush and woody plants. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g. drought).

BRUSH CONTROLLED:			
Ash	Cedar	Honey locust (suppression)	Sumac
Aspen	Cherry (except Black Cherry)	Honeysuckle	Sycamore
Beech		Kudzu	Trumpet creeper
Birch	Cottonwood	Multiflora Rose	Wild Grape
Blackberry	Dogwood	Oak	Wild Plum
Black Locust	Elm	Poison Ivy	Willow
Brambles	Gooseberry	Poison Oak	
Buckbrush	Hawthorn	Sycamore	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency,

or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

[Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[For Refillable Containers:]

CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

~~THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.~~ The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Tough Weed Control
- Tough Weed LOGO
- Tough Weed Formulation
- Targets [tough], [triclopyr] weeds

14/14

- Contains triclopyr, a proven ingredient for hard-to-control weeds
- To control some of your toughest [triclopyr] weeds
- Consistently fast control on tough weeds – wild violet, ground ivy, oxalis, thistles, dandelion, spurge, clover, plantain and [various other listed weeds].
- Rainfast in as little as 3 hours
- Rainfast [Rainproof] in 3 hours
- [weed name listed in the Directions for Use of this label] control in as little as one week
- Proven performance
- From the makers of Trimec® herbicides.
- Trimec® is a registered trademark of PBI/Gordon Corporation.
- For information call XXX-XXX-XXXX [contact www.xxx-xxxx.com]
- For use on: Bluegrass, Fescues, Perennial and Annual Ryegrass, [Bermudagrass, Zoysiagrass] [and other turf species listed]
- Fast visual response
- Weed control — FAST!
- Tough Weeds Controlled -- Fast!
- Suitable for applications up to 90°F
- Foliar absorption
- Reduces call backs
- Low odor
- Low-odor formulation
- PROFORM® and TRIMEC® are registered trademarks of PBI/Gordon Corporation.
- ® Checkered Flag/Label Design is a registered trademark of PBI/Gordon Corporation.



DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00XXX.20140514.new-proposed-EH1529-clean
2. **Reason for Issue:** EPA Comments #1