UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

June 14, 2016

James L. Kunstman, Ph.D. VP, Regulatory Services PBI/Gordon Corporation P. O. Box 014090 Kansas City, MO 64101

Subject: Label Amendment – Update Storage Statement/State Fertilizer Language/Ad Claims/Typo Corrections Product Name: EH-1558 Weed and Feed EPA Registration Number: 2217-997 Application Date: December 8, 2015 Decision Number: 511996

Dear Dr. Kunstman:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2 EPA Reg. No. 2217-997 Decision No. 511996

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Terri Stowe by phone at (703) 305-6117, or via email at stowe.terri@epa.gov.

Sincerely,

Vaytryn V. Wontaguo

Kathryn V. Montague Product Manager 23 Herbicides Branch Registration Division (7505P) Office of Pesticide Programs

Enclosure

#### NOTIFICATION

#### 2217-997

# EH-1558 Weed and Feed [EH-1414 WEED & FEED with MA]

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/14/2016

# EPA Reg. No. 2217-997

[Note to Reviewer: Label includes residential sites and outdoor domestic use with applications by homeowners]

| ACTIVE INGREDIENTS:       |          |
|---------------------------|----------|
| 2,4-D, 2-ethylhexyl ester | 1.108%   |
| Mecoprop-p acid           | 0.167%   |
| Dicamba acid              | 0.071%   |
| OTHER INGREDIENTS:        | 98.654%  |
| TOTAL                     | 100.000% |

#### THIS PRODUCT CONTAINS:

14.70 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%. CAS 1928-43-4.

3.34 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%. CAS 16484-77-8.

1.42 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.071%. CAS 1918-00-9 Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/Gordon Corporation.

| GUARANTEED ANALYSIS XX-X-X   |        |
|--|--------|
| Total Nitrogen (N)   | XX%    |
| x.x% ammoniacal nitrogen   |        |
| x.x% urea nitrogen*  |        |
| Available Phosphate (P2O5)   | X%     |
| Soluble Potash (K20)   | X%     |
| Sulfur (S)   | X%     |
| x.x% free sulfur (S)   |        |
| *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.<br>This product and/or its manufacture is covered by one or more of the following U.S. patents and internation<br>counterparts thereof: US Patent Numbers: xxxxx<br>Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may inclu  | de the |
| following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental register the states of the st |        |
| ALSO CONTAINS NON PLANT FOOD INGREDIENTS.<br>SOIL AMENDING GUARANTEED ANALYSIS<br>Active Fungal Ingredients:<br>Mycorrhizal Fungi1.12 props/gm   |        |
| Glomus intraradices       0.28 props/gm         Glomus mosseae       0.28 props/gm         Glomus etunicatum       0.28 props/gm         Glomus aggregatum       0.28 props/gm   |        |
| Inert Ingredients (as a soil amendment):<br>XX.XXX% Fertilizer ingredients<br>X.XXX% Herbicide ingredients   |        |



### **KEEP OUT OF REACH OF CHILDREN**

## CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

### **PRECAUTIONARY STATEMENTS**

#### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long sleeved shirt, long pants, shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

| First Aid  |  |
|--|--|
| If in eyes:  | <ul> <li>Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.<br/>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul> |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information. |  |

#### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

#### [Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use on dichondra, carpetgrass, 'Floratam; St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.
- Do not apply this product to newly seeded grasses until well-established or after the second or third mowing.
- Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the dripline of, or over surface roots of trees, shrubs, and other ornamentals.
- Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy.

| This herbicide controls weeds listed in the following table: |                        |                             |                            |
|--|------------------------|-----------------------------|----------------------------|
| Amaranth, purple   | Cupid shaving Brush    | Mustard, ball               | Speedwell, common          |
| Amaranth, slender  | Cudweed                | Mustard, black              | Speedwell, corn            |
| Aster, heath   | Daisy, English         | Mustard, blue               | Speedwell, ivyleaf         |
| Aster, white heath   | Daisy, oxeye           | Mustard, hedge              | Speedwell, persian         |
| Aster, purple  | Dandelion, common      | Mustard, Indian,            | Speedwell, purslane        |
| Aster, white prairie   | Dandelion, false       | Mustard, tansy              | Speedwell, slender         |
| Artichoke  | Day flower             | Mustard, tumble             | Speedwell, snow            |
| Austrian fieldcress  | Deadnettle             | Mustard, white              | Speedwell, thymeleaf       |
| Bedstraw, Catchweed  | Dichondra              | Mustard, wormseed           | Spiny amaranth             |
| Bedstraw, smooth   | Dock, broadleaf        | Nettle, stinging (burning)  | Spurge, spotted            |
| Bedstraw, Northern   | Dock, curly            | Nettle, tall                | Spurge, prostrate          |
| Bedstraw, pineywoods   | Dogbane                | Nettle, wood                | Spurweed                   |
| Beggarticks, hairy   | Dogfennel              | Parsley-piert               | Star of Bethlehem          |
| Betony, Florida  | Dollarweed             | Parsnip                     | Strawberry, India mock     |
| Bindweed, Field  | (Pennywort)            | Pearlwort                   | Tansy ragwort              |
| Bindweed, Hedge  | Dove weed              | Pennycress (frenchweed)     | Tanweed                    |
| Bitter Wintercress   | Elderberry             | Pepperweed, perennial       | Thistle, blessed           |
| Bittercress, hairy   | Falseflax              | Pepperweed, virginia        | Thistle, bull              |
| Bitterweed   | False sunflower        | (Peppergrass)               | Thistle, canada            |
| Black-eyed Susan   | Fiddleneck             | Pigweed, prostrate          | Thistle, flodman (prairie) |
| Black Medic  | Filaree, redstem       | Pigweed, redroot            | Thistle, musk              |
| Blood flower milkweed  | Filaree, whitestem     | Pigweed, russian            | Thistle, platte            |
| Blue lettuce   | Fleabane, daisy        | Pigweed, smooth             | Thistle, plumeless         |
| Brassbuttons   | (annual)               | Pigweed, tumble             | Thistle, russian           |
| Bristly oxtongue   | Fleabane, rough        | Pineapple weed              | Thistle, scotch            |
| Broomweed  | Galinsoga, hairy       | Plains coreopsis (tickseed) | Thistle, wavyleaf          |
| Buckhorn   | Galinsoga, smallflower | Plantain, blackseed         | Thistle, yellow (horrible) |
| Bullnettle   | Goldenrod              | Plantain, bracted           | Trailing crownvetch        |
| Burclover  | Ground ivy             | Plantain, broadleaf         | Velvetleaf                 |
| Burdock  | Groundsel              | Plantain, buckhorn          | Vervain, blue              |
| Burweed, lawn (spurweed)                                     | Gumweed                | Plantain, hoary             | Vervain, hoary             |
| Buttercup, bulbous   | Hairy fleabane         | Plantain, narrowleaf        | Vervain, prostrate         |
| Buttercup, creeping  | Hawkweed, meadow       | Plantain, slender           | Vervain, tall              |
| Buttonweed, common   | Hawkweed, mouseear     | Plantain, woolly            | Vetch, bird                |
| (poorjoe)  | Hawkweed, orange       | Poison ivy                  | Vetch, hairy               |
| Buttonweed, virginia   | Hawkweed, yellow       | Poison oak                  | Vetch, milk                |
| Carolina geranium (wild)                                     | Healall                | Pokeweed                    | Vetch, narrowleaf          |
| Carpetweed   | Heartleaf drymary      | Prairie Sunflower           | Vetch, wild                |

This herbicide controls weeds listed in the following table:

| Catnip                   | Hemp                  | Prickly lettuce (compass plant) | Virginia creeper             |
|--------------------------|-----------------------|---------------------------------|------------------------------|
| Catsear, spotted (false  | Henbit                | Prickly sida                    | Western clematis             |
| dandelion)               | Hoary cress           | Prostrate knotweed              | Western salsify              |
| Chamber bitter           | Horsenettle           | Puncturevine                    | Wild aster                   |
| Chickweed, common        | Horseweed             | Purslane, common                | Wild buckweat                |
| Chickweed, mouseear      | Jimsonweed            | Pusley, brazil                  | Wild carrot                  |
| Chickweed, sticky        | Johnny-Jumpup violet  | Pusley, florida                 | Wild four-o-clock            |
| Chicory                  | Knawel                | Pusley, large flower            | Wild garlic                  |
| Cinquefoil               | Knotweed, prostrate   | Ragweed, bur                    | Wild lettuce                 |
| Clover, crimson          | Kochia                | Ragweed, common                 | Wild marigold                |
| Clover, alyce            | Lambsquarters         | Ragweed, lancleaf               | Wild mustard (yellow rocket) |
| Clover, hop              | Lespedeza             | Ragweed, western                | Wild onion                   |
| Clover, rabbit foot      | Mallow, alkali        | Redstem filaree                 | Wild parsnip                 |
| Clover, red              | Mallow, bristly       | Rough cinquefoil                | Wild radish                  |
| Clover, strawberry       | Mallow, common        | Scarlet pimpernel               | Wild rape                    |
| Clover, yellow sweet     | Mallow, dwarf         | Shepherd's purse                | Wild strawberry              |
| Clover, white sweet      | Mallow, venice        | Smartweed, ladysthumb           | Wild sweet potato            |
| Clover, white            | Marcela               | Smartweed, pale                 | Wild violet                  |
| Cockle                   | Matchweed             | Smartweed, pennsylvania         | Woodsorrel, creeping         |
| Cocklebur                | Mexicanweed           | Smooth chaff-flower             | Woodsorrel, violet           |
| Cocklebur, oriental      | Moneywort             | Smoothdock                      | Woodsorrel, yellow (Oxalis)  |
| Cocklebur, spiny         | Morningglory, bigroot | Sorrel, red (sheep)             | Woolly croton                |
| Common mullein           | Morningglory, ivyleaf | Sowthistle, annual              | Wormseed                     |
| Corn Spurry              | Morningglory, mexican | Sowthistle, spiny               | Yarrow, common               |
| Creeping Beggarweed      | Morningglory, red     | Spanish needles                 | Yarrow, western              |
| Creeping Jenny (creeping | Morningglory, tall    | Speedwell, bi-lobed             | Yarrow, yellow               |
| charlie)                 | Morningglory, wolly   |                                 |                              |
|                          | Mugwort               |                                 |                              |

#### **INSTRUCTIONS:**

EH-1414 Weed & Feed with MA is a combination of three proven weed killers--2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, and spurge. In addition, a x-x-x [(grade percentage)] fertilizer provides a supply of plant nutrients.

#### **USE PRECAUTIONS:**

- Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.
- Reseeding or over seeding areas may occur three (3) weeks after the application of this product.

#### HOW MUCH TO APPLY [WHERE TO USE]:

[For use on: Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, bahiagrass, zoysiagrass, St. Augustinegrass, centipedegrass, hybrid bermudagrass, and colonial bentgrass.]: Apply 2.5 pounds of product per 1,000 square feet of turfgrass. If needed, a second application may be made after 30 days or more. The maximum number of broadcast applications per treatment site is 2 per year. Do not apply more than 5 pounds of product per 1,000 sq.ft. per year. [Note to Reviewer: The following information may be omitted from the final printed label: 2.5 lbs. of this product contains 0.018 lb 2,4-D ae, 0.042 lb mecoprop-p ae, and 0.018 lb dicamba ae.]

Typically a lawn utilizes [xx] pounds of nitrogen per 1,000 sq.ft. per growing season. When applied as directed, this product provides [yy] pounds of nitrogen per 1,000 sq.ft. Contact your local Extension Service for information specific to your geographic region.

#### WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

Apply when the air is calm (Choose a calm day) to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

#### **APPLICATION TIPS [APPLICATION INSTRUCTIONS]:**

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
- 3. Apply when broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
- 4. Avoid applying when temperatures are consistently above 90°F.
- 5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

# [SPREADER SETTINGS] [DISTRIBUTION AND SPREADER PATTERNS] [HOW TO APPLY - SPREADERS AND CALIBRATION]:

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

**Coverage** - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers [up to] XXX square feet.

IMPORTANT: It is important to follow directions as indicated on this quality lawn care product as overapplication may cause injury to your lawn. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.]

| Bag Size, Lbs. | Size, Lbs. Application Rate, Lbs./1,000 Sq.Ft. Treated Area for This Bag Size, Sq.Ft. |        |
|----------------|---|--------|
| 12.5           | 2.5   | 5,000  |
| 18.75          | 2.5   | 7,500  |
| 37.5           | 2.5   | 15,000 |

#### SINGLE PASS DISTRIBUTION:

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

#### CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS:

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.

• For 2.5 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 400 square feet and adjust to apply at the rate of 1 pound of product per 400 square feet.

**[BROADCAST SPREADERS:** For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to ensure complete coverage.]

**[DROP SPREADERS:** Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.]



[Optional Spreader Charts]

| Spreader                                  | Setting Delivers 2.5 lbs./1,000 sq.ft. |
|---|--|
| Scotts® Turf Builder® Classic Drop        | X                                      |
| Scotts® Turf Builder® Edgeguard® Mini     | Х                                      |
| Sta-Green® Broadcast                      | Х                                      |
| Pennington® Broadcast                     | Х                                      |
| LESCO® Rotary calibration gauge setting # | Х                                      |
| Ortho® Drop (5000)                        | Х                                      |
| Ortho® Rotary (6000)                      | Х                                      |
| Precision® Red Devil® Rotary              | Х                                      |
| Quaker® Rotary                            | Х                                      |
| Republic® EZ Drop                         | Х                                      |
| Republic® EZ Grow                         | Х                                      |
| Republic® EZ Rotary                       | Х                                      |
| Scotts® Drop (PF3)                        | Х                                      |
| Scotts® Easy Green                        | Х                                      |
| Scotts® Speedy Green                      | Х                                      |
| True Temper Rotary                        | Х                                      |

#### SETTING YOUR LAWN SPREADER:

THESE SETTINGS ARE A GUIDE. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch flow rate. If fertilizer appears to be coming out either slower or faster than you are covering your lawn, adjust the setting. Close the spreader when filling, turning or stopping.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a cool, dry, locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

#### **Alternate Warranty and Disclaimer**

WARRANTY & DISCLAIMER: To the extent consistent with applicable law, [company name] guarantees this fine product to the extent of the purchase price, only when used according to directions.

## APPENDIX FOR ALL SUBLABELS

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

# 2. Advertising claims and marketing phrases that may be presented on promotional materials and the container labels of supplemental registrants.

General Marketing Claims:

- [For use on residential lawns]
- If product is to be sold in California, add the following statement: [NOTICE This product contains chemicals known to the State of California to cause birth defects or other reproductive harm.]
- The Pennington® UltraGreen™ Difference

Pennington® UltraGreen<sup>™</sup> product line: Half of the nitrogen in this premium line is stabilized<sup>++</sup>, which allows grass to use more of the nutrients you put down and reduces nutrient loss to the environment. The other half of the nitrogen is slow-release for extended feeding for [up to] 3 months. Additionally, these unique fertilizer blends contain mycorrhizal fungi, natural organisms that have been present in vigorous soils for millions of years. These microscopic fungi increase the surface absorbing area of roots, greatly improving the ability of the lawn to access vital soil resources such as water and nutrients.

These tiny fertilizer granules are packed with essential plant nutrients and fungi, so you can feed more lawn with less fertilizer.

- <sup>++</sup> Excluding the Pennington Ultragreen Starter Fertilizer
- Benefits of Pennington® UltraGreen™ [Product Name]
- Promotes thick, green grass
- Half of the total nitrogen in this product is slow-release and feeds for [up to] 3 months
- Half of the total nitrogen is stabilized for less leaching into the environment
- Phosphorus-free
- Pennington® UltraGreen<sup>™</sup> [Product Name] WILL NOT BURN when used as directed. We're so confident you'll love the results of using Pennington® UltraGreen<sup>™</sup> [Product Name] that it comes with the Pennington Satisfaction Guarantee<sup>†</sup>.

† Pennington Satisfaction Guarantee: If, after following the instructions, you are not satisfied with the product results, send your name, address, and proof of purchase (original dated receipt) to: Pennington Refund

1000 Parkwood Circle, Suite 700 Atlanta, GA 30339

• Pennington® UltraGreen™ [Product Name] controls a wide variety of existing broadleaf weeds while feeding your grass.

Herbicide Benefit Claims:

- Starts working on contact
- Kills over [XXX] [Listed] weeds including henbit, dollar weed, white clover, plantain and chickweed
- [Brand Name] controls a wide variety of existing broadleaf weeds while they feed your desirable grass
- Kills over [XXX] weeds [down to the root]
- Covers [up to] [XXXX] sq.ft.
- Treats northern and southern grasses[\*] [\*not for 'Floratam' St. Augustinegrass]
- Kills [listed] weeds now for fewer weeds in spring
- Tough weed killer
- Kills even the tough[est] broadleaf weeds [like clover, dandelion, oxalis]
- Kill the toughest weeds like clover, dandelion, henbit, chickweed, spurge and many other listed broadleaf weeds
- Use for a beautiful weed-free lawn
- Kills weeds while feeding your lawn

Fertilizer Benefit Claims:

- Greens for [up to] 3 months (to be used only on formulations with controlled release technology)
- Quick[er] greening [and extended feeding] (to be used only on formulations with controlled release technology)
- Feeds for [up to] 3 months [guaranteed] (to be used only on formulations with controlled release technology)
- Helps strengthen lawn against heat & drought
- Rich in nitrogen for lush, green grass
- Enriched with potassium for overall plant vigor
- Phosphorus free (to be used only on formulations containing 0% phosphorus)
- [Contains] Nitrogen for lush, green plant growth
- [Contains] Potassium for overall plant vigor
- The only brand with MYCO ADVANTAGE™ Better results than fertilizer alone
- Enhanced efficiency fertilizer (to be used only on formulations with enhanced efficiency nitrogen)
- You only need [XX] pounds to cover [XXXX] square feet
- Feed more lawn with less fertilizer (to be used only on formulations with enhanced efficiency nitrogen)
- Promotes thick, full grass
- Delivers nutrients more efficiently than ordinary fertilizers (to be used only on formulations with enhanced efficiency nitrogen)
- Allows grass to use more of the fertilizer you buy & apply (to be used only on formulations with enhanced efficiency nitrogen)

#### Fungi Benefit Claims:

- Boosts nutrient uptake for a lush, green lawn
- Increases drought tolerance & reduces watering needs by extending root absorption area
- Increased root absorption area for more efficient use of water in [the] soil
- Delivers nutrients more efficiently than ordinary fertilizers
- Allows grass to use more of the fertilizer you buy & apply
- Contains mycorrhizal fungi for enhanced nutrient uptake
- Increased root absorption area to protect against heat & drought
- Increase absorption area of roots
- Enhance nutrient uptake
- Access and store more water
- Increased absorption of fertilizers

#### 3. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading "State Restrictions"]

- California, Oregon, Washington: "Information regarding the contents and levels of metals in this product is available on the internet at <u>http://www.aapfco.org/metals.htm</u>" [OR by calling [1-877-726-6275]
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
  - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
  - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
  - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida: "Apply only to actively growing turf."
- Florida: "Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- · Idaho: "Apply only as directed"
- Maryland: "This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area."
- New Jersey: "In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the "Soil Erosion and Sediment Control Act," P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots."
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: "In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency."
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]

- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supllies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program."
- When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials. Information regarding the contents and levels of metals in this product is available on the Internet at http://www.aapfco.org/metals.htm
- Certain states have limits on fertilizer application rates and timing. Consult your State Cooperative Extension for current information on annual application limitations for lawn fertilizers in your region.
- Information regarding the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.
- Each registrant shall substitute a unique alphanumeric identifier for "xx". This statement may be used only if the licensee establishes and maintains the Internet site: there is a clearly visible, direct hyperlink to a government website; and the Internet site contains no advertising or company specific information

### **DOCUMENT CONTROL INFORMATION**

- 1. Unique Label Identifier: 002217-00997.20160425.amend-proposed-highlighted
- 2. Reason for Issue: EPA Comments #1