

#### U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

Date of Issuance:

2217-1015

EPA Reg. Number:

03/30/2016

						D	$\Gamma$		T	71		
•			н.	( )	١н	$\mathbf{P}$	н 🔪			- 1		E:
. 1	,			` '				, ,	1		ш	<b>,</b> , ,

X Registration Reregistration (under FIFRA, as amended) Term of Issuance: Conditional

Name of Pesticide Product:

EH-1567 Herbicide

Name and Address of Registrant (include ZIP Code):

Mr. Eric D. Smith PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/registration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

**Signature of Approving Official:** 

Date:

03/30/2016

Kathryn Montague, Product Manager 23

astryn V. W Tontaguo

Herbicide Branch

Registration Division (7505P) Office of Pesticide Programs

- 2. You are required to comply with the data requirements described in the DCI identified below:
  - a. 2,4-D, Dimethylamine salt GDCI-030019-1358

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCIs listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1

- 3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 2217-1015."
- 4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 11/06/2015
- Alternate CSF 1 dated 11/06/2015
- Alternate CSF 2 dated 11/06/2015
- Alternate CSF 3 dated 11/06/2015
- Alternate CSF 4 dated 11/06/2015

If you have any questions, please contact Karen Samek by phone at (703) 347-8825, or via email at samek.karen@epa.gov.

Enclosure

# **EH-1567 HERBICIDE**

# 03/30/2016

ACCEPTED

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2217-1015

# EPA Reg. No. 2217-XXXX

Two sublabels represent the entire or master label of EH-1567 HERBICIDE. Please refer to the appropriate section of the labeling as shown as follows:

Section			
Label language common to all sublabels			
Sublabel 1. Concentrate			
Sublabel 2. Ready Spray			
Appendix			
Document Control Information			

# **Label Language Common To All Sublabels**

# EH-1567 HERBICIDE

# ACCEPTED

03/30/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 2217-1015

# EPA Reg. No. 2217-XXXX

ACTIVE INGREDIENT:	2217 1010
2,4-D, dimethylamine salt	7.24%
Quinclorac	4.00%
Dicamba, dimethylamine salt	0.48%
OTHER INGREDIENTS:	<u>88.28%</u>
TOTAL	100.00%

#### THIS PRODUCT CONTAINS:

0.517 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.01% 0.344 lb 3,7-dichloro-8-quinolinecarboxylic acid per gallon or 4.00% 0.035 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.40% Isomer Specific By AOAC Methods.

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

#### **KEEP OUT OF REACH OF CHILDREN**

# CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW **DIRECTIONS CAREFULLY.** 

# PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Harmful if swallowed. When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

## **User Safety Recommendations**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	First Aid					
If in eyes:	<ul> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>					
If swallowed:	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>					
Have the produc	et container or label with you when calling a poison control center or doctor or going for					

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

#### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

# **DIRECTIONS FOR USE**

[Note to Reviewer: The following section is for nonrefillable plastic containers and residential use only.]

#### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

# LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

<b>Alternate</b>	Warranty:
------------------	-----------

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for lack of performance or safety if not used according to the label directions.

# SUBLABEL 1 Concentrate

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### 1. Use Restrictions:

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
   Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85° F.
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

#### 2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Spot Treatments: Mix 2.5 fl.oz. (5 Tbs) in 1 gallon of water to treat 400 sq.ft. Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. per 1,000 sq.ft.
- [12 volt &] Tow-Behind Sprayers: Mix 6.4 fl.oz. in 0.5 to 1 gallon of water to treat 1,000 sq.ft.

Over-application or rates above those specified on this label can cause turf injury.

#### 3. When To Use

Apply this product when weeds are small and actively growing.

Weed species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 21 days later for dense populations.
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 21 days later if needed.

Weed species	Timing	Comments
	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with
listed broadleaf weeds		adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds		Apply when broadleaf weeds are young and actively growing with adequate soil moisture

The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.1 lb 2,4-D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.25 lb 2,4-D acid equivalent per acre]

## 4. How To Apply

#### 4.1 Pump Style Sprayers [sprayer icon]

Ideal for spraying individual weeds

- 1. Add the appropriate amount of water and concentrate to the sprayer tank.
- 2. Close sprayer, shake well, and pump handle to pressurize.
- 3. Adjust nozzle to deliver a coarse spray pattern.
- 4. To broadcast over large areas: apply evenly to treated area
- 5. To treat individual weeds: spray to wet weed leaf surfaces.
- 6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

#### 4.2 [12 Volt &] Tow-Behind Boom Sprayers [sprayer icon]

Ideal for treating large country and suburban lawns

- 1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1,000 sq.ft.. For example, at 4 mph a sprayer might deliver 1/2 gallon per 1,000 sq.ft. at 40 psi.
- 2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
- 3. Mix the appropriate amount of product and water in the sprayer using the following tables as a guide.

Table 1: 12 Volt & Tow-Behind Sprayers: Application Guide (Examples)					
Tank Size	Spray volume	Coverage Area	Concentrate to add to tank		
15 gol	0.5 gal/1000 sq.ft.	30,000 sq.ft.	192 fl.oz. (1.5 gal)		
15 gal	1 gal/1000 sq.ft.	15,000 sq.ft.	96 fl.oz. (0.75 gal)		
25 gol	0.5 gal/1000 sq.ft.	50,000 sq.ft.	320 fl.oz. (2.5 gal)		
25 gal	1 gal/1000 sq.ft.	25,000 sq.ft.	160 fl.oz. (1.25 gal)		

#### For spot treatments with [12 Volt &] Pull Behind Sprayers

See rates under "How Much To Apply," Spot Treatments.

#### 4.3 Dial Style Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

- 1. Fill sprayer with sufficient concentrate to cover the total area to be treated. DO NOT ADD WATER.
- 2. Close sprayer and set dial to 2.5 oz.
- 3. Connect sprayer to hose

- 4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
- 5. Pour any unused product back into this container.

#### 4.4 Concentrate Plus Water Hose-End Sprayers [sprayer icon]

Ideal for treating medium-sized lawns

- 1. Fill sprayer with sufficient concentrate to cover the total area to be treated. Pour product into sprayer jar to the "Fluid Oz." level desired (See sprayer jar for markings)
- 2. Add the appropriate amount of water to the "Gallons" level indicated on the side of the sprayer jar.
- 3. Close sprayer, connect to garden hose, shake well.
- 4. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.

#### 5. Use Precautions:

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- For new lawns, apply only when grass is well-established (approximately 6 weeks after seedling emergence) or after the third mowing.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

#### 6. Grass and Broadleaf Weeds Killed

GRASS WEEDS					
Crabgrass, large and	Foxtail, green, yellow and giant	Signalgrass, broadleaf			
smooth					

BROADLEAF WEEDS				
Annual yellow sweetclover	Cockle	Musk thistle	Speedwell	
Aster	Cocklebur	Mustard	Spiny amaranth	
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny cocklebur	
Bedstraw	Creeping Jenny (Creeping	Narrowleaf vetch	Spiny sowthistle	
Beggarticks	Charlie)	Nettle	Spotted catsear	
Betony, Florida	Cudweed	Orange hawkweed	Spotted spurge	
Bindweed, field	Curly dock	Oriental cocklebur	Spurweed	
Bird vetch	Daisy, English	Oxalis	Stinging nettle	
Bitter wintercress	Daisy fleabane	Parsley-piert	Strawberry, India mock	
Bittercress, hairy	Daisy, oxeye	Parsnip	Tall nettle	
Bitterweed	Dandelion	Pearlwort	Tall vervain	
Black-eyed Susan	Dichondra	Pennycress	Tansy ragwort	
Black medic	Dogbane	Pennywort	Tansy mustard	
Black mustard	Dogfennel	Peppergrass	Tanweed	
Blackseed plantain	Dollarweed	Pepperweed	Thistle	
Blessed thistle	False dandelion	Pigweed	Trailing crownvetch	
Bloodflower milkweed	Falseflax	Pineywoods bedstraw	Tumble mustard	
Blue lettuce	False sunflower	Plains coreopsis	Tumble pigweed	
Blue vervain	Fiddleneck	(tickseed)	Velvetleaf	
Bracted plantain	Florida pusley	Plantain	Venice mallow	

BROADLEAF WEEDS			
Brassbuttons	Frenchweed	Poison ivy	Virginia buttonweed
Bristly oxtongue	Galinsoga	Poison oak	Virginia creeper
Broadleaf dock	Goathead	Pokeweed	Virginia pepperweed
Broadleaf plantain	Goldenrod	Poorjoe	Wavyleaf bullthistle
Broomweed	Ground ivy	Prairie sunflower	Western clematis
Buckhorn	Gumweed	Prickly lettuce	Western salsify
Buckhorn plantain	Hairy fleabane	Prickly sida	White mustard
Bulbous buttercup	Hawkweed	Prostrate knotweed	Wild aster
Bull thistle	Healall	Prostrate pigweed	Wild buckwheat
Bullnettle	Heartleaf drymary	Prostrate spurge	Wild carrot
Burclover	Heathaster	Prostrate vervain	Wild four-o'clock
Burdock	Hedge bindweed	Puncturevine	Wild garlic
Burning nettle	Hedge mustard	Purslane, common	Wild geranium
Burweed	Hemp	Ragweed	Wild lettuce
Buttercup	Henbit	Red sorrel	Wild marigold
Buttonweed	Hoary cress	Redroot pigweed	Wild mustard
Canada thistle	Hoary plantain	Redstem filaree	Wild onion
Carolina geranium	Hoary vervain	Rough cinquefoil	Wild parsnip
Carpetweed	Horsenettle	Rough fleabane	Wild radish
Catchweed bedstraw	Jimsonweed	Russian pigweed	Wild rape
Catnip	Knawel	Russian thistle	Wild strawberry
Catsear	Knotweed	Scarlet pimpernel	Wild sweet potato
Chickweed, common	Kochia	Scotch thistle	Wild vetch
Chickweed, mouseear	Lambsquarters	Sheep sorrel	Wild violet
Chicory	Lespedeza	Shepherd's purse	Woodsorrel
Cinquefoil	Mallow	Slender plantain	Woolly croton
Clover, crimson	Matchweed	Smallflower galinsoga	Woolly morningglory
Clover, hop	Mexicanweed	Smooth dock	Woolly plantain
Clover, red	Milk vetch	Smooth pigweed	Wormseed
Clover, strawberry	Morningglory	Sorrel	Yarrow
Clover, sweet	Mouseear hawkweed	Sowthistle	Yellow rocket
Clover, white	Mugwort	Spanishneedles	Yellowflower pepperweed

# 7. Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

# SUBLABEL 2 Ready Spray

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### 1. Use Restrictions:

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
   Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Use on established lawns. Do not use on Bahiagrass, Bentgrass (colonial and creeping), Carpetgrass, Centipedegrass, Dichondra, Gardens (including vegetables, fruit trees, vines and berries), Ornamental plants (flowers, trees, groundcovers, landscape beds and shrubs), Seashore Paspalum, St. Augustinegrass, and turfgrass species that are not listed on this label.)
- Do not use clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.
- Do not apply this product to food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants including flowers, trees, shrubs, hedges, woody ornamentals, groundcovers established in landscape plantings and other nontarget plants.
- Do not spray exposed roots of ornamentals and trees.

#### 2. How Much To Apply

For Bermudagrass (common and hybrid), Buffalograss, Kentucky Bluegrass, mixtures of cool season grasses containing fine fescues (red, hard and chewings), Ryegrass (perennial and annual), Tall Fescue, and Zoysiagrass

Entire Lawn Treatments: This container covers XXX sq.ft. Measure the size of your yard. If the area to be treated is larger than XXX sq.ft., you may need to buy more containers.

Over-application or rates above those specified on this label can cause turf injury.

## 3. When To Use

Apply this product when weeds are small and actively growing.

Weed species	Timing	Comments
Crabgrass (large and smooth):	Apply in spring from crabgrass emergence until the crabgrass is 3 to 4 inches in height	Re-treat at the same rate at 21 days later for dense populations or for continued crabgrass emergence.
Foxtail (green, giant and yellow):	Apply in spring and early summer during the early growth stages or when foxtail is 1 to 4 inches in height.	Re-treat at the same rate at 21 days later for dense populations.
Broadleaf signalgrass	Apply in spring and early summer during the early growth stages or during the 3 to 7 leaf stage.	Re-treat at the same rate at 21 days later if needed.
Dandelion, plantain, buttercup, speedwell, knotweed and other listed broadleaf weeds	Apply in spring when broadleaf weeds are small and actively growing.	Apply when broadleaf weeds are young and actively growing with adequate soil moisture.
Henbit, chickweed, white clover, ground ivy, lawn burweed and other listed broadleaf weeds	Apply in fall when broadleaf weeds are young and actively growing	Apply when broadleaf weeds are young and actively growing with adequate soil moisture

Weed species Timing		Comments
---------------------	--	----------

The maximum application rate is 6.4 fl.oz. of product per 1,000 sq.ft. per application [1.1 lb 2,4-D acid equivalent per acre per application]. The maximum number of applications is limited to 2 per year with a minimum of 21 days between applications. The maximum seasonal rate is 12.8 fl.oz. of product per 1,000 sq.ft. [2.25 lb 2,4-D acid equivalent per acre]

#### 4. How To Apply

- 1. Connect your garden hose to the hose-end sprayer. With your thumb, snap off the safety locking tab that prevents accidental discharge of spray solution. Leave control valve in the OFF position.
- 2. To BEGIN spraying: Turn on water at faucet. Extend hose to the furthest point from the faucet. Point nozzle toward the lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Walk backwards as you apply in order to avoid contact with the treated area. For best results, spray evenly over the measured area.
- 3. To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

#### 5. Use Precautions:

- Apply when weeds are young and actively growing.
- Delay mowing 2 days before and until 2 days after the application of this product.
- If the lawn is dry, water 24 hours before using this product.
- Do not water lawn within 24 hours of treatment.
- Certain hard-to-kill weeds may need to be spot treated in 21 days.
- For new lawns, apply only when grass has reached a height of at least 2 inches.
- Bare spots may be seeded 3 to 4 weeks after application.
- Spray when air temperatures are cooler than 90°F.
- Avoid applications during the spring transition of the warm season grasses and during periods of extremely high temperatures.
- Application to bermudagrass may cause temporary yellowing or discoloration but full recovery can be expected. Do not apply to bermudagrass when daytime temperatures exceed 85°F.

#### 6. Grass and Broadleaf Weeds Killed

GRASS WEEDS			
Crabgrass, large and smooth	Foxtail, green, yellow and giant	Signalgrass, broadleaf	

BROADLEAF WEEDS			
Annual yellow sweetclover	Cockle	Musk thistle	Speedwell
Aster	Cocklebur	Mustard	Spiny amaranth
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spiny cocklebur
Bedstraw	Creeping Jenny (Creeping	Narrowleaf vetch	Spiny sowthistle
Beggarticks	Charlie)	Nettle	Spotted catsear
Betony, Florida	Cudweed	Orange hawkweed	Spotted spurge
Bindweed, field	Curly dock	Oriental cocklebur	Spurweed
Bird vetch	Daisy, English	Oxalis	Stinging nettle
Bitter wintercress	Daisy fleabane	Parsley-piert	Strawberry, India mock
Bittercress, hairy	Daisy, oxeye	Parsnip	Tall nettle
Bitterweed	Dandelion	Pearlwort	Tall vervain
Black-eyed Susan	Dichondra	Pennycress	Tansy ragwort
Black medic	Dogbane	Pennywort	Tansy mustard
Black mustard	Dogfennel	Peppergrass	Tanweed
Blackseed plantain	Dollarweed	Pepperweed	Thistle
Blessed thistle	False dandelion	Pigweed	Trailing crownvetch

DDG ADI EAS WESS			
BROADLEAF WEEDS	T	_	1
Bloodflower milkweed	Falseflax	Pineywoods bedstraw	Tumble mustard
Blue lettuce	False sunflower	Plains coreopsis	Tumble pigweed
Blue vervain	Fiddleneck	(tickseed)	Velvetleaf
Bracted plantain	Florida pusley	Plantain	Venice mallow
Brassbuttons	Frenchweed	Poison ivy	Virginia buttonweed
Bristly oxtongue	Galinsoga	Poison oak	Virginia creeper
Broadleaf dock	Goathead	Pokeweed	Virginia pepperweed
Broadleaf plantain	Goldenrod	Poorjoe	Wavyleaf bullthistle
Broomweed	Ground ivy	Prairie sunflower	Western clematis
Buckhorn	Gumweed	Prickly lettuce	Western salsify
Buckhorn plantain	Hairy fleabane	Prickly sida	White mustard
Bulbous buttercup	Hawkweed	Prostrate knotweed	Wild aster
Bull thistle	Healall	Prostrate pigweed	Wild buckwheat
Bullnettle	Heartleaf drymary	Prostrate spurge	Wild carrot
Burclover	Heathaster	Prostrate vervain	Wild four-o'clock
Burdock	Hedge bindweed	Puncturevine	Wild garlic
Burning nettle	Hedge mustard	Purslane, common	Wild geranium
Burweed	Hemp	Ragweed	Wild lettuce
Buttercup	Henbit	Red sorrel	Wild marigold
Buttonweed	Hoary cress	Redroot pigweed	Wild mustard
Canada thistle	Hoary plantain	Redstem filaree	Wild onion
Carolina geranium	Hoary vervain	Rough cinquefoil	Wild parsnip
Carpetweed	Horsenettle	Rough fleabane	Wild radish
Catchweed bedstraw	Jimsonweed	Russian pigweed	Wild rape
Catnip	Knawel	Russian thistle	Wild strawberry
Catsear	Knotweed	Scarlet pimpernel	Wild sweet potato
Chickweed, common	Kochia	Scotch thistle	Wild vetch
Chickweed, mouseear	Lambsquarters	Sheep sorrel	Wild violet
Chicory	Lespedeza	Shepherd's purse	Woodsorrel
Cinquefoil	Mallow	Slender plantain	Woolly croton
Clover, crimson	Matchweed	Smallflower galinsoga	Woolly morningglory
Clover, hop	Mexicanweed	Smooth dock	Woolly plantain
Clover, red	Milk vetch	Smooth pigweed	Wormseed
Clover, strawberry	Morningglory	Sorrel	Yarrow
Clover, sweet	Mouseear hawkweed	Sowthistle	Yellow rocket
Clover, white	Mugwort	Spanishneedles	Yellowflower pepperweed

## 7. Spray Drift Management

It is the responsibility of the applicator to avoid spray drift at the application site. Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable broadleaf plants, as small amounts of this product can damage sensitive plants near the treated area. If desirable plants are accidentally sprayed, immediately rinsing leaves with water may reduce or eliminate plant damage.

# **APPENDIX**

# 1. Statements which may appear on different label components depending on packaging configuration.

nfiguration.			
•	See next panel for additional Precautionary Statements and First Aid		
•	Net Contents:		

-	riot Contonto.	
•	EPA Est. No	

#### 2. Advertising claims that may be presented on the retail containers of supplemental registrants.

#### **Efficacy Claims**

- Controls crabgrass, dandelion, chickweed, knotweed, henbit, plantain, clover, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- Kills Crabgrass and over XXX Listed Weeds in Country & Suburban Lawns\* \*See inside booklet for a complete list of weeds controlled
- Kills [controls] [Crabgrass,] [Dandelion,] [Clover,] and other common [lawn weeds] [weeds found in your lawn] (See inside booklet for a complete list of weeds controlled)
- Controls crabgrass and broadleaf weeds in lawns (See inside booklet for a complete list of weeds controlled)
- Kill crabgrass & [any listed weed(s)] too!
- Kills Crabgrass [plus [over XXX] Broadleaf Weeds] (See inside booklet for a complete list of weeds controlled)
- Kills both Crabgrass & Broadleaf weeds (See inside booklet for a complete list of weeds controlled)
- Kills lawn weeds plus crabgrass. (See inside booklet for a complete list of weeds controlled)
- Kills [Controls] 200+ broadleaf weeds. (See inside booklet for a complete list of weeds controlled)
- Controls (Number) weeds! (See inside booklet for a complete list of weeds controlled)

#### CONTROLS

- **SENSITIVE WEEDS:** Crabgrass, Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed and Wild Lettuce.
- HARD-TO-KILL WEEDS: Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and other broadleaf weeds. (See inside booklet for a complete list of weeds controlled)
- Kills [Lawn] Weeds [plus] [including] Crabgrass in One [Easy] Step (See inside booklet for a complete list of weeds controlled)
- Kills Over 200 Lawn Weeds Plus Crabgrass in One Easy Step (See inside booklet for a complete list
  of weeds controlled)
- Kills [Controls] Listed Weeds (See inside booklet for a complete list of weeds controlled)
- Kills Hard-to-Control Weeds (See inside booklet for a complete list of weeds controlled)
- Kills [Controls] Both Broadleaf and Grass Weeds [in lawns] (See inside booklet for a complete list of weeds controlled)
- [Also] Kills [Dandelion], [Crabgrass], [Clover], plus XXX Other [Listed] Weeds [As Listed] (See inside booklet for a complete list of weeds controlled)
- Three-in-One Product that Kills Crabgrass & Broadleaf Weeds (See inside booklet for a complete list
  of weeds controlled)
- Effectively Kills Both Broadleaf and Certain Weed Grasses (See inside booklet for a complete list of weeds controlled)
- This product kills Crabgrass PLUS broadleaf weeds. The dual-action formula goes beyond crabgrass and foxtail to kill over XXX other broadleaf weeds listed in country and suburban lawns. (See inside booklet for a complete list of weeds controlled)
- Lawn Weed & Crabgrass Killer (See inside booklet for a complete list of weeds controlled)
- Kills Dandelions, Clover, and Other Broadleaf Weeds! Plus Kills Crabgrass and Other grass Weeds! (See inside booklet for a complete list of weeds controlled)

- Kills the Toughest Weeds in these Lawns (any lawn listed on label) (See inside booklet for a complete list of weeds controlled)
- Kills Over 200 Lawn Weeds plus [including] Crabgrass [in one easy step]. (See inside booklet for a complete list of weeds controlled)
- Kills Major [Lawn] Broadleaf Weeds (See inside booklet for a complete list of weeds controlled)
- Kills the weeds not the lawn when used as directed
- Kill [control] the most common weeds in your lawn [grass] (See inside booklet for a complete list of weeds controlled)
- Kills [controls] [hard-to-control] [tough] weeds (See inside booklet for a complete list of weeds controlled)
- Tough weed killer
- Kills [controls] all types of [lawn] weeds [in your grass] (See inside booklet for a complete list of weeds controlled)
- Controls [Kills] even the tough weeds...roots and all!
- Kills the root.
- Kills Even Tough Weeds Down to the root
- Kills Crabgrass and Broadleaf weeds...-Down to the root
- Gets to the root [of your weed problem]
- Kills [lawn] weeds [down] to the root[s]
- Kills to the root
- Kills [controls] XX types of weeds [- broadleaf or grass weeds] (See inside booklet for a complete list
  of weeds controlled)
- Kill[s] [controls] [over] [up to] [XXX+] [lawn] [types of] weeds (See inside booklet for a complete list of weeds controlled)
- Won't harm lawn grass when used according to label directions.
- Won't harm [lawn] grass when used as directed.
- Won't harm lawns\* \*When used as directed
- Kills Weeds, Not Lawns(TM)\* \*when used as directed
- [Even] kills [controls] poison ivy and poison oak
- [Only] targets the weeds, leaving your lawn [grass] unharmed when used as directed
- Targets weeds invading your lawn [grass]
- [Visible] [See] results in X hours
- Kills by contact
- Starts working on contact
- Results [works] with 1 [one] application
- One [1] [easy] solution for crabgrass, dandelions and clover

#### **Formula Claims**

- Keep from freezing.
- Water based.
- Mixes with Water Concentrate
- Proprietary Formula that Kills Both Crabgrass and Broadleaf Weeds
- 2-in-1
- All-in-One [complete] [total] [2-in-1] weed control [for lawns]
- All in One Weed Solution
- Colorless formula
- Non-staining
- Stain-free [formula]

#### Characteristics

- XX covers up to XXXX square feet.
- Makes more than XX Gallons!
- One quart covers up to 5,000 square feet.

- One [1] [bottle size] covers up to XXXX square feet
- 1 quart treats [up to] 5,000 sq.ft.
- Covers [treats up to] [XXXX] sq.ft.
- [XX] [Pint, Quart, etc] Treats Up To [YYYY] Sq.Ft.
- Bonus Size
- 25% More Free!
- Easily covers large areas
- Economical use to cover large areas
- [Great] [Ideal] for broadcast treatment [application]
- Great value [- use to cover small or large areas]
- Designed for use on Southern lawns [warm season grasses/list approved grass types here]
- Treats many Northern & Southern grass types
- For use on [many Northern & Southern] grasses [lawns] including XXX
- XX% more than X gallon size
- The value of XX [size RTU] in one concentrate

#### Weather & Speed

- Starts working [overnight] [immediately]
- Rainfast in 1 Hour
- [Rainproof] [rainfast] in [1] [X] Hour[s]
- [See] results in hours
- · Overnight results
- Fast-acting [formula]

#### **Applicator Claims**

- [Easy] [Ergonomic] Comfort Grip [Design] [Handle]
- Ergonomic [design]
- Easy grip [control]
- Easy [to use]
- · Simple to use
- Convenient [Simple] [Design]
- Simple [and easy] [to use]
- [Simply] attach to [garden] hose and spray [go]
- · Hand doesn't get wet
- · Hand stays dry
- · Hands stay dry
- No messy clean-up
- Includes water [only] setting [option]
- Instant/Easy [One-handed] on/off applicator/switch/action
- One-handed operation [use] [spraying]
- Ready-Spray®/Ready-to-Spray/Applicator Name
- Ready-Spray
- [Requires] no measuring, [no mixing]
- · No mixing required
- Just connect to garden hose & spray
- [Quick and] Easy [and simple] to use
- [The] Easy way to spray
- Easy [Simple] to attach [to [garden] hose]
- One step prep
- [Applicator name]
- Save[s] time [and energy]
- Mess free cap [top] [lid] [cup]
- Measure with ease

- Easy-Measure [Cap] [Top] [Lid] [Cup]
- Drain-Back [Cap] [Top] [Lid] [Cup]
- · Quick, convenient, and mess free
- [reusable] [drain-back/measuring] cap

#### Other

- Local Solutions for Local Problems
- For more Lawn and Garden Products Visit GordonsUSA.com
- Open Resealable Label for Directions & Precautions (Press to Reseal)
- For [Use on] Lawns
- For home lawn care
- For Home Use [Only]
- Weed control for your lawn
- Weed control made easy
- For outdoor use
- Easy to mix
- Proven performance
- Systemic [Herbicide] [Activity]
- Be in control of weeds in your yard
- From the lawn care experts at Scotts
- Help[s] [keep[s]] the lawn look[ing] beautiful
- Lawn friendly [weed control] when used as directed
- Guaranteed not to harm the lawn when used as directed
- How it works: The [insert brand name] formula kills BOTH broadleaf weeds and Crabgrass in one
  easy step [without harming your lawn\* \*when used as directed guaranteed]. (See inside booklet for
  a complete list of weeds controlled)
- Read Entire label before use
- · Recyclable bottle
- [Recycle symbol]
- Takes the guesswork out of weed control [in the lawn]
- XX% less plastic . . . As compared to [alternate product]
- Optional Instructions for Side Clip Tape

PULL / PEEL HERE [TO REMOVE TAPE]

Open Resealable Label For Directions and/& Precautions

Press to Reseal

- Questions & Comments: [icon phone] [icon computer]
- Call 1-800- 884-3179 or Visit our Website at GordonsUSA.com
- [Superscript is first letter of lot number]

[The following Product Facts format is optional and the information may appear inside or outside of a box.]

#### PRODUCT FACTS:

What It Does: Kills major broadleaf and troublesome grass weeds including dandelion, chickweed and clover as well as crabgrass and foxtail. (See inside for complete list)

Where to Use: Lawns: Kills the weeds not the lawn when used as directed. (icon weeds)

When to Use: Apply when daytime temperatures are below 90°F. (do not apply above 85°F for bermudagrass).

**Questions & Comments:** [icon phone] [icon computer] Call 1-800-884-3179 or Visit our Website at GordonsUSA.com

#### 3. Trademark acknowledgements

- Ortho® Dial'N Spray® Hose-End Sprayer is a registered trademark of The ORTHO Group.
- Hudson® Hose End Liquid Concentrate Sprayer is a registered trademark of the H.D. Hudson Manufacturing Company.
- Gilmour® Dial-A -Mix® Lawn and Garden Sprayer is a registered trademark of the Gilmour Manufacturing Company.
- Trimec® is registered trademark of PBI/Gordon Corporation

# **DOCUMENT CONTROL INFORMATION**

1. Unique Label Identifier: 002217-0XXXX.20160229.new-proposed-EH1567-clean

2. Reason for Issue: EPA Comments #1