



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 20, 2022

Eric D. Smith
Director, Regulatory Affairs
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286

Subject: Label Amendment – Addition of NY Restriction and Revised Weeds List, Use
Directions, and Storage and Disposal
Product Name: EH-1566 ORP GRANULE
EPA Registration Number: 2217-1024
Application Date: 8/24/2020
Decision Number: 566022

Dear Eric D. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 2217-1024
Decision No. 566022

with FIFRA section 6. If you have any questions, please contact Sarah Meadows by phone at 202-566-2828, or via email at meadows.sarah@epa.gov.

Sincerely,



Emily Schmid, Product Manager 25
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs


Enclosure

PYRIMISULFAN	GROUP	2	HERBICIDE
--------------	-------	---	-----------

EH-1566 ORP GRANULE

EPA Reg. No. 2217-1024

Alternate Brand Names: Vexis Herbicide Granular



Not for sale, distribution or use in Nassau or Suffolk Counties in New York State.

ACTIVE INGREDIENT:

Pyrimisulfan	0.025%
OTHER INGREDIENTS:	<u>99.975%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS:

0.5 lb Pyrimisulfan per ton or 0.025% (0.00025 lb pyrimisulfan per pound).

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See [back panel] [side panel] for complete Precautionary Statements, including First Aid and Directions for Use, including Storage and Disposal.

Net Contents: _____
EPA Est. No. _____

Company Name and Address:
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

Personal Protective Equipment (PPE)

All mixers, loaders, and applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Removing clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Advisory: This product has a potential for reaching surface water via runoff after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of pyrimisulfan from runoff water. Runoff of this product will be reduced by avoiding applications when heavy rainfall or irrigation is expected to occur within 48 hours.

Physical and Chemical Hazards

Do not mix or allow to come into contact with an oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

[Note to reviewer: The Ag Use and Non-Ag Use Requirements boxes are only required when container labels include use on sod farms.]

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls,
- chemical-resistant gloves made of any water-proof material,
- shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until dust has settled.

1. Product Description

EH-1566 ORP GRANULE is a selective herbicide for post-emergent control of listed broadleaf weeds, sedges, kyllinga, and rush species commonly found in turfgrasses. It can be used on established warm-season turfgrasses (including improved varieties of St. Augustinegrass such as Bitterblue and Floratam) and cool-season turfgrasses (excluding bentgrass mowed under 1/2 inch). Fall applications provide control of winter weeds such as henbit, false dandelion and chickweed.

2. Resistance Management

For resistance management, this product contains Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 2 herbicides. The resistant biotypes

may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, call 877-800-5556.

3. Weeds Controlled

EH-1566 ORP GRANULE will control the following weeds:

Sedges, Kyllinga, and Rushes (including ALS resistant annual sedges)

Yellow Nutsedge	Green Kyllinga	Annual Kyllinga *
Purple Nutsedge	False Green Kyllinga	Path Rush
Annual Nutsedge *	Cockscomb Kyllinga	

Broadleaf Weeds

Buckhorn Plantain	Ground Ivy (Creeping Charlie)	Rough Fleabane *
Common Chickweed	Hairy Bittercress	Virginia Buttonweed *
Dollarweed (Pennywort)	Henbit	Wild Garlic
False Dandelion	Lawn Burweed	Wild Onion
Florida Betony *	Purple Deadnettle *	

[* Not for use in California]

4. Where To Use

This product provides weed control on the following sites.

- **Ornamental Turfgrass sites:**
 - **Residential/domestic sites** including areas associated with household or home life including apartment complexes and condominiums.

- **Ornamental turf sites** including turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, playgrounds, fairgrounds, and athletic fields.
- **Institutional sites** including properties or facilities providing a service to public or private organizations including, hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses, and office buildings.
- **[Non-cropland sites:** including farmyards, fencerows or fence lines, highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians; Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas; Municipal, state, and federal lands: Airports and military installations; railroad rights-of-ways, railroad yards, railroad crossings and railroad bridge abutments; Utility rights-of-way: telephone, pipeline, electrical powerlines, and communication transmission lines.]
- **[Agricultural site:** Commercial sod production]

Turfgrass Species Which May be Treated:

EH-1566 ORP GRANULE may be used on the following established turfgrass species:

- **Established Cool-Season Turfgrass:** Kentucky bluegrass, fine fescues, tall fescue, perennial ryegrass, bentgrass.
- **Established Warm-Season Turfgrass:** Bermudagrass (common or hybrid), bahiagrass, buffalograss, centipedegrass, kikuyugrass, seashore paspalum, St. Augustinegrass (including improved varieties such as Bitterblue and Floratam), zoysiagrass.

Use Restrictions:

- Do not apply more than 8.26 lb of product per 1000 sq.ft. per year.
- Do not retreat within 30 days of initial application.
- Do not use on carpetgrass, dichondra, legumes, or turfgrass where desirable clovers are present.
- Do not apply to bentgrass mowed under 1/2 inch.
- Do not collect grass clippings from treated turfgrass for use as mulch around plants or in compost for 3 weeks after treatment. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- Do not apply directly to flowers, vegetables, groundcovers, gardens or landscape plantings including ornamental shrubs or trees.

Use Precautions:

- Delay application of this product to grass seedlings until after the second mowing.
- Delay application of this product until 3 weeks after sodding, sprigging, or plugging operations.
- Delay sodding, sprigging, seeding or plugging operations until 3 weeks after application of this product.
- Delay application of this product on Centipedegrass and improved varieties of St Augustinegrass until after the second mowing in the spring when the lawn is fully green and actively growing.

5. Application Directions

How Much to Apply:

Apply 2.5 to 5.5 lb of product per 1000 sq.ft. (109 to 240 lb of product per acre). Retreatment after 30 days is allowed but do not apply more than a total of 8.26 lb per 1000 sq.ft. per year (360 lb of product per acre per year).

When to Apply:

- Apply when weeds are actively growing. Retreatment may be necessary when weed pressures are high, when soil moisture is low, or when weeds are not actively growing.
- To avoid injury to improved varieties of St. Augustinegrass or Centipedegrass, apply when night-time temperatures are above 50°F for 7 consecutive nights.

Application Tips:

- Moisture on leaf surface is not required. Apply when weed leaves are wet or dry.
- Temperatures below 50°F will reduce weed control activity.
- For best results, water-in granules within 48 hours after treatment.
- For best results, postpone mowing until granules are watered-in.
- For best winter annual control (henbit, false dandelion, and chickweed) apply in early fall.

[Begin Optional Language]

Optional paragraph for shaker can packaging:
 [This 1 lb canister will cover 180 to 400 sq.ft.]

[How to Use the EH-1566 ORP Granule Shaker Can:

Applying EH-1566 ORP Granule with the shaker can is simple:

1. Be sure to read product label instructions before you begin.
2. Open the flip-top on the lid; Notch 1 provides ideal coverage.
3. Shake out product in a steady, consistent motion over target weeds.
4. Water in product per label instructions.
5. Close the shaker can lid for storage.]

Optional Language for bag packaging:

Select the appropriate bag size and present only that information. For example, "This 15 lb bag covers 2700 to 6000 sq.ft."]

Bag Size	Application Rate	Treated Area for This Bag Size	Bag Size	Application Rate	Treated Area for This Bag Size
12.5 lb	2.5 lb/1000 sq.ft.	5000 sq.ft.	25 lb	2.5 lb/1000 sq.ft.	10,000 sq.ft.
	3.1 lb/1000 sq.ft.	4000 sq.ft.		3.1 lb/1000 sq.ft.	8100 sq.ft.
	3.7 lb/1000 sq.ft.	3400 sq.ft.		3.7 lb/1000 sq.ft.	6800 sq.ft.
	5.5 lb/1000 sq.ft.	2300 sq.ft.		5.5 lb/1000 sq.ft.	4550 sq.ft.
14 lb	2.5 lb/1000 sq.ft.	5600 sq.ft.	30 lb	2.5 lb/1000 sq.ft.	12,000 sq.ft.
	3.1 lb/1000 sq.ft.	4500 sq.ft.		3.1 lb/1000 sq.ft.	9700 sq.ft.
	3.7 lb/1000 sq.ft.	3800 sq.ft.		3.7 lb/1000 sq.ft.	8100 sq.ft.
	5.5 lb/1000 sq.ft.	2550 sq.ft.		5.5 lb/1000 sq.ft.	5450 sq.ft.
15 lb	2.5 lb/1000 sq.ft.	6000 sq.ft.	35 lb	2.5 lb/1000 sq.ft.	14,000 sq.ft.
	3.1 lb/1000 sq.ft.	4800 sq.ft.		3.1 lb/1000 sq.ft.	11,300 sq.ft.
	3.7 lb/1000 sq.ft.	4100 sq.ft.		3.7 lb/1000 sq.ft.	9500 sq.ft.
	5.5 lb/1000 sq.ft.	2700 sq.ft.		5.5 lb/1000 sq.ft.	6400 sq.ft.
16 lb	2.5 lb/1000 sq.ft.	6400 sq.ft.	40 lb	2.5 lb/1000 sq.ft.	16,000 sq.ft.
	3.1 lb/1000 sq.ft.	5200 sq.ft.		3.1 lb/1000 sq.ft.	12,900 sq.ft.
	3.7 lb/1000 sq.ft.	4300 sq.ft.		3.7 lb/1000 sq.ft.	10,800 sq.ft.
	5.5 lb/1000 sq.ft.	2900 sq.ft.		5.5 lb/1000 sq.ft.	7300 sq.ft.
20 lb	2.5 lb/1000 sq.ft.	8000 sq.ft.	50 lb	2.5 lb/1000 sq.ft.	20,000 sq.ft.
	3.1 lb/1000 sq.ft.	6500 sq.ft.		3.1 lb/1000 sq.ft.	16,100 sq.ft.
	3.7 lb/1000 sq.ft.	5400 sq.ft.		3.7 lb/1000 sq.ft.	13,500 sq.ft.
	5.5 lb/1000 sq.ft.	3600 sq.ft.		5.5 lb/1000 sq.ft.	9100 sq.ft.

Calibration Instructions and Spreader Settings:

Calibrate the spreader according to the manufacturer’s directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

Spreader	Delivers ___ lb/1000 sq.ft.		
	2.5 lb	3.1 lb	5.5 lb
LESCO® Rotary calibration gauge setting #	X	X	X
Ortho® Drop (5000)	X	X	X
Ortho® Rotary (6000)	X	X	X

Distribution and Spreader Patterns:

Always apply with a properly calibrated spreader designed for granular materials such as a rotary or drop type fertilizer spreader or a shaker applicator [such as EH-1566 ORP Granule Shaker Can][such as Gordon's Spred-Rite® G Granule Spreader]. For best results, apply this product evenly and uniformly avoiding streaking, skips, or overlaps.

1. For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage.
2. First, spread the borders of your lawn and header strips for turning around. Then apply this product to the center with straight line passes
3. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

How to Refill:

[For Ready-To-Use Containers]

To refill this product container, use the EH-1566 ORP GRANULE bag and carefully pour granules directly into this container.]

[End Optional Language]

[Nonrefillable Container with refill use directions]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container unless allowed by the directions for use. Then offer for recycling if available or dispose of empty container in a sanitary landfill or by incineration.

[For Nonrefillable Container without refill use directions]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration.”

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable,

return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner or use of application, etc. Such factors and conditions are beyond the control of the manufacturer, and **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** To the extent consistent with applicable law, buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THE PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this LIMITED WARRANTY AND DISCLAIMER in any manner.

APPENDIX

1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Apply in the fall to control [kill] listed winter weeds such as henbit, false dandelion and chickweed
- Kill [Control] listed winter weeds such as henbit, false dandelion and chickweed with a fall application
- Provides [Delivers] [Offers] post-emergence control of listed broadleaf weeds, sedges, kyllinga, and rushes
- Can be used on established warm-season turfgrasses and cool-season turfgrasses as listed
- Provides [Delivers] [Offers] dependable season-long [120-day] [4-month] control
- 3-week reseed/resod interval
- Reseed/resod in just three weeks
- Short reseed/resod interval: 3 weeks
- Reseed/resod in just 21 days
- Apply to wet or dry turf
- Can be applied to wet or dry turf

- Flexible: can be applied to wet or dry turf
- Flexible application to wet or dry turf
- Turf can be wet or dry at application
- Turf does not have to be wet before application
- Effective [Performs] applied to either wet or dry turf
- Can be used on Bermudagrass
- Labeled for use on Bermudagrass.
- Can be used on many turfgrass species as listed [list species]
- Provides [Delivers] [Offers] [excellent] control of sedges, kyllinga, and rushes
- Kills sedges, kyllinga, and rushes fast [quickly]
- Provides [Delivers] [Offers] fast [rapid] [quick] control of sedges, kyllinga, and rushes
- Dependable control of sedges, kyllinga, and rushes
- A wide range of labeled use sites for versatile application options
- Multiple applications are allowed, not to exceed 8.26 lb of product per 1000 sq.ft. per year.
- No maximum temperature restriction allows for application when sedges are most active
- Simple [Easy-to-read] label
- Label instructions are simple to read and easy to understand
- Shaker can [canister] application makes spot applications easy [simple]
- Innovative shaker [can] [canister] for easy [simple] application
- Shaker can [canister] allows for targeted applications
- Shaker can [canister] [ideal] [excellent] for application to small properties and micro lots
- [Kills] [Controls] yellow nutsedge
- [Kills] [Controls] purple nutsedge
- [Kills] [Controls] yellow nutsedge and purple nutsedge without damaging most lawn grasses when used as directed
- [Kills] [Controls] yellow nutsedge, purple nutsedge, kyllinga species, and listed broadleaf weeds
- For use on cool-season and warm-season grasses
- For use on [residential] lawns
- [For best results], apply when weeds are actively growing
- May be applied when weed leaves are wet or dry
- Apply to weed leaves when wet or dry
- For best results, water-in granules within 48 hours after treatment
- Getting Started
- Kills
- Controls
- Where to use/apply
- How much to use/apply
- How to use/apply
- When to use/apply
- Amount to use
- Coverage
- For questions or comments call toll-free 800.884.3179 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST-or visit: GordonsProfessional.com
- Powered by Vexis [Technology]

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-01024.20220322.amend-proposed-clean.doc

2. Reason for Issue: Update weed list per latest efficacy data, add application equipment instructions, shaker can reuse instructions, Storage & Disposal, EPA Comments, Incorporate Notification 4/5/2021.