



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

2217-1065

Date of Issuance:

6/23/22

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

EH-1672 Herbicide

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation
P.O. Box 860350
Shawnee, KS 66286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Continues page 2

Signature of Approving Official:

Mindy Ondish, Product Manager 23
Herbicide Branch, Registration Division (7505T)

Date:

6/23/22

2. You are required to comply with the data requirements described in the generic data call-in (GDCI) identified below:
 - a. Fluazifop-P-butyl GDCI-122809-1513

You must comply with all of the data requirements within the established deadlines. If you have questions about the GDCI listed above, you may contact the Chemical Review Manager in the Pesticide Re-Evaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirement for storage stability (Guideline 830.6317) is not satisfied. You have 18 months from the date of registration to provide this data.
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.


If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 11/13/2020
- Alternate CSF 1 dated 11/13/2020
- Alternate CSF 2 dated 11/13/2020
- Alternate CSF 3 dated 11/13/2020
- Alternate CSF 4 dated 11/13/2020
- Alternate CSF 5 dated 11/13/2020
- Alternate CSF 6 dated 11/13/2020

If you have any questions, please contact Endia Blunt at 202-566-2505 or at blunt.endia@epa.gov.

Enclosure



Fluazifop-p-butyl	GROUP	1	HERBICIDE
Imazapyr	GROUP	2	HERBICIDE
2,4-D Dicamba	GROUP	4	HERBICIDE
Pyraflufen-ethyl	GROUP	14	HERBICIDE

EH-1672 HERBICIDE

EPA Reg. No. 2217-1065

ACTIVE INGREDIENTS:

Pyraflufen-ethyl	0.0079%
Imazapyr, isopropylamine salt	1.74 %
Fluazifop-p-butyl	0.63 %
2,4-D, 2-ethylhexyl ester	3.02 %
Dicamba	0.51 %
OTHER INGREDIENTS:	94.0921%
TOTAL	100.0000%

THIS PRODUCT CONTAINS:

- 0.00067 lb Pyraflufen-ethyl per gallon or 0.0079%
- 0.12 lb Imazapyr acid per gallon or 1.42%
- 0.053 lb Fluazifop-p-butyl per gallon or 0.63%
- 0.17 lb 2,4-D acid per gallon or 2.00%
- 0.043 lb Dicamba acid per gallon or 0.51%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See [attached booklet] [inside pages] for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.

Net Contents: _____

EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286



PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks,
- chemical-resistant gloves made of barrier laminate, butyl rubber (≥ 14 mils), nitrile rubber (≥ 14 mils), neoprene rubber (≥ 14 mils), polyvinyl chloride (PVC) (≥ 14 mils) or viton (≥ 14 mils)

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing• Rinse skin immediately with plenty of water for 15-20 minutes• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may contact the poison control center toll free at 1-877-800-5556 for emergency medical treatment information.

Environmental Hazards

This pesticide is toxic to plants, fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. Do not apply when weather conditions favor drift from treated areas.

Groundwater Advisory: These chemicals (2,4-D, Fluazifop-p-butyl and dicamba) have properties and characteristics associated with chemicals detected in groundwater. These chemicals may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Application around a cistern or well may result in contamination of drinking water or groundwater.

Non-Target Organism Advisory: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of fluazifop-P-butyl from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Physical or Chemical Hazards

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

EH-1672 HERBICIDE is a nonselective herbicide that kills most annual and perennial grasses, broadleaf weeds, vines, brush, and trees. The product is absorbed through leaves, stems, and roots and translocates throughout the plant. Treated plants stop growing 1 to 2 weeks after application. EH-1672 HERBICIDE will also provide residual control for up to 12 months in the treated area. Heavy rainfall will reduce the length of control. Heavy clay soils may increase length of control.

2. Use Restrictions

- Do not enter or allow people or pets to enter treated areas until sprays have dried. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Only protected handlers may be in the area during application. Keep people and pets out of the area during application.
- Do not use on feed or food crops.
- Do not allow livestock to graze on any areas treated with this product.
- Use only in areas where you want to 1) kill all existing vegetation and 2) suppress regrowth for up to 12 months.
- Do not use in gardens, flower beds or in areas intended to be seeded or planted within 12 months.
- Do not use on indoor plants including interior plantscapes. For outdoor use only.
- Do not spray in mulch around newly planted and established trees. Do not use in areas where the spray solution may contact the root system of desirable plants. [See also Section 9: Avoiding Injury to Non-Target Plants.]
- Do not treat aquatic plants.
- For ground application only; aerial application is prohibited.
- Chemigation: Do not apply this product through any type of irrigation.
- In lawns, do not use for individual weed control as this product will suppress the growth of all vegetation, for up to 12 months.
- Thoroughly flush application equipment with water after using this product. Do not flush equipment near desirable plants or their root zones.
- Do not apply more than a total of 36 fl.oz. per 1000 sq.ft. per year, the equivalent of, 2.08 lb 2,4-D ae, 1.47 lb Imazapyr ae, 0.65 lb Fluazifop-p-butyl, 0.53 lb dicamba ae, and 0.008 lb Pyraflufen-ethyl per acre per year.
- Not for use on turf or plants being grown for sale or other commercial use, or for commercial seed or sod production, or for research purposes.
- **[New York:** Not for sale, distribution, or use in New York State.]

3. Weed Resistance Management

For resistance management, this product contains Group 1, 2, 4 and 14 herbicides. Any weed population may contain or develop plants naturally resistant to this product and other Group 1, 2, 4 or 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of this product or other Group 1, 2, 4 or 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or pest control advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural (e.g., timing to favor vegetation control), biological (weed-competitive varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- For further information or to report suspected resistance, contact PBI-Gordon Corporation at 800-884-3179.

4. Mandatory Spray Drift Management

Boomless Ground Applications:

- Applicators are required to use a fine or coarser droplet size (ASABE S572) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Temperature and Humidity: When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions: Drift potential is high during temperature inversions. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind: Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications: Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications: Take precautions to minimize spray drift

5. Where To Use

EH-1672 HERBICIDE may be used to control vegetation in: [In fencerows, hedgerows, agricultural uncultivated areas, shelter belts, farm yards, and farm premises. Around domestic (residential) dwellings, outdoor buildings, structures and shelters such as beef barns, hog barns, dairy barns, stables, dog kennels and poultry houses. Along fences of paddocks, corrals, feedlots, and livestock holding areas. For the establishment and maintenance of wildlife openings. In roadways, roadsides, airports, landing fields, industrial sites, storage yards, lumber yards, tank farms, pumping installations, fuel tanks, pipelines, non-agricultural uncultivated areas or soils, paved areas (including crack and crevice), curbs, sidewalks, driveways, parking areas, brick walks, gravel paths, paths, patios, and vacant lots, urban areas, cemeteries, recreational areas, golf courses, play areas, and tennis courts. Along fences.]

6. How Much To Apply For Non-Brush Vegetation Control

Mix 18 fl.oz. of product in 3 to 5 gallons of water to cover 1000 sq.ft. A maximum of 2 broadcast or spot applications per year with a minimum of 30 days between applications. Add a surfactant to the spray solution to enhance performance. A second application may be necessary to control some species.

Use hand-operated or mechanical spray equipment such as backpack sprayers, pump-up sprayers, sprinkling can, hose-end sprayers, or pull-behind sprayers. Spray the area where bareground is desired. Spot treatments may be useful to control weed escapes after a broadcast treatment.

Example: Amount to mix when spraying at 5 gallons per 1000 sq.ft.			
	To make 1 gallon of finished spray solution	To make 5 gallons of finished spray solution	To make 25 gallons of finished spray solution
Amount of product to add to tank	3.6 fl.oz.	18 fl.oz.	90 fl.oz.
Coverage area	200 sq.ft.	1000 sq.ft.	5000 sq.ft.

For spot treatments made by mechanically-pressurized handguns to landscaping trees and shrubs follow the mixing directions above, so as to not exceed the maximum concentration of 0.01 lb Fluazifop-p-butyl per gallon of application solution or to fall below the minimum of 3 gallons of spray solution per 1000 sq. ft.

7. How Much To Apply For Brush Control

Mix 36 fl.oz. of product in 1 to 5 gallons of water. A maximum of 1 application per year. Add a surfactant to the spray solution to enhance performance.

Use hand-operated or mechanical spray equipment such as backpack sprayers, pump-up sprayers, sprinkling can, hose-end sprayers, or pull-behind sprayers. Uniformly spray the brush foliage and the soil directly under the brush. Excessive wetting of foliage is not recommended.

8. For Best Results

This product is absorbed through leaves, stems, and roots and moves throughout the plant. Treated plants stop growing soon after application. Residual control will last up to 12 months.

Speed of control: This product controls vegetation systemically. Treated plants stop growing 1 to 2 weeks after application. Control may take 3 to 4 weeks. Cooler weather may delay the effects of the herbicide.

Conditions that will reduce control: A reduction in effectiveness may occur when applications are made to vegetation under stress due to heat or drought. Heavy rainfall will reduce the length of residual control.

Conditions that will improve control: Make applications when daytime high temperatures are between 50°F and 85°F. Heavy clay soils may increase the length of residual control. The product is rainfast within 2 hours of application. Vegetation and brush should be growing vigorously for best results.

9. Avoiding Injury to Non-Target Plants

Movement of the product from the target plant or site may injure non-target plants. Follow these precautions to manage off-target movement.

Runoff: Do not over apply causing runoff from the treated foliage. Applications made to dry soils may not be absorbed thoroughly. Soil moisture from rainfall or irrigation prior to application may reduce runoff. Do not apply to slopes.

Spray Drift: Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Avoiding spray drift is the responsibility of the applicator. Applications with wind speeds greater than 10 mph are prohibited. Applications into temperature inversions are prohibited.

Roots of Trees and Shrubs: Untreated plants can be affected by root uptake. Do not use over the root zone of desirable trees and shrubs. Roots extend beyond the drip line and can extend under fences, driveways, patios, sidewalks, paths, and similar landscape elements.

Movement of Treated Soil: Do not treat powdery, dry or light sandy soils when the soil may be blown by wind. If necessary soil surface should be settled by rain or irrigation.

10. Vegetation Controlled

EH-1672 HERBICIDE will control or suppress most grasses, broadleaf weeds, woody brush, vines, and trees as listed. The degree of control and residual control varies by plant species.

[Vegetation table(s). A subset or all of the following weeds/vegetation may be included on the container label.

GRASSY WEEDS			
Annual bluegrass	Dallisgrass*	Japanese stiltgrass	St. Augustinegrass*
Annual ryegrass	Downy brome	Johnsongrass	Thin paspalum
Bahiagrass*	Fall panicum	Kentucky Bluegrass	Timothy
Beardgrass	Feathertop	Lovegrass	Torpedograss*
Bermudagrass*	Fescue*	Orchardgrass	Vaseygrass*
Big bluestem	Foxtail (green, giant, yellow)	<i>Panicum</i> spp.	Wild barley
Broadleaf signalgrass	Giant reed	Prairie cordgrass	Wild oats

Canada bluegrass	Globe sedge	Prairie three awn	Wirestem muhly
Centipedegrass	Goosegrass	Quackgrass	Witchgrass
Cheat	Green kyllinga	Reed canarygrass	Woolly cupgrass
Cogongrass	Guineagrass	Sand dropseed	Yellow nutsedge
Crabgrass	Italian ryegrass	Sandbur	Zoysiagrass*
Crowfootgrass	Itchgrass	Smooth brome	

* sequential application may be necessary

BROADLEAF WEEDS			
Arrowwood	Fiddleneck	Nettleleaf goosefoot	Smartweed
Blackberry briar	Field madder	Oxalis	Sorrell
Broadleaf Plantain	Filaree	Oxeye daisy	Sowthistle
Broom snakeweed	Fleabane	Pepperweed	Spotted spurge
Bull thistle	Goldenrod	Pigweed	Spurge, annual
Burclover	Gray rabbitbrush	Plantain	Stinging nettle
Burdock	Ground ivy	Pokeweed	Sunflower
Bushy aster	Hairy buttercup	Prickly sida	Sweet clover
California burclover	Henbit	Primrose	Tansymustard
Camphorweed	Horseweed	Puncturevine	Texas thistle
Canada thistle*	Hoary vervain	Purple deadnettle	Velvetleaf
Carolina Geranium	Indian mustard	Purple loosestrife	Virginia buttonweed
Carpetweed	Japanese	Purple vetch	White clover
Chickweed, common	bamboo/knotweed	Purslane	Wild carrot
Chickweed, mouse ear	Knotweed, prostrate	Pusley, Florida	Wild lettuce
Cocklebur	Kochia	Ragweed, common	Wild mustard
Cudweed	Lambsquarters	Ragweed, giant	Wild parsnip
Dandelion	Lawn burweed	Redroot pigweed	Wild strawberry
Desert camelthorn	Lespedeza	Rocket, London	Wild turnip
Dichondra	Little mallow	Rush skeletonweed	Wild violet
Diffuse knapweed	Lyre-leaf sage	Russian knapweed	Woollyleaf bursage
Dock	Milkweed	Russian thistle	Yellow starthistle
Dogfennel	Miner's lettuce	Saltbush	Yellow woodsorrel
Dollarweed	Mullein	Shepherd's purse	
		Silverleaf nightshade	

* sequential application may be necessary

VINES			
Field bindweed	Morningglory	Virginia creeper	Wild rose* (Including: Multiflora and Macartney rose)
Hedge bindweed	Poison ivy	Wild buckwheat	
Honeysuckle*	Redvine	Wild grape	
Kudzu	Trumpet creeper		

* sequential application may be necessary

WOODY BRUSH AND TREES			
Alder	Chinaberry	Madrone	Sassafras
American beech	Chinese tallow-tree	Maple	Sourwood
Ash	Chinquapin	Melaleuca	Sumac
Aspen	Cottonwood	Mesquite	Sweetgum
Autumn olive	Cypress	Mulberry	Sycamore
Bald cypress	Dogwood	Oak	Tanoak
Bigleaf maple	Eucalyptus	Persimmon	Titi
Birch	Hawthorn	Poison oak	Tree of heaven
Black oak	Hickory	Popcorn-tree	<i>Vaccinium</i> spp.
Blackgum	Huckleberry	Poplar	(Including: Blueberry and Sparkleberry)
Boxelder	<i>Lyonia</i> spp. (Including: Fetterbush and Staggerbush)	Privet	Willow
Brazilian peppertree		Red alder	Yellow-poplar
Ceanothis		Red maple	
Cherry		Saltcedar	

End optional weed list table.]

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

[For Plastic Containers – Nonrefillable with capacities equal to or less than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

[For Plastic Containers – Nonrefillable with capacities greater than 5 gallons:]

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Triple rinse [or pressure rinse] container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

[Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.]

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED

WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner or use of application, etc. Such factors and conditions are beyond the control of the manufacturer, and **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** To the extent consistent with applicable law, buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OF HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THE PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this LIMITED WARRANTY AND DISCLAIMER in any manner.

APPENDIX

Information that may optionally be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

[Back Panel]

Getting Started [What It Does/Kills/Controls]:

Kills and prevents: listed [weeds, grass, brush, and vegetation] for up to one year

Where To Use:

Along fences, fencerows, hedgerows, driveways, patios, sidewalks, around buildings, curbs, parking areas, brick walks, gravel paths, walkways, paths

Apply to any area where no vegetation is desired

When To [Use/Apply]:

Apply in spring, summer, or fall.

[Amount To Use/How Much To [Use/Apply]]:

Amount to mix when spraying at 5 gallons per 1000 sq.ft.			
	To make 1 gallon of finished spray solution	To make 5 gallons of finished spray solution	To make 25 gallons of finished spray solution
Amount of product to add to tank	3.6 fl.oz.	18 fl.oz.	90 fl.oz.
Coverage area	200 sq.ft.	1000 sq.ft.	5000 sq.ft.

[Coverage:]

1 quart covers 1,777 sq.ft.

1 gallon covers 7,100 sq.ft.

2.5 gallon covers 17,750 sq.ft.

Additional claims and label statements:

- Offers long-term weed [vegetation] control in places where no growth is wanted
- Kills [Controls] [unwanted] [tough] [weeds], [vegetation], [brush], [plants], and [grasses] as listed
- Prevents [weeds and] [vegetative] [new] growth for up to 1 year.
- Apply only where no vegetation growth is wanted for up to 1 year.
- Kills [Controls] [most common][unwanted] vegetation as listed
- Year-long control of tough weeds & brush [vegetation] as listed
- Kills and prevents listed weeds, grass and brush [vegetation] for up to 12 months
- Goes to the roots to stop growth fast
- [Where to Use:] Apply where no vegetation is desired in areas such as fencerows, driveways, around residential buildings, propane tanks, around residential storage areas, parking areas, patios, and pathways, [and more areas listed on label].
- Kills: tough weeds, brush and grasses [vegetation] [as listed] and prevents new growth for up to 1 year
- Amount to use: Mix 18 fl.oz. of product with 3-5 gallon of water to cover 1,000 sq.ft.
- Kills and prevents listed weeds, grass and brush [vegetation] for up to [12 months] [1 year]
- Kills & prevents listed vegetation for up to 1 year
- Results in hours
- Kills the root
- Offers long-term weed [vegetation] control
- Kills unwanted weeds, brush and grass [vegetation] as listed
- Year-long control
- Year-long control of tough weeds and brush [vegetation] as listed
- Kills listed vegetation – weeds, grasses, and brush
- Visible results in hours
- Works in hours
- Kills unwanted weeds [vegetation] as listed
- Same [dependable] [reliable] [trusted] [effective] results* *when compared to Barrier Year-Long Vegetation Killer Concentrate
- [New] [Updated] [improved] [enhanced] Formulation* *when compared to Barrier Year-Long Vegetation Killer Concentrate [Note: “New” may only be used for 6 months following commercial launch]
- Same year-long control [performance]* *when compared to Barrier Year-Long Vegetation Killer Concentrate
- Improved Formulation – Same Dependable Results** **when compared to Barrier Year-Long Vegetation Killer Concentrate
- 18 fl.oz. of product treats up to 1,000 sq.ft
- 1 quart covers 1,777 sq.ft.
- 1 gallon covers 7,100 sq.ft.
- 2.5 gallon covers 17,750 sq.ft.

- For driveways, patios, fence rows, and other places where no vegetation is desired.
- For patios, fence rows, and other listed places where no vegetation is desired
- Use Where No Growth Is Desired
- Apply to any area where no vegetation is desired
- Fencerows, Driveways, And More Sites As Listed . . . Around Residential Buildings, Residential Storage Areas, Patios, Walkways, Sidewalks
- Getting Started
- Amount to Use: Mix 18 fl.oz. of product with 3-5 gallons of water to cover 1000 sq.ft.
- [Telephone Icon][Computer Icon] For questions or comments call toll-free 1-800-884-3179 Mon. – Fri. 8:00 a.m. – 4:30 p.m. CST or visit: GordonsUSA.com
- Rainproof [Rainfast] in 2 hours
- Guaranteed* to work or your money back [*If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.]
- Questions or Comments? Call 1-800-322-5553
- NOTICE: To the extent required by applicable law, buyer assumes all responsibility for safety and use not in accordance with directions



Add icons:

Length of Control:



Use sites:



Concentrated:



Applicator Claims:

- Easy to use
- Easy to use sprayer
- Attach to hose & spray

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-0XXXX.20220622.new-proposed-EH1672 - clean.doc

2. **Reason for Issue:** New Registration

Internal Usage Notes