

# 20% TOXAPHENE COTTON DUST

**ACTIVE INGREDIENTS:**

**TOXAPHENE** ..... 20.00%

[Technical Chlorinated Camphene (Chlorine Content 67 to 69%)]

**INERT INGREDIENTS** ..... 80.00%

**TOTAL** ..... 100.00%

**THIS DUST WILL PASS 90% THROUGH A 325 MESH SCREEN**

## SUGGESTIONS FOR USE

This material when used on cotton should be applied at the rate of 10 to 20 pounds per acre for controlling the Cotton Boll Weevil, Cotton Bollworm, Cotton Fleahopper, and certain other cotton insect pests. For best results, dust when there is little or no wind, preferably in late afternoon. Use good hand or power dusting equipment. Consult your local county or state agricultural authorities for further information concerning the use of this material.

## CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid inhalation or contamination of feed and foodstuffs. Wash exposed parts of body thoroughly with soap and warm water. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Keep away from children and domestic animals. Do not feed treated cotton trash to dairy animals or animals being finished for slaughter. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season. Highly Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

USDA REG. NO. 2269-013

**ACCEPTED**  
**MAR 13 1967**

UNDER THE FEDERAL INSECTICIDE  
 ACT AND THE FEDERAL  
 FOOD, DRUG AND COSMETIC  
 ACT  
 22104-19

