

10% SEVIN* DUST

ACTIVE INGREDIENTS:	BY WT.
CARBARYL (1-Naphthyl N-Methylcarbamate)	10.00%
INERT INGREDIENTS	90.00%
	<u>Total 100.00%</u>

*Trade Mark of Union Carbide Corp. for CARBARYL (1-NAPHTHYL N-METHYLCARBONATE)

DIRECTIONS

Pesticide products used contrary to directions may cause serious plant or personal injury.

COTTON—Against early season pests such as Thrips and Cotton Fleahopper use 5 to 10 lbs. per acre every 5 to 10 days as needed beginning when insects increase to injurious numbers.

Against mid-season and late-season Boll Weevil, Cotton Bollworm, Cotton Leafworm, Lygus Bugs and Cotton Leaf Perforator, use 10 to 20 lbs. per acre, the amount depending upon plant size and the severity of infestation. Begin dusting on a 5-day schedule when infestations reach injurious numbers.

APHIDS are usually suppressed by a regular Sevin schedule. Do not apply after bolls open. Do not graze treated fields or feed cotton trash to dairy or meat animals.

CAUTION

HARMFUL IF INHALED OR SWALLOWED. Do Not Take Internally. Avoid Breathing of Dust. *Skin Contact May Be Harmful.* Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Work. *Note For Physician:* Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Burn empty bags immediately. Stay out of smoke and bury ashes and residue.

EPA Reg. No. 2269-74 2M 9/73

Gold Kist, Inc.
Atlanta, Ga. 30301

ITEM NO. 4000237

ACCEPTED
2269-74
AUG 9 1974
UNDER THE FEDERAL PESTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

2269-74

08/09/1974

1/1