

10% SEVIN* DUST

ACCEPTED

8-28-74

ACTIVE INGREDIENTS:

CARBARYL (1-Naphthyl N-Methylcarbamate)

BY WT.

10.00%

INERT INGREDIENTS

~~UNDER THE FEDERAL INSECTICIDE~~

90.00%

~~ACTING UNDER THE FEDERAL INSECTICIDE~~

Total 100.00%

FOR ECONOMIC FOREIGN REGISTRATION

NO TRADE NO. SUBJECT

TO APPLICABLE REGULATIONS

*Trade mark of Union Carbide Corp. for CARBARYL (1-NAPHTHYL N-METHYLCARBONATE)

DIRECTIONS

Pesticide products used contrary to directions may cause serious plant or personal injury.

COTTON—Against early season Thrips and Cotton Fleahopper use 5 to 10 lbs. per acre every 5 to 10 days as needed beginning when insects increase to injurious numbers.

Against mid-season and late season Boll Weevil, Cotton Bollworm, Cotton Leafworm, Lygus Bugs and Cotton Leaf Perforator, use 10 to 20 lbs. per acre, the amount depending upon plant size and the severity of infestation. Begin dusting on a 5-day schedule when infestations reach injurious numbers.

~~Do not apply to plants or animals. Do not apply to plants or animals. Do not apply to plants or animals. Do not apply to plants or animals.~~

To control the insects specified on the crops listed below, apply this dust uniformly and thoroughly to upper and lower surfaces of all foliage. Dust coverage of foliage should be complete, but do not apply in excessive amounts. Begin applications when insects or their damage first appear. Repeat is necessary, generally at 7-day intervals unless shorter interval is stated. May be applied on day of harvest unless otherwise indicated.

BEANS—to control Mexican bean beetle use 5 lb. per acre, to control bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle adults and leafhoppers, use 10 pounds per acre.

CUCUMBERS, MELONS, PUMPKINS AND SQUASH—to control pickleworm, melonworm use 5 lb. per acre, to control cucumber beetles, flea beetles, leafhoppers and squash bug, use 10 pounds per acre. Avoid excessive applications. Some leaf injury may occur if treatments are made when foliage is wet or during periods of prolonged high humidity. Do not use on watermelons.

POTATOES, TOMATOES, EGGPLANT AND PEPPERS—to control Colorado potato beetle, flea beetles, leafhoppers use 10 lb. per acre, to control European corn borer, fall army-worm, tomato fruitworm, tomato hornworm, larnished plant bug and stink bug, use 20 pounds per acre.

CABBAGE, BROCCOLI AND CAULIFLOWER—to control flea beetles, Harlequin bug, use 10 lbs. per acre, to control corn earworm, use 20 pounds per acre. Do not apply within 3 days of harvest.

POULTRY

CHICKENS, PARTRIDGES, PHEASANTS AND PIGEONS—Northern fowl mite, chicken red mite, lice, fleas, and bed bugs in houses or quarters. Apply 1 pound per 80 sq. ft. of floor and litter, and to roosts and interior surfaces. Do not apply directly to eggs or nest litter. Do not house birds in treated houses or areas within 7 days of slaughter. For Northern fowl mite and lice on the birds, as a supplement to treatment of premises, apply 1 pound per 200 birds. Direct dust to vent and fluff areas, and distribute throughout feathers. Do not repeat within 4 weeks. Do not apply within 7 days of slaughter. DO NOT CONTAMINATE FEED OR DRINKING WATER.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF INHALED OR SWALLOWED. Do Not Take Internally. Avoid Breathing of Dust. Skin Contact May Be Harmful. Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Note For Physician. Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine in antidotal.

"This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service."

Burn empty bags immediately. Stay out of smoke and bury ashes and residue.

EPA Reg. No. 2269-74
EPA EST 2269-GA-1

Gold Kist, Inc.
Atlanta, Ga. 30301
SM-6 74

Item No. 4000237-50 lb. bag
Item No. 4000238-25 lb. bag