

5.0% SEVIN* DUST

ACTIVE INGREDIENTS		BY WT.
CARBARYL (1-Naphthyl N-Methylcarbamate)	5.0%
INERT INGREDIENTS:	95.0%
	Total	100.0%

ACCEPTED

CAUTION AUG 6 1974

TO ATTACHED COMMENTS.

*Sevin, Trade Mark of UCC for Carbaryl 1-Naphthyl N-Methylcarbamate

HARMFUL IF INHALED OR SWALLOWED. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. Skin Contact May Be Harmful. Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Use. Note For Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine Antidote.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

DIRECTIONS UNDER NO. _____

To Control the insects specified on the corps listed below, apply this dust uniformly and thoroughly to upper and lower surfaces of all foliage. Dust coverage of foliage should be complete, but do not apply in excessive amounts. Begin applications when insects or their damage first appear. Repeat as necessary, generally at 7-day intervals unless shorter interval is stated. May be applied on day of harvest unless otherwise indicated.

- BEANS** — Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetles, Japanese beetle adults and leafhoppers.
- CUCUMBERS, MELONS, PUMPKINS AND SQUASH** — Pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bug. Avoid excessive applications. Some leaf injury may occur if treatments are made when foliage is wet or during periods of prolonged high humidity. Do not use on watermelons.
- POTATOES, TOMATOES, EGGPLANT AND PEPPERS** — Colorado potato beetle, flea beetles, leafhoppers, European corn borer, fall armyworm, tomato fruitworm, tomato hornworm, tarnished plant bug and stink bug.
- CABBAGE, BROCCOLI AND CAULIFLOWER** — Flea beetles, Harlequin bug, cabbage caterpillars, and corn earworm. For cabbage looper, apply at 4 to 5-day intervals. Do not apply within 3 days of harvest.

POULTRY

CHICKENS, PARTRIDGES, PHEASANTS AND PIGEONS — Northern fowl mite, chicken red mite, lice, fleas, and bed bugs in houses or quarters. Apply 1 pound per 40 sq. ft. of floor and litter, and to roosts and interior surfaces. Do not apply directly to eggs or nest litter. Do not house birds in treated houses or areas within 7 days of slaughter. For Northern fowl mite and lice on the birds, as a supplement to treatment of premises, apply 1 pound per 100 birds. Direct dust to vent and fluff areas, and distribute throughout feathers. Do not repeat within 4 weeks. Do not apply within 7 days of slaughter. **DO NOT CONTAMINATE FEED OR DRINKING WATER.**

SPECIAL NOTES

The rate of application described as uniform and thorough dust coverage on upper and lower surfaces of all foliage is equivalent to 20 pounds per acre on cucumbers, melons, pumpkins and squash. 40 pounds per acre on beans, potatoes, tomatoes, eggplant, peppers, cabbage, broccoli, and cauliflower.