

ACTIVE INGREDIENTS:

BY WT.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

DO NOT EAT OR DRINK WHILE APPLICATING. STAY OUT OF SIGHT.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

HARMFUL IF INHALED OR SWALLOWED. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. Skin Contact May Be Harmful. Avoid Contact. Wash Hands Before Eating. Take Shower or Bath After Work.

Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidote.

For protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

DIRECTIONS

PEANUTS: BLISTER BEETLES, MEXICAN BEAN BEETLE — 10 to 20 lbs. per acre. THIRIPS, LEAFHOPPERS, THREE-CORNERED ALFALFA HOPPER, CUCUMBER BEETLES, BEAN LEAF BEETLE, JAPANESE BEETLE, ALFALFA CATERPILLAR, VELVET BEAN CATERPILLAR — 20 lbs. per acre.

ARMYWORM, WEBWORMS, CORN EARWORM, STINK BUGS — 20 to 30 lbs. per acre.

GRASSHOPPERS — 10 to 30 lbs. per acre. May be applied up to the day of harvest or greening.

**COTTON PRODUCERS ASSOCIATION
Atlanta, Georgia**

IM-20

ACCEPTED

JUL 21 1970

Under the Federal Insecticide, Fungicide and Rodenticide Act

ESTATE WILSON CO.