

# ZINEB DUST No. 10

(For Tobacco Blue Mold Control)

**ACTIVE INGREDIENTS:**

Zineb (Zinc Ethylene Bisdithiocarbamate)  
(Metallic Zinc equivalent 1.54%)

**INERT INGREDIENTS**

**TOTAL**

100.00%

**DIRECTIONS FOR USE ON TOBACCO PLANT BEDS**

Begin applications as soon as the first signs of blue mold are observed. Apply 100 lbs. of dust per acre. A second application may be made 10 to 14 days later. Applications should be made when plants are in the early stages of growth. Do not apply to plants showing a uniform maturity. Apply as soon as possible. Usually 2 to 3 applications are required. For the best results, apply at intervals of 10 to 14 days. All other applications should be made at intervals of 10 to 14 days.

**DIRECTIONS FOR USE ON OTHER CROPS**

Apply 100 lbs. of dust per acre. A second application may be made 10 to 14 days later. Applications should be made when plants are in the early stages of growth. Do not apply to plants showing a uniform maturity. Apply as soon as possible. Usually 2 to 3 applications are required. For the best results, apply at intervals of 10 to 14 days. All other applications should be made at intervals of 10 to 14 days.

**CAUTION**

KEEP OUT OF REACH OF CHILDREN

Atlanta, Ga. 30301