

5% CUTWORM BAIT

ACTIVE INGREDIENT (Contains 5% Dylox®)
Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate†
INERT INGREDIENTS

†DYLOX, a registered trademark of the German Proprietor, Leverkusen.

ACCEPTED
8-2-74
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
UNDER NO. _____ SUBJECT
ATTACHED DOCUMENTS.

5%
95%
100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. DO NOT BREATHE DUST. Avoid contact with skin. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

To Physician: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DO NOT APPLY 5% CUTWORM BAIT TO OR ALLOW TO DRIFT ONTO VARIETIES OF SORGHUM WHICH ARE SENSITIVE TO PHOSPHATES TO DO SO MAY CAUSE BURNING OF THE SORGHUM OR MILO.

Do not use on other crops used for food or forage. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes. This product is toxic to birds and other wildlife. Do not contaminate water by cleaning of equipment or disposal wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

DIRECTIONS FOR USE

APPLY 5% Cutworm Bait at specified rate per acre for the control of Beet Armyworm, Western Yellow-striped Armyworm, Variegated Cutworm, Black Cutworm, Greasy Cutworm, Granulated Cutworm, Western Bean Cutworm, Mole Crickets, Salt-marsh Caterpillar. Controls surface feeding cutworms only.

If field has a history of previous damage or if cutworms are suspected, apply 5% Cutworm Bait before or soon after planting to obtain best results. Protect your growing plants by applying a thin film of 5% Cutworm Bait over infested area. **AVOID APPLYING DIRECTLY TO EDIBLE PARTS. USE ONLY AS RECOMMENDED!**

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

RECOMMENDATIONS

VEGETABLES

BEANS (Dry and Snap): Use 20 to 30 lb. per acre. Repeat as necessary. Do not apply within 14 days of harvest. Do not use treated vines as food for dairy or meat animals. Apply when plants are small for best results.

LIMA BEANS: Use 20 to 30 lb. per acre. Do not apply more than twice per season. Do not harvest for food, feed or forage within 30 days of treatment.

TOMATOES: Use 20 lb. per acre. Repeat as necessary. Do not apply within 21 days of harvest.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Use 20 lb. per acre. Repeat as necessary. Do not apply within 21 days of harvest. Avoid application directly to edible parts.

COWPEAS, SOUTHERN PEAS, BLACK-EYED PEAS, CROWDER PEAS: Use 20 to 30 lb. per acre. Apply only twice per season. Do not use treated vines as food for dairy or meat animals. Do not apply within 14 days of harvest.

PUMPKIN: Apply only once per season and not within 14 days of harvest. Use 20 to 30 lb. per acre.

COLLARDS, LETTUCE: Use 20 lb. per acre. Repeat as necessary until heads begin to form. Do not apply within 28 days of harvest.

FIELD CROPS

ALFALFA: Use 20 lb. per acre. One application per cutting. Do not apply within 14 days of cutting or grazing.

COTTON: Use 30 lb. per acre. Repeat as necessary. Do not pasture treated fields within 14 days of application. Do not apply within 7 days of picking.

PEANUTS: Use 20 to 30 lb. per acre. Treat as needed when worm infestations are suspected.

TOBACCO: Use 20 to 30 lb. per acre. Do not apply directly into bud, but apply to base of plants.

SEED FIELD CROPS

SOYBEANS: Use 20 to 30 lb. per acre. Repeat as necessary. Do not pasture.

EPA REG. NO. 2269-154

EPA EST. 2269-GA-1

1M 4 74

Manufactured by
GOLD KIST INC., Atlanta, Georgia 30301

ITEM NO. 4000158

5% CUTWORM BAIT

ACTIVE INGREDIENT

(Contains 5% Dylox®)

Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate †

INERT INGREDIENTS

®DYLOX, a registered trademark of the German Proprietor, Leverkusen.

ACCEPTED

8-2-74

5%
95%
100%

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT
FOR ECONOMIC PESTS REGISTERED
UNDER NO. _____ SUBJECT
ATTACHED COMMENTS.

CAUTION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN, DO NOT BREATHE DUST. Avoid contact with skin. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food.

If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

To Physician: Atropine sulfate is antidotal. 2 PAM is also antidotal and may be administered in conjunction with atropine.

DO NOT APPLY 5% CUTWORM BAIT TO OR ALLOW TO DRIFT ONTO VARIETIES OF SORGHUM WHICH ARE SENSITIVE TO PHOSPHATES. TO DO SO MAY CAUSE BURNING OF THE SORGHUM OR MILO.

Do not use on other crops used for food or forage. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes. This product is toxic to birds and other wildlife. Do not contaminate water by cleaning of equipment or disposal wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Keep out of lakes, streams or ponds. Apply this product only as specified on this label.

DIRECTIONS FOR USE

APPLY 5% Cutworm Bait at specified rate per acre for the control of Beet Armyworm, Western Yellow-striped Armyworm, Variegated Cutworm, Black Cutworm, Greasy Cutworm, Granulated Cutworm, Western Bean Cutworm, Mole Crickets, Salt-marsh Caterpillar. Controls surface feeding cutworms only.

If field has a history of previous damage or if cutworms are suspected, apply 5% Cutworm Bait before or soon after planting to obtain best results. Protect your growing plants by applying a thin film of 5% Cutworm Bait over infested area. **AVOID APPLYING DIRECTLY TO EDIBLE PARTS. USE ONLY AS RECOMMENDED!**

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

RECOMMENDATIONS

VEGETABLES

BEANS (Dry and Snap): Use 20 to 30 lb. per acre. Repeat as necessary. Do not apply within 14 days of harvest. Do not use treated vines as food for dairy or meat animals. Apply when plants are small for best results.

LIMA BEANS: Use 20 to 30 lb. per acre. Do not apply more than twice per season. Do not harvest for food, feed or forage within 30 days of treatment.

TOMATOES: Use 20 lb. per acre. Repeat as necessary. Do not apply within 21 days of harvest.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Use 20 lb. per acre. Repeat as necessary. Do not apply within 21 days of harvest. Avoid applying directly to edible parts.

COWPEAS, SOUTHERN PEAS, BLACK-EYED PEAS, CROWDER PEAS: Use 20 to 30 lb. per acre. Apply only twice per season. Do not use treated vines as food for dairy or meat animals. Do not apply within 14 days of harvest.

PUMPKIN: Apply only once per season and not within 14 days of harvest. Use 20 to 30 lb. per acre.

COLLARDS, LETTUCE: Use 20 lb. per acre. Repeat as necessary until heads begin to form. Do not apply within 28 days of harvest.

FIELD CROPS

ALFALFA: Use 20 lb. per acre. One application per cutting. Do not apply within 14 days of cutting or grazing.

COTTON: Use 30 lb. per acre. Repeat as necessary. Do not pasture treated fields within 14 days of application. Do not apply within 7 days of picking.

PEANUTS: Use 20 to 30 lb. per acre. Treat as needed when worm infestations are suspected.

TOBACCO: Use 20 to 30 lb. per acre. Do not apply directly into bud, but apply to base of plants.

SEED FIELD CROPS

SOYBEANS: Use 20 to 30 lb. per acre. Repeat as necessary. Do not pasture.

EPA REG. NO. 2269-154

EPA EST. 2269-GA-1

Manufactured by

GOLD KIST INC., Atlanta, Georgia 30301

ITEM NO. 4000158

1M/4/74