


TOXAPHENE EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:

Toxaphene *
Xylene range aromatic solvent
INERT INGREDIENTS

*Technical chlorinated camphene containing 67% to 69% chlorine.



	60%
	35%
	5%
Total	100%

CONTAINS 6 POUNDS OF TOXAPHENE PER GALLON

DIRECTIONS FOR USE

Toxaphene Emulsifiable Concentrate should be diluted or mixed with water for application as a spray. To mix, pour correct amount of concentrate into known volume of water in spray tank and stir or agitate thoroughly. If allowed to stand after mixing, re-agitate thoroughly before use. Some agitation is required during application.

COTTON—To control boll weevil, bollworm (does not include pink bollworm), fleahopper, thrips, lygus bug and other mirids, cutworms, and leafworm. Use $\frac{1}{3}$ to $\frac{2}{3}$ gal. (2 to 4 lbs. toxaphene) in the amount of water required for uniformly thorough spraying of 1 acre of cotton. Apply at the lower rate during the early season and at the higher rate during and after the blooming period.

For cutworms, apply when worms appear during early season, usually at 2 to 4-leaf stage; repeat when necessary. For thrips and fleahoppers, begin at 2 to 4-leaf stage and repeat at 7-day intervals if needed. For lygus bug and other mirids, apply when damage first appears; repeat as needed. For bollworm, apply when eggs or 3 to 4 small larvae per 100 terminal buds are found; repeat application at weekly intervals until infestation is controlled. Late season control of bollworm may require applications to be repeated at 4 to 5-day intervals as needed.

For boll weevil, during early season, apply at 2 to 4-leaf stage or at first squaring, if weevils are present, and make 3 to 4 applications at weekly intervals as necessary; during mid and late-season, apply when 10 percent of the squares or bolls are punctured and repeat at 4 to

5-day intervals as necessary. For leafworm, apply when worms appear, usually during the late season or maturing period, and repeat as needed.

DO NOT GRAZE DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER IN FIELD TREATED LATE IN THE SEASON.

**DO NOT USE, POUR, SPILL
OR STORE NEAR HEAT
OR OPEN FLAME**

MAY BE FATAL IF SWALLOWED! Do not breathe spray mist. Do not get on skin or clothing, or in eyes. In case of skin contact, wash thoroughly with soap and water; for eyes, flush with plenty of water and get medical attention. Wash thoroughly after using. Do not contaminate feed and foodstuffs. Not for use or storage in or around the home.

Highly toxic to fish & wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate any body of water nor apply to any area not specified on the label.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

This product is toxic to bees and should not be applied when bees are actively visiting the area.