


[2269-168]

2269-168



DIPEL 150 DUST*

(*Bacillus thuringiensis*)

CAUTION

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS

Bacillus thuringiensis, Berliner, Potency of 320 International Units per mg. (at least 9.5 billion viable spores per g.)*.....0.064%

INERT INGREDIENTS 99.936%

*Equivalent to potency of 0.15 billion International per pound of this product.

REASONABLE CARE SHOULD BE EXERCISED TO AVOID INHALATION OR CONTACT WITH OPEN WOUNDS. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

E.P.A. Registration No. 2269-168
E.P.A. Est. 2269 GA 1

DIRECTIONS FOR USE

DIPEL 150 Dust is a biological insecticide intended for use against chewing larvae (caterpillars) of certain lepidoptercous species.

DIPEL 150 Dust is recommended for use by ground or by air. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

APPLICATION RATE

For the control of Cabbage Looper and Imported Cabbage Worm on Broccoli, Cabbage, Cauliflower, Collards, Kale, Mustard Greens, apply 25 to 40 lbs. per acre.

For the control of Cabbage Looper on Celery, Lettuce, Melons, Potatoes, Spinach, Tomatoes and Cucumbers (Southeastern U.S. only) apply 25 to 40 lbs per acre.

For the control of Hornworm on Tobacco and Tomatoes, apply 12 to 25 lbs per acre.

Item No.

*Dipel is the registered trademark of Abbott Laboratories, North Chicago, Illinois 60064.

GOLD BUST, INC.
Atlanta, Georgia 30301