

\LABEL\22PREMRG.LAB HD #16
EPA File Symbol: 2269-__

105
CMA S
2269-174

Page 1 of 4
25 LB BAG TREATS
UP TO 5,000 SQ.FT.
OF LAWN



PRE-WEED AND FEED
FOR SOUTHERN LAWNS

**GOLD KIST
FERTILIZER + CRABGRASS PREVENTER**

Contains SURFLAN®, a Pre-emergent Herbicide for the Control of Crabgrass, Henbit, and Most Other Annual Broadleaf & Grassy Weeds in Lawns, Golf Courses, Parks, and other Ornamental and Recreational Turf areas.

For Established Southern Lawns

Bermudagrass - Zoysiagrass - St. Augustinegrass - Bahiagrass - Centipedegrass

**CONTAINS
20-0-10 PLANT FOOD**

PLUS>>>High Potash Ratio ...For Winter & Summer Survival

MAY BE APPLIED IN THE FALL OR SPRING
FOR PRE-EMERGENT CONTROL OF ANNUAL GRASSES AND BROADLEAF WEEDS

ACTIVE INGREDIENT:

ORYZALIN: 3,5-dinitro-N4,N4-

dipropylsulfanilamide 0.70%

INERT INGREDIENTS: 99.30%

Total 100.00%

SURFLAN® is a Registered Trade Mark of the Elanco Products Company

KEEP OUT OF REACH OF CHILDREN

W A R N I N G

STATEMENTS OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg white, gelatin, or, if these are not available, large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: If breathing difficulty occurs, move individual to fresh air and get medical attention.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 2269-__

EPA EST. NO. 2269-GA-1

NET WEIGHT 25 POUNDS (50 POUNDS)

**GOLD KIST INC.
ATLANTA GA 30301**

EST 1901
Use of Fertilizer and Insecticides,
Fungicides and Rodenticides Act,
as amended, for the pesticides
registered under
EPA Reg. No. 2269-174

FERTILIZER + CRABGRASS PREVENTER

FALL Application - 25 Pounds covers 5,000 Sq.Ft.
SPRING Application - 25 pounds covers 4,000 Sq.Ft. up to 5,000 Sq.Ft.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WHERE TO USE

When applied according to Directions this product can be used safely on the following established Southern Lawn, warm season turfgrass varieties: BAHIAGRASS, BERMUDAGRASS, CENTIPEDEGRASS, ST. AUGUSTINEGRASS VARIETIES, TALL FESCUE VARIETIES, & ZOYSIAGRASS lawns and ornamental turf. It may be used for Home Lawns, Apartment and Condominium complexes, Park, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turfgrass areas. Will not harm most nearby ornamental trees and shrubs when used as directed.

GOLD KIST® FERTILIZER + CRABGRASS PREVENTER

A BROAD SPECTRUM PRE-EMERGENT HERBICIDE + FERTILIZER FOR SOUTHERN LAWNS

**CONTROLS
GRASSY WEEDS & BROADLEAF WEEDS**

This New product, backed by ELANCO research, gives excellent pre-emergent control of GRASSY AND BROADLEAF WEEDS as listed. It contains Oryzalin Herbicide (commonly known as SURFLAN®, a registered Trade Mark of Elanco Products Co.) in a free flowing combination herbicide-fertilizer product.

GRASSY WEEDS

It should be applied to established turf for the control of ANNUAL BLUEGRASS (Poa Annuu), BARNYARD GRASS, BROWNTOP PANICUM, CRABGRASS (Large Crabgrass, Smooth Crabgrass), CROWFOOTGRASS, CUPGRASS, FOXTAILS (Bottlegrass, Bristlegrass, Giant Foxtail, Green Foxtail, Pigeongrass, Robust Foxtail, Yellow Foxtail), GOOSEGRASS (Silvercrabgrass), JOHNSON GRASS SEEDLINGS, LOVEGRASS (Mexican, Orcutt), FALL PANICUM, RYEGRASS (Annual), SANDEUR, TEXAS PANICUM (Buffalograss, Coloradograss), and WITCHGRASS.

BROADLEAF WEEDS

Will also control many Broadleaf Weeds: BITTERCRESS, CARPETWEED, COMMON CHICKWEED, FIDDLENECK (Coast), FILAREE (Redstem, Whitestem), GROUNDSEL, HENBIT, KNOTWEED, LAMBSQUARTER, PIGWEEDS, PUNCTURE VINE, COMMON PURSLANE, FLORIDA PUSLEY, LONDON ROCKET, ROCK PURSLANE, SHEPARDSPURSE, SPRANGLETOP, PROSTRATE SPURGE, WOODSORREL (Creeping).

THE FOLLOWING WEEDS MAY BE SUPPRESSED BY APPLICATIONS

Black Nightshade, Common Ragweed, Ladysthumb, Morningglory, Smartweed, Teaweed (Prickly Sida), Velvetleaf, Horseweed, Prickly Lettuce, Common Mallow, Black Mustard, Wild Mustard, Annual Sowthistle, Climbing Milkweed, Spotted Spurge, and Volunteer Wheat.

IMPORTANT PRECAUTIONS.

DO NOT apply in the spring to Bermuda Grass which has been overseeded with winter grasses (annual rye) since thinning will occur. DO NOT do not use on Dichondra, nor to newly sprigged areas of Bermudagrass, St. Augustine Grass or Centipede Grass, until these turfs are well established, nor to Bentgrass or Putting Greens. DO NOT apply in spring to new Turfgrass planted the previous fall. DO NOT use this product at rates higher than recommended, e.i. 25 lbs./4,000 Sq.Ft. DO NOT OVERDOSE OR OVERLAP.

WHEN TO APPLY

FOR BEST RESULTS APPLY AT THE PROPER TIME

Apply only to healthy established turf at least 1 to 2 weeks prior to germination of weed seeds since established weeds will not be controlled. All summer annual grasses do not germinate at the same time. Many weeds germinate in the fall. For this reason your application must coincide with the normal germination period. If applied too early or too late, erratic weed control will result. See Chart for proper timing.

WATER THE LAWN AFTER APPLICATION: Best results will be obtained if treated areas are irrigated with 1/4 to 1/2 inch water soon after application. Any cultural practices that disturb the soil such as verticutting, should be done prior to treatment.

WINTER ANNUAL WEEDS

**FOR CONTROL OF WINTER ANNUAL BROADLEAF AND GRASSY WEEDS
IN SOUTHERN LAWNS**

For pre-emergence weed control of listed "winter annual" grassy and broadleaf weeds apply 25 lbs. per 5,000 Sq.Ft. of Lawn; (this is equal to 1.54 lbs herbicide per acre), Apply in the fall (as early as Mid-September) or early winter at least one to two weeks prior to warm moist weather which is ideal for the germination of weed seeds. See Chart; (No.4)

"Sun-Turf" Bermudagrass has shown sensitivity to this herbicide e.i. first test apply on a small inconspicuous plot before treating the entire area.

**SPRING ANNUAL WEEDS
IN SOUTHERN LAWNS**

For control of weeds as listed in late winter or spring apply 25 lbs. to cover 4,000 to 5,000 Sq.Ft. (equivalent to 1.54 lbs. to 1.92 lbs herbicide per acre). Apply 1 to 2 weeks prior to the onset of conditions favorable for weed seed germination. See Chart; (No.1)

When making a "first ever application" or where the turf has been completely overrun by weeds use 25 lbs. to cover 4,000 Sq.Ft. When weeds have been sparse and/or the turf has been or is now in a Yearly Weed Control Program 25 lbs. may be used to cover 5,000 Sq.Ft.

\LABEL\22PREMRG.LA5 HD #16
EPA File Symbol: 2269--

OPTIONAL LABEL COPY OR FOR BROCHURE
FERTILIZER + CRABGRASS PREVENTER

WEED CONTROL IS BASED ON PROPER APPLICATION

The following spreader settings are for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreader may require a slight adjustment.

If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 500 Sq.Ft. into your spreader and apply to your pre-measured 500 Sq.Ft. (5 ft. wide by 100 ft long)

HOW TO MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

Using a Scott Spreader for Weed Control place 2.50 lbs. granules in your spreader using No. 5 1/2 setting & apply to 500 Sq.Ft. With a 20 inch spreader this will equal 3 spreader widths wide (3 passes) by 100 feet long. If you run out before you cover the pre-measure area close the spreader setting by 1/2 the number. If you have granules left over, open (increase) the spreader setting by 1/2 the number. For further help refer to your Spreader Guide Book.

POOR WEED CONTROL WILL RESULT IF YOUR SPREADER APPLIES LESS GRANULES THAN RECOMMENDED.... MAKE SURE TO APPLY THE TOTAL POUNDS PER SQUARE FEET AS SHOWN IN THE FOLLOWING CHART.

SPREADER SETTINGS

		4,000 SQ. FT.	5,000 SQ. FT.
		6 1/2 lbs./1,000 Sq.Ft.	5 lbs./1,000 Sq.Ft.
CYCLONE B-1	25 lbs.	4 1/2	3 1/2
SCOTTS #35	25 lbs.	6 1/2	5 1/2

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

When using a 50 lb. bag; spreader settings will be the same as for the 25 lb. bag but it will treat only have the amount of Sq.Ft. as the 50 lb bag. These settings are based on a "One Time Over" application.

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.