

OTHER EXCELCIDE PRODUCTS

For Rodent Control

Bait Boxes

For dispensing of dry or liquid bait

In cardboard, wood, plastic or metal

Bait

Besides Pival — also dry Warfarin, PMP in cereal and liquid form, and Diphacin.

Traps

Ketch-All Mouse Traps

Metal Extended Trigger Mouse Trap

New Type Wood Snap Mouse Trap

DIRECTIONS
FOR RATS: EXCELCIDE Pival, used in food processing areas, non-food processing areas, or any rooms adjacent, should be dispersed in EXCELCIDE Fibreboard, Wood, or Metal Bait Boxes. Four ounces per station should be the minimum amount used. An uninterrupted supply of bait should be maintained for at least 10 days. Check all baits daily and keep containers full of bait. Continue to feed the rats until kill and control occurs. This usually takes from 5-14 days, but may take longer.

CONTINUOUS BAITING IS ESSENTIAL FOR RESULTS. Protect the bait from children and domestic animals by means of bait boxes where necessary. Where possible, choose clean, dry places for placement to avoid bait souring or decomposing. Place baits in locations not accessible to children, pets, and domestic animals.

FOR MICE: Follow the same directions as for rats except that bait placements may be smaller in size. Bait, ¼ to ½ oz., should be placed at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days and placements should be made in more places over infested areas. Baits should be placed not over 10-20 feet apart in infested areas for best results.

ACTIVE INGREDIENT

* 2-Pivalyl-1,3-indandione 0.025%

INERT INGREDIENTS

Food Mixture 99.975%

* Pival

USDA Registration No. 2270-12



**"EXCELCIDE"
PIVAL**

RAT and MOUSE KILLER

**READY-TO-USE
CAUTION:**

**Keep out of reach of children.
(See right panel for additional cautions)**



Net Weight 6 lbs.

THE HUGE COMPANY, INC.
Sanitation Consultants to the Food Industry
MANUFACTURERS OF

ACCEPTED

FEB 10 1960

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE ACT OF 1947 AS AMENDED
 FOR ECONOMIC PEST CONTROL REGISTERED
 UNDER NO. 2270-12

CAUTION

This product contains as its active ingredient an anticoagulant agent (Pival) an anti-Vitamin K agent. If swallowed by humans, domestic animals or birds, this material may reduce the clotting of the blood and cause bleeding. In severe case, intravenous and oral administration of Vitamin K, combined with transfusions, are indicated, and in case of hemorrhage caused by this product, the use of bishydroxycoumarin.

NOTICE

The Huge Company, Inc. warrants that this product conforms to its description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but no warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, is made by the use of this product contrary to the instructions, or under abnormal conditions not reasonably foreseeable, and buyer assumes the risk of such use.

**OTHER EXCELCIDE
PRODUCTS**

For Rodent Control

Bait Boxes

For dispensing of dry or liquid bait

In cardboard, wood, plastic or metal

Bait

Besides Pival — also dry Warfarin, PMP in cereal and liquid form, and Diphacin.

Traps

Ketch-All Mouse Traps

Metal Extended Trigger Mouse Trap

New Type Wood Snap Mouse Trap

DIRECTIONS
FOR RATS: EXCELCIDE Pival, used in food processing areas, non-food processing areas, or any rooms adjacent, should be dispersed in EXCELCIDE Fibreboard, Wood, or Metal Bait Boxes. Four ounces per station should be the minimum amount used. An uninterrupted supply of bait should be maintained for at least 10 days. Check all baits daily and keep containers full of bait. Continue to feed the rats until kill and control occurs. This usually takes from 5-14 days, but may take longer.

CONTINUOUS BAITING IS ESSENTIAL FOR RESULTS. Protect the bait from children and domestic animals by means of bait boxes where necessary. Where possible, choose clean, dry places for placement to avoid bait souring or decomposing. Place baits in locations not accessible to children, pets, and domestic animals.

FOR MICE: Follow the same directions as for rats except that bait placements may be smaller in size. Bait, ¼ to ½ oz., should be placed at 8-12 foot intervals. An uninterrupted supply of bait should be maintained for at least 15 days and placements should be made in more places over infested areas. Baits should be placed not over 10-20 feet apart in infested areas for best results.

ACTIVE INGREDIENT

* 2-Pivalyl-1,3-indandione 0.025%

INERT INGREDIENTS

Food Mixture 99.975%

* Pival

USDA Registration No. 2270-12



EXCELCIDE

PIVAL

RAT and MOUSE KILLER

CAUTION:

Keep out of reach of children.

(See right panel for additional cautions)



Net Weight 6 lbs.

THE HUGE COMPANY, INC.

Sanitation Consultants to the Food Industry

MANUFACTURERS OF

884-886 HODIAMONT AVE. • ST. LOUIS, MISSOURI 63112

ACCEPTED

FEB 10 1960

2270-12

CAUTION

This product contains as its active ingredient an anticoagulant agent (Pival) an anti-Vitamin K agent. If swallowed by humans, domestic animals or birds, this material may reduce the clotting ability of the blood and cause bleeding. In severe cases, intravenous and oral administration of Vitamin K, combined with blood transfusions, are indicated, in case of hemorrhage caused by the use of bishydroxycoumarin.

NOTICE

The Huge Company, Inc. warrants that this product conforms to its description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but no warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, is made by the use of this product contrary to the instructions, or under abnormal conditions not reasonably foreseeable. Buyer assumes the responsibility for such use.

DIRECTIONS
 EXCELCIDE Pival, used in baiting areas, non-food processing areas, or any rooms adjacent to food, should be dispersed in EXCELCIDE Bait Boxes, Metal Bait Boxes, or Metal Bait Stations. Baiting station should be the only one used. An uninterrupted supply of bait should be maintained for 15 days. Check all baits daily and replenish when full of bait. Continue until rats are killed and control is usually taken from 5-7 days longer.

BAITING IS ESSENTIAL
 1. Protect the bait from domestic animals by means of bait boxes where necessary. Where possible, use clean, dry places for baiting to avoid bait souring or degrading. Place baits in locations inaccessible to children, pets, and livestock.

Follow the same directions as above except that bait placements should be made at 8-12 foot intervals. A continuous supply of bait should be maintained for at least 15 days and could be made in more restricted areas. Baits should be replaced every 10-20 feet apart in order for best results.

ACTIVE INGREDIENT
 Pival 0.025%

OTHER INGREDIENTS
 99.975%

Pat. No. 2270-12



"EXCELCIDE"

PIVAL

RAT and MOUSE KILLER

**READY-TO-USE
 CAUTION:**

Keep out of reach of children.

(See right panel for additional cautions)



Net Weight 6 lbs.

THE HUGE COMPANY, INC.

Sanitation Consultants to the Food Industry

MANUFACTURERS OF

884-886 HODIAMONT AVE. • ST. LOUIS, MISSOURI 63112

ACCEPTED

FEB 10 1950

UNDER THE
 FEDERAL
 FOOD, DRUG,
 AND COSMETIC ACT
 2270-12

CAUTION

This product contains as its active chemical an anticoagulant agent (Pival), an anti-Vitamin K agent. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K, combined with blood transfusions, are indicated, as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

NOTICE

The Huge Company, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

OTHER EXCELCIDE PRODUCTS

For Insect Control

EXCELCIDE Residual Spray
 EXCELCIDE Fly Spray
 EXCELCIDE Vapo Spray
 EXCELCIDE Mill Spray
 EXCELCIDE Resifume
 EXCELCIDE Vapo Fly Spray
 (with Vapona)

EXCELCIDE Larvae Concentrate

EXCELCIDE Outside Residual

EXCELCIDE Fly Tid-Bits

EXCELCIDE Formulas for use in Dyna-Fog machines

EXCELCIDE Residual No. 2A

EXCELCIDE Residual No. 2B

EXCELCIDE Stro-Mal Residual No. 1

EXCELCIDE Stro-Mal Residual No. 2

Also space, residual and hand sprayers

Overhead spraying equipment