

Rsp. 77 227

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Strictly follow all safety precautions and directions for use on this labeling.

### STORAGE AND DISPOSAL

**STORAGE:** Store in a secure area separated from water, food or feed. Keep container tightly closed when not in use. Do not transfer this product to unlabeled storage containers. Store away from heat, strong oxidizers or open flame.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinsate that is not used according to label instructions must be disposed of according to approved federal, state and local procedures.

**CONTAINER DISPOSAL:** Metal Containers—Triple rinse (or equivalent). Then offer for reconditioning or recycling, or puncture and dispose of in a sanitary landfill, or by other approved federal, state and local procedures. Plastic Containers—Triple rinse (or equivalent). Then offer for reconditioning or recycling, or dispose of in a sanitary landfill, or by professional incineration if allowed by federal, state and local authorities.

**FOR CATTLE**—To protect dairy stock from biting flies, houseflies, mosquitoes and gnats, as well as certain other flying pests, spray daily using 1-2 ounces per head. Cover the animal with spray with particular attention to back, flanks and legs. Since continued daily treatments create a buildup of the repellent on the animal, spraying may be reduced to once every other day after the first three or four days.

**FOR HORSES**—Spray lightly or spray horses, using 1-2 ounces per adult animal, proportionately less for smaller animals. Give special attention to shanks, necks and shoulders. Apply until hair is moistened but do not saturate.

**FOR PIGS**—Spray 1-2 ounces where flies are present. Repeat as needed. Do not apply more than 2 ounces per animal per application to cattle, horses and pigs. Apply to animals as a mist spray to the outer hair.

EXCELCIDE Dairy Cattle, Livestock & Barn Spra contains a combination of ingredients which have been selected for use on warm blooded animals, milk and meat, when used as directed, for killing and repelling horseflies, stable flies, houseflies, gnats and mosquitoes, and provides protection against these insects during and after spraying. Also kills horseflies that are hit by the spray.


EXCELCIDE Dairy Cattle Livestock & Barn Spra is a special formulation containing a repellent which provides protection from biting flies in the pasture between milkings. This reduces the discomfort and annoyance caused by flies which often decreases milk production and weight gain.

### CAUTION

Do not use or store near heat or open flame. Avoid wetting the hides of animals. Harmful if swallowed.

SAMPLE LABEL

NET CONTENTS GALS.



# DAIRY CATTLE, LIVESTOCK & BARN SPRA

For the protection of Dairy Animals From Certain Biting Flies. Also Kills Flies, Mosquitoes and Gnats in Barns and Stables. For Use On Dairy Animals.

For Practical Pasture Protection.

ACTIVE INGREDIENTS	100%
Petroleum Distillate	63.215%
Mineral Seal Oil	36.000%
Di-n-propyl isocincomeronate*	.500%
N-octyl bicycloheptene dicarboximide	.150%
Technical Piperonyl Butoxide**	.090%
Pyrethrins	.045%

\*MCK Repellent 326  
\*\*Consists of .072% (butylcarbityl) (6-propylpiperonyl) ether and .018% related compounds.



EPA EST. NO. 2270-MO-01  
EPA Registration No 2270 26

**CAUTION:**  
Keep out of reach of children.

**THE HUGE COMPANY, INC.**  
*Sanitation in the Food and Allied Industries*  
Manufacturers of EXCELCIDE INSECTICIDES and PEST CONTROL TOOLS  
7625 PAGE AVE. • ST. LOUIS, MISSOURI 63133

### FOR BARN, STABLES AND MILK ROOMS

When used as a space spray direct power sprayer toward upper portion of the enclosure until the area is filled with a fine mist. Do not apply this product in poultry houses.

Doors and windows need not be closed during or after application. When coarse spray is used, apply directly on walls, stanchions, ceilings, window and door frames wherever flies alight or congregate until surfaces are moistened.

Avoid direct contact with food or food utensils. Leave doors and windows open during application to permit flies to leave. After fly population is reduced, doors and windows may be closed. Spray outdoor surfaces also to discourage flies from resting and repeat as needed.



### NOTICE

Seller and manufacturer warrant that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

See side panel for additional cautions.