


IMPORTANT NOTICE

GENERAL DIRECTIONS: Pour contents into a dry container next to burrows, in runways, or in locations where rats and mice are accustomed to feeding, drinking or travelling. Desirable bait placements are along walls, in corners, and in concealed locations where rats and mice can feed unobserved. Replace stale, damp or moldy bait with fresh bait as necessary. Continue baiting until all signs of feeding have stopped.

This rodenticide contains as its active ingredient an anticoagulant chemical that reduces the clotting ability of the blood. Upon repeated feedings, it causes fatal hemorrhage in rats and mice. Bait should be placed in locations not accessible to children, pets or domestic animals, or in tamper proof bait boxes.

RATS: Use 4 or more ounces of bait at each location and maintain an uninterrupted supply of bait for at least 10 days.

MICE: Place tablespoon amounts (1/4 to 1/2 ounce) of bait in shallow containers at 8-10 foot intervals and maintain an uninterrupted supply of bait for at least 15 days.



PIVAL PELLETS

KILLS RATS AND MICE


ACTIVE INGREDIENT:
2-Pivalyl-1, 3-indandione* 0.025%
INERT INGREDIENT: 99.975%
*Also known as Pivalyl

CAUTION:
Keep out of reach of children.
(See additional cautions on side panels)



Net Wt. Lbs.

Manufactured for
THE HUGE COMPANY, INC.
Sanitation Consultants to the Food Industry
MANUFACTURERS OF
EXCELCIDE INSECTICIDES, RODENTICIDES AND EQUIPMENT
884-886 HODIAMONT AVE. • ST. LOUIS, MISSOURI 63112



CAUTION: Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes. If swallowed by humans, animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, a physician or veterinarian should be called immediately. Intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

EPA Reg. No. 2270-161


IMPORTANT NOTICE

GENERAL DIRECTIONS: Pour contents into a dry container next to burrows, in runways, or in locations where rats and mice are accustomed to feeding, drinking or travelling. Desirable bait placements are along walls, in corners, and in concealed locations where rats and mice can feed unobserved. Replace stale, damp or moldy bait with fresh bait as necessary. Continue baiting until all signs of feeding have stopped.

This rodenticide contains as its active ingredient an anticoagulant chemical that reduces the clotting ability of the blood. Upon repeated feedings, it causes fatal hemorrhage in rats and mice. Bait should be placed in locations not accessible to children, pets or domestic animals, or in tamper proof bait boxes.

RATS: Use 4 or more ounces of bait at each location and maintain an uninterrupted supply of bait for at least 10 days

MICE: Place tablespoon amounts (1/4 to 1/2 ounce) of bait in shallow containers at 8-10 foot intervals and maintain an uninterrupted supply of bait for at least 15 days.



PIVAL PELLETS

KILLS RATS AND MICE


ACTIVE INGREDIENT:
2-Pivalyl-1, 3-indandione* 0.025%
INERT INGREDIENT: 99.975%
*Also known as Pivalyl

CAUTION:
Keep out of reach of children.
(See additional cautions on side panels)



Net Wt. Lbs.

Manufactured for
THE HUGE COMPANY, INC.
Sanitation Consultants to the Food Industry
MANUFACTURERS OF
EXCELCIDE INSECTICIDES, RODENTICIDES AND EQUIPMENT
884-886 HODIAMONT AVE. • ST. LOUIS, MISSOURI 63112



CAUTION: Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes. If swallowed by humans, animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, a physician or veterinarian should be called immediately. Intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

Treated baits should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamper-proof bait boxes. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

EPA Reg. No. 2270-161