

[220-242]

EXCELICIDE FOGICIDE #131

ACCEPTED
NOV 20, 1970

READY-TO-USE

UNDER THE FEDERAL
INSECTICIDE AND
RODENTICIDE ACT
AND ECONOMIC
INSECTICIDE
REGISTRATION NO.

DIRECTIONS

For control of flying insects either inside or outside--Drosophila Gnats (Vinegar and Fruit Flies), Houseflies, Blowflies, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths; and for such crawling insects as Cockroaches, Confused Flour Beetles, Ants, Caddisflies, Silverfish, Carpet Beetles that would be exposed at the time of application.

RECOMMENDATIONS

Most effective inside fogging of food processing plants and inside of structures normally found on parks, golf courses, and outdoor theatres, is accomplished during dry weather. External openings to buildings should be closed with exception of probably one providing an up-draft. All parts of building or enclosure should receive equal concentration of fog. Exposed foods should be guarded by proper protective covering or by temporary relocation. Plan fogging operation so that operator may continually move away from fog. Fogging should be done from windward side when applying fog on outside of building.

For use in thermal-type generators such as the Dyna-Fog machine for control of certain crawling and flying insects.

THIS FORMULA MUST BE USED BY TRAINED OPERATORS ONLY

ACTIVE INGREDIENTS	100%	
Petroleum Distillate		99.625%
Technical Piperonyl Butoxide*		.250%
Pyrethrins		.125%

CAUTION: KEEP OUT OF REACH OF CHILDREN!
(See additional cautions below)

*Equivalent to .200% (butylcarbityl) (6-propyl-piperonyl) ether and .050% related compounds.

DIRECTIONS

For outdoor fogging the area to be treated (parks, golf courses, outdoor theatres, and the immediate adjacent areas of structures normally found in the above areas and around food processing plant buildings) should be completely covered with visible fog, or at the rate of about 2 1/2 pints per acre. The same is true of indoor treatment, although in the latter case no more than 1 gallon per 50,000 cubic feet should be applied, and the particle size of the insecticide should be 1/2 to 10 microns. Machine is adjustable for these sizes. Repeat application should be effected as often as is necessary.

Spray EXCELICIDE Fogicide #131 upward at walls and ceilings so that as many insects as possible can be contacted with direct mist. After spraying, sweep up all dead and dying insects, thereby reducing infestation.

CAUTION: Harmful if swallowed. Do not contaminate or store near feed or foodstuff. Use in food processing plants only when plant is not in operation. All food processing surfaces should be thoroughly cleaned before using. Do not reuse container. Destroy when empty. This product is not intended for household use.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE: The manufacturer and seller warrant that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Registration No.2270-242

THE HUGE COMPANY, INC.
Sanitation Consultants to the Food Industry
MANUFACTURERS OF
EXCELICIDE INSECTICIDES, RODENTICIDES AND ROACHING
884-886 HODIAMONT AVE. • ST. LOUIS, MISSOURI 63112