

FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
NO. UNDER REG. 2-12-73

2272-13

ACTIVE INGREDIENT — Sodium Hypochlorite, 12.5%
INERT INGREDIENTS — 87.5%

Note on composition: Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine.

KEEP IN A COOL DARK PLACE TO SLOW DECOMPOSITION; VENT DAILY TO RELIEVE PRESSURE INSIDE.

CONTAINER DISPOSAL; RINSE EMPTY CONTAINER THOROUGHLY WITH WATER AND DISCARD — OR RETURN FOR DEPOSIT IF A DEPOSIT CONTAINER, WITH ALL PLUGS INSERTED AND TIGHTENED.

DIRECTIONS FOR USE:

As a germicidal rinse, for cleaned non-porous surfaces such as metal or glass, dilute 1 oz. to 6 oz. of water.

For disinfection on porous surfaces, such as concrete, tile, or wood, after cleaning dilute 1 oz. to 1 gal. of water and apply so as to thoroughly cover area to be cleaned by wet mopping, scrubbing, or flushing.

FOR USE AS A SOURCE OF SODIUM HYPOCHLORITE IN CLEANING EQUIPMENT, FLOORS, & WALLS.

WARNING

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED — STRONG OXIDANT — MAY PRODUCE CHEMICAL BURNS. DO NOT GET ON SKIN OR IN EYES OR ON CLOTHING.

In case of contact with skin or eyes, flush with plenty of water. If skin irritation persists, get medical attention. For eyes get prompt medical attention.

FIRST AID: If taken internally, drink raw egg white, milk, or rice gruel. Follow with a tablespoon of mustard in warm water. Call a physician.

