

DIXIE MALATHION DUST

CAUTION: KEEP OUT OF REACH OF CHILDREN

AVOID BREATHING OF DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL. DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. - STAY AWAY FROM SMOKE OR FUMES.

DIRECTIONS

USDA Reg. No. 2301-9

To control the insects listed below dust at the rate of 25 to 35 pounds per acre. Repeat the applications as necessary. Usually two or more applications at 5 to 7 day intervals will be necessary. To control Mexican Bean Beetle direct the dust to the underside of the leaves.

VEGETABLES

BEANS: Aphids, Leafhoppers, Mexican Bean Beetle, Spider Mites, apply 25 to 35 pounds per acre. No application should be made to beans within 3 days of harvest, providing the dosage does not exceed 35 pounds of the product per acre.

CABBAGE, BROCCOLI, TURNIP: Aphids, Imported Cabbage Worm and Cabbage Looper.

CUCUMBERS, SQUASH AND MELONS: Aphids and Spider Mites. Do not apply to plants unless they are dry or injury may result.

EGGPLANT: Spider Mites and Eggplant Lace Bug.

ONION: Thrips.

PEAS: Aphids and Pea Weevil

PEPPERS: Aphids.

POTATO and TOMATO: Aphids, Spider Mites and Leafhoppers.

Do not use on Beans, Cucumbers, Melons or Squash within 1 day of harvest; on Broccoli, Eggplant, Onions, Peas, Peppers, Tomatoes, Turnips or Potatoes within 3 days of harvest; on Brussel Sprouts, Cabbage, Cauliflower, within 7 days of harvest.

POULTRY

CHICKENS; DUCKS; GEESE AND TURKEYS: To control Northern Fowl Mite, Lice and as a supplement to premise treatment for Chicken Red Mite. Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary.

PREMISE TREATMENTS: To control Northern Fowl Mite, Chicken Red Mite and Poultry Lice apply at the rate of 1 lb. to 50-60 sq. ft. of litter and floor space and to nests, roosts and adjacent areas. Repeat application as necessary. **AVOID CONTAMINATION OF FEED AND AND FEEDING TROUGHS AND WATER FOUNTAINS.**

DOGS AND CATS

FLEAS: Apply just enough dust to treat animal thoroughly. Apply to the back and work into the coat with the hand. Wash hands thoroughly after using. Repeat application every second or third week.

ACTIVE INGREDIENTS: "Malathion: O,O-dimethyl dithiophosphate of diethylmercaptosuccinate ~5%

INERT INGREDIENTS: 95%

TOTAL 100%

NOTICE

C.D. TERHUNE CO. makes no warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risks of handling or use whether in accordance with directions or not.

Net Weight 5 Lbs.

Mfg. By: C. D. TERHUNE CO. George West Rd., Cedartown, Ga.

ACCEPTED
July 2, 1971
U.S. DEPARTMENT OF AGRICULTURE
FEDERAL INSECTICIDE

ACCEPTED WITH COMMENTS