

DIXIE 5% SEVIN DUST

CAUTION: KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN! AVOID BREATHING DUST OR SPRAY MIST. AVOID CONTACT WITH EYES OR SKIN. WASH THOROUGHLY AFTER HANDLING. WASH ALL CONTAMINATED CLOTHING WITH SOAP AND WATER BEFORE RE-USE. AVOID STORAGE NEAR FEED AND FOODSTUFFS. THIS PRODUCT IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT TREATMENT OR RESIDUE ON CROPS. PROTECTIVE INFORMATION MAY BE OBTAINED FROM YOUR COOPERATIVE AGRICULTURAL EXTENSION SERVICE. - DO NOT REUSE EMPTY CONTAINER. - DESTROY IT BY BURYING WITH WASTE OR BURNING. - STAY AWAY FROM SMOKE OR FUMES.

DIRECTIONS

BEANS: Apply thoroughly to upper and lower leaf surfaces at first sign of infestation before feeding occurs: **FOR CONTROL OF MEXICAN BEAN BEETLE** apply 10 lbs. per acre: **FOR LEAFHOPPERS** apply 20 lbs. per acre. **FOR LYGUS AND STINK BUGS** apply 40 lbs. per acre. If needed, repeat applications every 7 to 10 days up to and including day of harvest.

CUCUMBER, SUMMER SQUASH: **FOR CONTROL OF SQUASH BUGS, CUCUMBER BEETLES, FLEA BEETLES, PICKLE WORM AND MELONWORM** apply 20 lbs. per acre. Apply when insects or their damage appear. Repeat at 7-10 day intervals or as necessary. May be applied up to and including day of harvest.

TOMATO, EGGPLANT, PEPPER: **FOR CONTROL OF TOMATO FRUITWORM, FALL ARMYWORM, FLEA BEETLES:** Apply at the rate of 20-40 lbs. per acre: **FOR CONTROL OF TOMATO HORNWORM, FALL ARMYWORM AND STINK BUGS** apply at the rate of 40 lbs. per acre. Apply when insects or their damage appear. Repeat at 7-10 day intervals or as necessary, up to and including day of harvest.

POTATOES: Apply on appearance of insect. **FOR CONTROL OF COLORADO POTATO BEETLE** apply 10 to 15 lbs. per acre: **FOR LEAFHOPPERS AND FLEA BEETLES** apply 20 to 25 lbs. per acre: **FOR LYGUS BUGS** apply 30 to 40 lbs. per acre. Repeat if needed every 7-10 days up to and including day of harvest.

CHICKENS, DUCKS, GAMEBIRDS, GEESE, PARTRIDGES, PHEASANTS, PIGEONS, TURKEYS. For Northern Fowl Mite, Lice and Chicken Red Mite use 1 lb. per 100 birds. Do not apply within 7 days of slaughter. Do not repeat application within four weeks. Do not contaminate feed or drinking water. For housing the above use 25 lbs. per 1000 square feet. Apply thoroughly to floors, roosts and interior surfaces. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter.

DOGS AND CATS: **FOR CONTROL OF FLEAS AND BROWN DOG TICKS** apply liberally over the animal and rub thoroughly into skin. Apply also to the feet, between toes and on the legs. Dust in and around sleeping quarters and other areas where ticks and fleas occur. Apply at weekly intervals if needed. **AVOID CONTACT WITH EYES AND MUCOUS MEMBRANES.**

ACTIVE INGREDIENTS

Carbaryl (1-Naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS 95%

Total 100%

NOTICE Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to labeling instructions.

EPA REG. NO. 101-111

NET WT. 10 LBS. EPA EST. REG. NO. 101-111

MANUFACTURED BY C. I. TERHUNE CO., George West, Rte. 1, California