

Directions for Use

For sanitizing and disinfecting floors, walls and lacinate hard surfaces, apply solution with mop or cloth. Sanitize with 1 ounce to 4 gallons of water. Disinfect with 3 ounces to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

For the Five Step Sanitization Procedure necessary for use in the State of Wisconsin, see attached brochure.

Hospitals
For floors, walls and other hard surfaces in Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use 3 1/2 ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Food Processing Equipment
For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce per 4 gallons of water to provide 200 ppm of active quaternaries. At this level No Potable Water Rinse is required.

Dairies
To sanitize dairy equipment such as tanks, lines, pails and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce to 4 gallons of water (200 ppm). At this level, No Potable Water Rinse is required. Follow recommendations of local Health Board.

Restaurant and Bar Rinse
Dishes, glassware, silverware, cooking utensils wash with soap or synthetic detergent, rinse thoroughly and immerse in a sanitizing solution containing 1 ounce to 4 gallons of water. No Terminal Potable Water Rinse Required.

This product when used on environmental inanimate, hard surfaces at 3 1/2 ounces per 5 gallons of water is effective against influenza A₂, Herpes Simplex, Adenovirus type 1 and Vaccinia Virus.

Net Contents 1 U.S. Gallons

QAT 1000

DISINFECTANT DEODORIZER SANITIZER

ACTIVE INGREDIENTS:

- n Alkyl (60% C₁₂, 30% C₁₄, 5% C₁₆, 5% C₁₈) dimethyl benzyl ammonium chlorides
- n Alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium chlorides

INERT INGREDIENTS:

QAT 1000 is a concentrate and should be diluted before using

DANGER

Keep Out of Reach of Children

See other precautions and first aid on side panel

EPA REG. NO. 2311-11 -

EPA EST NO. 023111L01

ONLY FOR SALE TO, USE, AND STORAGE BY SERVICE PERSONS

SOLED BY

HAAG LABORATORIES INC.

14000 S. SEELEY AVE.

BLUE ISLAND, IL 60406



ACCEPTED
 MAR 20 1964
 U.S. Environmental Protection Agency
 registered under
 EPA Reg. No. 2311-11

BEST DOCUMENT AVAILABLE

Cold Disinfection
For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 ounces per gallon of water for 10 minutes.

FOR ROOM FOGGING

DIRECTIONS:

To minimize the danger of cross-contamination from environmental surfaces apply solution as a fog as described below, before applying standard routine terminal cleaning and disinfecting practices. 1. Remove all human, animal, and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount Jet FOG-MASTER TRI-JET MODEL 6200 or other fogging device delivering equivalent spray at 3 RPM. 5. Fill sprayer reservoir with 15 oz. to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before entering room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN


Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed. Do not reuse empty container. Rinse thoroughly with water before discarding or returning to drum reconditioner.

Registration Division (TS 707)

STATE OF WISCONSIN FIVE STEP SANITIZATION PROCEDURE
FOR PRODUCTS IDENTICAL TO QAT 1000, EPA REG. NO. 2311-11

- 1) Scrap and pre-wash utensils and glasses whenever possible.
- 2) Wash with a good detergent or compatible cleaner (a commercial cleaner or detergent can be recommended as long as it is not of an anionic nature).
- 3) Rinse with clean water.
- 4) Sanitize in a solution of 1 ounce to 4 gallons of water (200ppm). Immerse all utensils for at least two minutes or contact time specified by governing sanitary code.
- 5) Place sanitized utensils on rack or drainboard to air dry.

Note: A clean potable water rinse following sanitization is not permitted under Section H 96.06 (3) (h) of the Wisconsin Administrative code.



Rec. No. 11-15107