



NET CONTENTS
5 POUNDS

NU-TRI-COP

PRODUCT
No. 3-0500-2

ACTIVE INGREDIENT:	By Wt.
Copper expressed as Metallic (Derived from Basic Copper Sulphate)	46.25%
INERT INGREDIENTS:	53.75%
	<hr/>
	Total — 100.00%

SPECIMEN LABEL
® Trade Mark Reg. U.S. Patent Office

**CAUTION: Keep Out of the Reach of Children.
See additional cautions on this label.**

FASCO Nu-Tri-Cop is a very fine particle size Neutral Copper in Tri-Basic Copper Sulphate form. Special wetting, spreading and adhesive agents have been added.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

FOR SMALL AMOUNTS — One level measuring tablespoonful per gallon is equal to 1½ pounds per 100 gallons of spray.

TOMATOES, PEPPER, POTATOES — Against Early and Late Blight of Potatoes and Tomatoes, Bacterial Spot of Tomatoes and Peppers, Septoria Leaf Spot of Tomatoes, and Late Frog-eye Spot of Peppers use 5 pounds per 100 gallons of spray. Begin spraying when diseases appear. Repeat every 3 to 7 days as needed.

CUCURBITS (Cucumbers, Watermelons, Cantaloupes, Squash) — Against Downy Mildew use 5 pounds per 100 gallons of spray every 5 to 10 days as needed. Begin spraying at first sign of disease.

CELERY — Against Early Blight use 5 pounds per 100 gallons of spray every 5 to 10 days as needed. Begin spraying at first sign of disease.

CITRUS — Use 1½ pounds per 100 gallons spray solution in Petal Fall or Post Bloom application against Scab and/or Melanose. For dormant application against Scab use 3 lbs. per each 100 gallons spray. FASCO Nu-Tri-Cop contains deflocculator and all necessary spreader sticker. It is compatible with all sprays used on Citrus. The use of Copper and oil combinations should be avoided after the fruit has attained a size of ¾ inch in diameter and when the temperature is excessively high.

CAUTION

Harmful if swallowed! If taken internally, induce vomiting by giving mustard and water and tickling the throat. Give large doses of magnesia or lime water, followed with doses of Epsom Salts. Avoid inhalation of dust.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

DRAFT H 71-6637 USDA Reg. No. 2342-9

ACCEPTED
6/15/71


Manufactured by FLORIDA AGRICULTURAL SUPPLY COMPANY, Jacksonville Florida U.S.A.

800-L

comments

UNDER THE FEDERAL INSPECTION
PROGRAM
FOR FOOD AND DRUGS
EST. UNDER NO. 2342-9

Cameraman's Note
Poor Copy



SUSPERSE NU-TRI-HI-COP


Product No. 20774G11

ACTIVE INGREDIENTS:	100.00%
INERT INGREDIENTS:	0.00%
TOTAL —	100.00%

CAUTION: HARMFUL IF SWALLOWED

KEEP OUT OF REACH OF CHILDREN

Use only as directed on label and for the purpose intended.



EPA Reg. No. 2342-FL-1

NOTE: The above poundages are suggested on an approximate basis, and should be applied in enough water to insure thorough distribution over the area.

Do not mix this spray with other commonly used pesticides such as DDT, Dieldrin, Chlorobenzilate, Kelthane, Zineb, Parathion and Lead Arsenate.

TOMATOES, PEPPERS, POTATOES — Against Early and Late Blight of Potatoes and Late Blight of Potatoes, Sprota Leaf Spot of Tomatoes and Late Blight of Potatoes. Use 3 1/2 to 4 pounds per 100 gallons of spray every 2 to 7 days as needed.

CUCURBITS (Cucumbers, Watermelons, Cantaloupes, Squashes) — Against Downy Mildew. Use 3 1/2 to 4 pounds per 100 gallons of spray every 5 to 10 days as needed.

CELERY — Against Early Blight. Use 4 pounds per 100 gallons of spray every 5 to 10 days as needed.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy by burning, incineration, crushing and filling.

CAUTION TO USER

Do not use this product unless the use of this product is specifically recommended on the label. Do not use this product on any crop or plant which has not been specifically mentioned on the label. The user of this product will be liable for any damage to crops or plants which may result from the use of this product.

EPA Reg. No.

7-73 W

Manufactured by **FARR-MORRIS CHEMICAL CORP.** Jacksonville, Florida, U.S.A.

BEST DOCUMENT AVAILABLE