

(TOXAPHENE - SULPHUR)

FASCO 20-40 Cotton Dust is an insecticidal composition containing Toxaphene and Sulphur, specifically designed for use on cotton. It may be applied by ground equipment or by plane.

### DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, execessive residues, unsatisfactory control or other unintended consequences.

Against Boll Weevil, Cotton Bollworm, Leaf- 🥋 worm, Fleahopper and to suppress Spider Mites use for 10 to 15 pounds of dust per acre. Start dusting at the rate of 10 to 12 pounds per acre and increase rate higher later in the season, depending upon plant growth and degree of infestation. Against Grasshoppers, apply 20 pounds per acre. Do not apply over 20 lbs. per acre on Cotton.

NOTE-The U.S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

### CAUTION

Harmful If Swallowed! Avoid breathing of dust or spray mist. Avoid contact with eyes, skin

> Uroft E 71 6955

CAUTION: Keep out of the reach of children. See additional cautions on this label. ACCEPTED 1111 343.26 SECTICIDE TODE AJD SIER.

ACTIVE INGREDIENTS:

Toxaphene (Chlorinated Camphene) (Chlorine Content 67% to 69%) Sulphur as S INERT INGREDIENTS

R Frade Markes Reg. U. S. Patent Offfice

KERR-MCGEE CHEMICAL CORP. Jacksonville, Florida, USA Manufactured by

No. 3-0522-1

# **CAUTION** (Cont'd)

and clothing. Wash thoroughly after using. Do not contaminate feed or foodstuffs. Do not feed treated Cotton trash to dairy animals or animals being finished for slaughter, nor graze such animals on Cotton receiving late applications of this dust.

## DANGER TO FISH AND WILDLIFE


This product is poisonous to fish and wildlife. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of this product from areas treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke), Birds feeding on treated areas may be killed.

## CONDITIONS OF SALE

All statements concerning the use of this prod-40.0% uct apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or Total ---- 100.00% its use, which extend beyond the description on the label. Read all directions carefully.

USDA Reg No 2042 20



By Wt. 20.0%

40.0%