



Net Contents
5 Pounds

CHLORDANE 40-WP

Product
No. 623

ACTIVE INGREDIENT:

Technical Chlordane
(Equivalent to 24% Octachloro-4
7-methanotetrahydroindane and 16% of related compounds)

By Wt.
40%

INERT INGREDIENTS:

60%
TOTAL 100%

USDA Reg. No. 2342-55

CAUTION: Keep out of the Reach of Children.
See Below for Additional Cautions.

FASCO CHLORDANE 40-WP is a dry wettable insecticidal compound containing 40% Chlordane by weight. It is a contact and stomach poison with residual effect. This product is designed for use with water in sprays.

DIRECTIONS

NOTE: For small amounts of spray use one heaping tablespoonful per gallon of water. Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

BEANS, CELERY, CABBAGE, ONIONS, PEPPERS — Against Serpentine Leaf Miner, Thrips, Cabbage Worms, Stink Bugs, Cutworms (Spray soil surface), Armyworms, Flea Beetles, use 2½ to 5 pounds per acre in sufficient water for good coverage. Repeat every 7 to 10 days or as needed. Do not use on Green or Snap Beans after pods begin to form; on Celery after bunches begin to form or plants are half grown; or on Cabbage or Pepper after edible parts start to form. Do not use on Spring or Green Onions.

NOTE — The U.S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should NOT be marketed and to do so may violate State law and will violate Federal law when sold in Interstate Commerce. Such crops should not be marketed without removal of such excess residues.

SOIL BORNE INSECTS — Against Wireworms, White Grubs, Mole Crickets and Southern Corn Rootworm Larvae that attack Corn. Before planting use 5 Pounds per acre on light sandy soils or 10 Lbs. per acre on muck soils as a broadcast spray in sufficient water for coverage and work into the top 3 or 4 inches of soil immediately.

LAWNS and GREENS — Against Grasshoppers, Mole Crickets, Armyworms, Sod Webworms, Lygus Bugs, Chiggers, Ants, Crickets and certain other pests, use 5 oz. per 1,000 Sq. ft. in sufficient water for coverage.

ORNAMENTAL SHRUBS and FLOWERS — Against Thrips, Serpentine Leaf Miner, Flea Beetles, certain other beetles, Armyworms and Cutworms use 2½ to 3 lbs. per 100 Gallons water (1 Meas. Teaspoon per 1 Gallon) Repeat as needed.

CAUTION

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid contamination of feed and foodstuffs. Avoid breathing spray mist.

Any residue at harvest should be removed by washing, wiping or other effective means.

Do not use on forage crops to be fed to or grazed by dairy animals or animals being finished for slaughter. Do not use on dairy cattle or in dairy barns. Do not use on chlordane sensitive plants such as cucurbits.

Toxic to fish and wildlife. Do not use where any body of water will be contaminated by direct application or run-off.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

DRAFT H 71 6218

ACCEPTED
MAY 26 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTRATION
ED UNDER NO. 2342-55 SUBJECT
TO ATTACHED COMMENTS.

Manufactured by **FLORIDA AGRICULTURAL SUPPLY COMPANY**, Jacksonville, Florida, U.S.A.