



NET CONTENTS
50 POUNDS

ZINEB DUST-6 1/2

PRODUCT
NO. 3-602-1

FASCO Zineb Dust-6 1/2 contains 6.5% Zineb (Zinc Ethylene Bis-Dithiocarbamate) fungicide for use against certain plant diseases of economic plants.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

CUCUMBERS, WATERMELONS, CANTALOUPE, SQUASH — Against Anthracnose, Downy Mildew, Gummy Stem Blight and certain other diseases, use 25 to not over 45 Pounds per acre, beginning when runners begin to form or when first signs of disease appear, repeating every 4 to 10 days as required to keep new growth covered. Do not use on summer squash within 7 days of harvest.


TOMATOES AND PEPPERS — Against Early Blight and Late Blights, Grey Leafspot, and against Freeze Spot of Pepper, use 25 to 40 Pounds (not over 45 Pounds per acre for Tomatoes) every 3 to 7 days, beginning with first indications of diseases.

PEPPERS FREEZE SPOT
CABBAGE, CAULIFLOWER, BROCCOLI, COLLARDS, TURNIPS — Against Alternaria Leafspot, Downy Mildew use at rate of 25 to 40 Pounds per acre (not over 45 Pounds for Collards and Turnips), 2 or 3 times each week beginning when plants come up in seed beds and every 5 days in the field. Do not apply within 7 days of harvest.

CELERY — Against Early Blight use 25 to not over 45 Pounds per acre every 4 to 7 days. Dust plant beds every 4 to 7 days, beginning when plants come up. In field start when plants are established and dust every 5 days. Remove residues at harvest by stripping, trimming and washing.

DRAFT H 886097

CAUTION: Keep out of the reach of children.
See side panel for additional cautions.



| | |
|--|-----------------------|
| ACTIVE INGREDIENTS: | BY WT. |
| Zineb (Zinc Ethylene Bis-Dithiocarbamate) (Equivalent to Zinc, expressed as metallic 1.54%) | 6.5% |
| INERT INGREDIENTS: | 93.5% |
| | TOTAL — 100.0% |

® Trade Marks Reg. U. S. Patent Office

TOBACCO PLANT BEDS—Against Blue Mold, use 2 to 3 1/2 Pounds per 100 sq. yards. Begin dusting when leaves are the size of a dime and before Blue Mold is first reported in the area. Dust every 3 days (every 2 days after Blue Mold appears). Change to or alternate with FASCO P-D-Q Plant Bed Dust where insects appear.

NOTE — The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State Law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

CAUTION

May cause irritation of nose and throat. Do not breathe dust.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

USDA Reg. No. 2342-67

Manufactured by **FLORIDA AGRICULTURAL SUPPLY COMPANY**, Jacksonville, Florida U.S.A.



FASCO

NET CONTENTS
50 POUNDS

ZINEB DUST-6 1/2

PRODUCT
NO. 3-602-1

FASCO Zineb Dust-6 1/2 contains 6.5% Zineb (Zinc Ethylene Bis-Dithiocarbamate) fungicide for use against certain plant diseases of economic plants.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

CUCUMBERS, WATERMELONS, CANTALOUPE, SQUASH — Against Anthracnose, Downy Mildew, Gummy Stem Blight and certain other diseases, use 25 to not over 45 Pounds per acre, beginning when runners begin to form or when first signs of disease appear, repeating every 4 to 10 days as required to keep new growth covered. Do not use on summer squash within 7 days of harvest.


TOMATOES AND PEPPERS — Against Early ^{blight} and Late Blights, Grey Leafspot, and against Frogeye Spot of Pepper, use 25 to 40 Pounds (not over 45 Pounds per acre for Tomatoes) every 3 to 7 days, beginning with first indications of diseases.

PEPPERS ^{FROGEYE SPOT}
CABBAGE, CAULIFLOWER, BROCCOLI, COLLARDS, TURNIPS — Against Alternaria Leafspot, Downy Mildew use at rate of 25 to 40 Pounds per acre (not over 45 Pounds for Collards and Turnips), 2 or 3 times each week beginning when plants come up in seed beds and every 5 days in the field. Do not apply within 7 days of harvest.

CELERY — Against Early Blight use 25 to not over 45 Pounds per acre every 4 to 7 days. Dust plant beds every 4 to 7 days, beginning when plants come up. In field start when plants are established and dust every 5 days. Remove residues at harvest by stripping, trimming and washing.

DRAFT H 886097

CAUTION: Keep out of the reach of children.
See side panel for additional cautions.



| ACTIVE INGREDIENTS: | BY WT. |
|--|---------------|
| Zineb (Zinc Ethylene Bis-Dithiocarbamate) (Equivalent to Zinc, expressed as metallic 1.54%) | 6.5% |
| INERT INGREDIENTS: | 93.5% |
| TOTAL | 100.0% |

® Trade Marks Reg. U. S. Patent Office

TOBACCO PLANT BEDS—Against Blue Mold, use 2 to 3 1/2 Pounds per 100 sq. yards. Begin dusting when leaves are the size of a dime and before Blue Mold is first reported in the area. Dust every 3 days (every 2 days after Blue Mold appears). Change to or alternate with FASCO P-D-Q Plant Bed Dust where insects appear.

NOTE — The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should **not** be marketed and to do so may violate State Law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

CAUTION

May cause irritation of nose and throat. Do not breathe dust.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

USDA Reg. No. 2342-67

FLORIDA AGRICULTURAL SUPPLY COMPANY, INC. • TAMPA, FLORIDA U.S.A.