



NET CONTENTS
5 POUNDS

LINDANE 25-WP

PRODUCT
NO. 776

ACTIVE INGREDIENTS	BY WT.
Gamma Isomer of Benzene Hexachloride (from Lindane)	25.0%
INERT INGREDIENTS	75.0%
TOTAL — 100.0%	

® Trade Marks Reg. U.S. Patent Office
USDA Reg. No. 2342-252

FASCO Lindane 25-WP is a wettable powder insecticidal spray concentrate containing technically pure Gamma Isomer of Benzene Hexachloride, from Lindane.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

MIXING RATES — Unless otherwise directed below use 1 to 2 lbs. per 100 gals., beginning when insects appear, and apply at least 100 gals. of spray per acre on large plants. For SMALL AMOUNTS OF SPRAY use 1 heaping tablespoonful per 3 gals. water.

CUCUMBERS, CANTALOUPES, MELONS, SQUASH — Aphids, Pickleworm, Melonworm, Thrips, Cucumber Beetles, Squash Bug and Grasshoppers begin spraying when runners begin to develop and continue on a 5 to 10 day schedule. For Squash Bug increase to 2 pounds. Lindane may adversely affect the flavor of melons and squash. Do not use more than two lbs. per acre on cucurbits. Do not apply to cucumbers or squash within one day of harvest.

EGGPLANTS — Aphids and Flea Beetles, use as needed, not over 2 lbs. per acre. Do not apply after fruit starts to form.

SPINACH — Aphids, Flea Beetles and Leaf Miner, not after seedling stage, not over 1.6 lbs. per acre.

ONION — Thrips and Onion Maggot, not over 4 lbs. per acre. Do not use on green or spring onions (Shallots).

WIREWORM SEED TREATMENT — Apply as dust to seed at planting time — following amounts per 100 lbs. of seed; Corn — 8 oz.; Beans, Peas, Oats, Rye, Barley, Wheat, Milo — 4 oz. Do not use treated seed as food or feed.

NOTE — The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

ORNAMENTAL PLANTS — Aphids, Thrips, certain Caterpillars, Plant Bugs, Flea Beetles, certain Beetles, Leaf Rollers.

TICKS, CHIGGERS (RED-BUGS) — Spray vegetation at 2 lbs. per 100 gals. when needed. Do not contaminate crops used for food or forage.

GLADIOLUS BULBS — Against Wireworms dust bulbs at planting time with 2 to 3 oz. per 100 lbs. bulbs.

WARNING: Keep Out of the Reach of Children

May be Fatal If Swallowed. May be absorbed through skin! Do not breathe dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

DANGER TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not use where runoff is a frequent problem. Birds feeding on treated areas may be killed.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read directions carefully.

DRAFT M

71-6221

May 3, 1971
3342 154

Manufactured by FLORIDA AGRICULTURAL SUPPLY COMPANY, Jacksonville, Florida, U.S.A.