

FASCO

Net Contents  
50 Pounds

Product  
No. 3-0792-1

MALATHION 25-WP

ACTIVE INGREDIENTS	BY WT.
Malathion* -----	25.00%
INERT INGREDIENTS -----	75.00%
TOTAL -----	100.00%

\*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate

Made from CYTHION ® Insecticide  
The Premium Grade Malathion  
® Reg. Trademark American Cyanamid Company

® Trade Marks Reg. U.S. Patent Office

USDA Reg. No. 2342-258

CAUTION: Keep out of the reach of Children  
See side panel for additional cautions.

FASCO MALATHION 25-WP is a wettable concentrate insecticide and miticide for use in water suspension sprays. This product contains Premium Grade Malathion.

#### DIRECTIONS


The contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

ORNAMENTAL FLOWERS AND SHRUBS - Against Aphids, Spider Mites, Mealybugs and Whitefly-use 2½ Lbs. per 100 Gals. or 2 level tablespoons per 1 Gal. water. Against Soft Brown Scale, Florida Red Scale and Black Scale Crawlers, use 6 lbs. per 100 gals. or 1 oz.

(4 level tablespoons) per each gallon of water. Spray for thorough, full coverage and repeat as needed. Injury may occur on Boston, Maidenhair and Pteris Ferns, and some species of Crassula.


#### VEGETABLES

Against Aphids or Spider Mites attacking certain vegetables use 2 lbs. per 100 Gals. of spray for thorough coverage when needed but not to exceed the pounds of this product per acre given below nor nearer to harvest than given below.



VEGETABLES (CONT'D)

<u>Crop</u>	<u>Max. lbs./Acre</u>	<u>Days Between Last Application and Harvest</u>
Pepper	5 lbs.	3
Okra	5 lbs.	Not after pods start to form
Endive (Escarole)	2 lbs.	7
Eggplant	2 lbs.	3
Beans	5 lbs.	1
Cucumber	6 lbs.	1
Lettuce (Leaf)	5 lbs.	7
Carrots,	2 lbs.	7
radish	2 lbs.	7
Tomato	2 lbs.	1
Strawberry	2½ lbs.	3
Cabbage,	2 lbs.	7
Cauliflower,	2 lbs.	7
Collards,	2 lbs.	7
Mustard	2 lbs.	7
Broccoli,	2 lbs.	3
Turnips	2 lbs.	3



Do not apply less than 2 pounds of this product per acre on any of the above vegetable crops.

CITRUS

Oranges, Grapefruit, Tangelos, Limes, Lemons, Kumquats and Tangerines) - Against Florida Red Scale and Purple Scale use 3 lbs. per 100 gals. for light infestations to 5 or 6 lbs. for medium infestations. Against Aphids, use 2 lbs. per 100 gals. Do not use when trees are in bloom nor later than 7 days before harvest. Do not exceed 40 lbs. per acre. PEACHES - Against Plum Curculio and Oriental Fruit Moth use 3 lbs. per 100 Gals. of water. Make 4 or more cover applications beginning when petals fall, not within 7 days of harvest. Do not exceed 28 lbs. per acre.

PEANUTS - For the protection of stored peanuts (unshelled) against infestations of Red Flour Beetle, Indian Meal Moth, Confused Flour Beetle, Rice Weevil, Flat Grain Beetle, Rusty Grain Beetle, Lesser Grain Borer, Granary Weevil and Saw-Toothed Grain Beetle, apply FASCO Malathion 25-WP as follows:

Bulk Spray Treatment - Peanuts Going into Storage: Apply 2½ pints of 57% Premium Grade EC in 5 gallons of water per 15 tons of farmers' stock peanuts as they go into storage. Apply coarse spray uniformly. Preferable use a suitable mechanical spray applicator which regulates rate of application to flow of peanuts. Otherwise, adjust pressure and size of nozzle opening to rate of flow of peanuts.

Residual Warehouse Spray - Before Storing Peanuts: Clean warehouse thoroughly of trash and old remains of peanuts 1 to 2 weeks before new peanut crop is stored. Thoroughly spray with sufficient pressure interior of empty warehouse (including cracks and protected places) outside walls to height of 6 to 8 feet, and the ground to a distance of about 6 feet from warehouse by diluting 1 pint of 57% premium grade EC in sufficient water to make 2½ gallons of spray or 1 gallon with 19 gallons of water. Apply finished spray at the rate of 2 gallons per 1000 Square Feet or to run-off.

SUPPLEMENTAL SURFACE SPRAY: After warehouse has been filled, level surface of peanuts before spraying. Apply surface spray using 1-3/4 lbs. of premium grade malathion 25% WP in 2 gallons of water and apply at the rate of 2 gallons per 1000 square feet surface. Apply first treatment as soon as bin is filled and leveled, but not later than the first week in October. Apply the second surface treatment 1 month later, followed by treatments at 2 month intervals. Use a piston type power sprayer equipped with agitator and nozzle capable of delivering a coarse spray. Repeat surface application 60 days later if warm temperatures prevail. Otherwise delay further treatment until late winter.

#### CAUTION

Harmful if swallowed. Avoid prolonged breathing of spray mist or dust. Avoid repeated or prolonged contact with skin. Wash thoroughly. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.

#### DANGER TO FISH

This product is poisonous to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of this product from areas treated.

#### CONTAINER DISPOSAL

Reuse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke and fumes.)

#### NOTE

The U.S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

#### CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

DRAFT K

Manufactured by FLORIDA AGRICULTURAL SUPPLY COMPANY  
Jacksonville, Florida, U.S.A.

ACCEPTED

Jan-6-1970

2372-

28