

ACCEPTED WITH COMMENTS

ACCEPTED



JACKSONVILLE, FLORIDA, U.S.A.

(R) TRADE MARKS REG. U. S. PAT. OFFICE

* *

IDEAL Brand OILAN — is an emulsifiable concentrate insecticide containing a special highly refined insecticidal oil fortified with malathion. This product should be diluted in water for use as a spray against insects of certain ornamental and fruit plants as directed below.

DIRECTIONS

Read the entire label and follow directions and cautions. Use contrary to these directions may result in injury to valuable plants, excessive residues, or in poor insect control.

hATES — Unless directed otherwise below mix 1.28 Fl. Oz. $(2\frac{1}{2}$ Measuring Tablespoons) per each gallon of water; 4 Fl. Oz. $(\frac{1}{4}$ Pt.) per 3 Gals.; 8 Fl. Oz. per 6 Gals.; $1\frac{1}{2}$ Pts. per 18 Gals.; or 2 Qts. per 50 Gals. Spray for even and thorough coverage of upper and lower leaf surfaces, stems and fruit.

CAMELLIAS: Use against Tea Scale and other Scale Insects, Spider Mites, Aphids, Mealybug and White Fly. Against heavy Tea Scale infestations increase strengths (above) by one-third. Use 1.7 Fl. Oz. (3½ Measuring Tablespoons) per each gallon of water or 10 Fl. Oz. per 6 Gals. water. Spray in Spring and again in late September if needed.

AZALEAS: Against Spider Mites, Mealybugs, Lace bug, Scale Insects, and Leaf Miner (roller), spray just after blooming, again in late May or early June; and in late September. Use at other times, if needed, but within limits of the "Cautions" given below.

GARDENIA, HOLLY, IXORA, LIGUSTRUM, and JUNIPER — Against Scale Insects, Mealybugs, White Fly, Spider Mites and Aphids. To avoid plant injury observe "Cautions" given below.

CITRUS, AVOCADOS — Use against Scale Insects, Spider Mites, Mealybugs, White Fly and Aphids. To avoid plant injury follow "Cautions" given below. Do not use within 7-days before harvesting Citrus or Avocados. Do not apply during full bloom on Citrus.

CAUTIONS

Harmful if swallowed. Avoid breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep out of reach of children.

This product corrains petroleum oils and should never be combined with Sulphur or Lime-Sulphur sprays. Do not use within 30 days before or after Sulphur or Lime-Sulphur applications. Do not repeat oil sprays in less than thirty days. Do not spray plants with this product if the plants are suffering from lack of moisture, wilted or near wilted, or in otherwise weakened condition. Do not spray with this product when temperatures exceed 90 F. or when temperatures are expected to approach freezing. Oil sprays after October 1 may increase susceptibility to cold injury. Oil sprays should not be used on bearing citrus during August or September and until harvest as fruit maturity may be delayed. Application on new tender growth is more likely to produce injury than after foliage is hardened.

Do not use on Maiden Hair Ferns, Cocos Palms, Maples, or on non-woody plants as injury may occur.

Do not use or store near heat or open flame.

Do not re-use container. Destroy when empty.

DANGER TO FISH

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully

DRAFT D

USDA Reg. No. 2342-465

71-6228

FORM 1290-