

NET CONTENTS  
50 POUNDS

# SEVIN\* DUST-10

PRODUCT  
NO. 927

FASCO Sevin\* Dust-10 is an insecticidal dust designed for use against certain insects attacking field and vegetable crops as described below.

### DIRECTIONS

Pesticide products used contrary to directions may cause serious plant or personal injury.

**SWEET CORN** — Against Corn Earworms, use at rate of 18 to 20 lbs. per acre (no more). Begin dusting when first silks appear and repeat at 2 to 3 day intervals until silks are withered. May be applied up to harvest, at rates directed, without exceeding residue tolerances. Application of this product to tassel region of corn during the pollen shed period will seriously reduce bee populations.


**POTATO, TOMATO, EGGPLANT, PEPPER** — Against Colorado Potato Beetle, Flea Beetles and Leaf hoppers apply 8 to 10 lbs. per acre when insects appear. Repeat at 7 to 10 day intervals. Against Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs and Stink Bugs apply 10 to 20 lbs. (no more) per acre when insects appear. Repeat at 7 to 10 day intervals.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, and KOHLRABI** — Against Corn earworm, Armyworm, Cabbage Caterpillar, Flea Beetle, and Harlequin Bug, apply 10 to 20 lbs. (no more) per acre when insects appear. Repeat at 7 to 10 day intervals. Do not apply later than 3 days before harvest.

**PASTURE, FORAGE GRASSES, CLOVER, ALFALFA** — Against Thrips, Armyworms, Grasshoppers, and Stink bugs, apply 10 to 15 lbs. (no more) per acre when insects appear. Repeat 7 to 14 days later if necessary. May be applied up to harvest at rates directed, without exceeding residue tolerances.

**NOTE** — The U.S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

**COTTON** — Against early season pests such as Thrips, FLEA BEETLES, and COTTON FLEA HOPPER, use 5 to 10



(not over 25) lbs. per acre every 5 to 10 days as needed beginning when insects increase to injurious numbers.

Against mid-season and late-season BOLL WEEVIL, COTTON BOLLWORM, FALL ARMY WORM, COTTON LEAF PERFORATOR, LEAF HOPPERS, LYGUS BUGS, and LEAF ROLLERS, use 10 to 20 (not over 25) lbs. per acre, the amount depending on plant size and severity of infestation. Applications for BOLL WEEVIL should be made every 5 to 7 days as long as control is necessary.

Against PINK BOLLWORM, use 15 to 25 (not over 25) lbs. per acre, the amount depending on size of cotton and degree of infestation. Apply every 5 to 7 days as needed.

Against GRASSHOPPERS, use 5 to 15 (not over 25) lbs. per acre, applied on appearance of insects and their feeding damage. Highest dosage should be used against mature grasshoppers.

APHIDS are usually suppressed by a regular Sevin schedule. SEVIN may be applied after the bolls open. Allow 7 days before grazing or harvesting crop residue for dairy or meat animals.

### CAUTION

HARMFUL IF INHALED OR SWALLOWED. Do NOT take internally. Avoid prolonged or Repeated Breathing of Dust. Skin Contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to Clean Clothing Daily.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**NOTE FOR PHYSICIANS:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**  
Do not reuse container. Destroy when empty.

### CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

### ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methyl-  
carbamate)

### BY WT.

10.00%

### INERT INGREDIENTS:

90.00%

**Total — 100.00%**

\*Trade Marks Reg. U. S. Patent Office U. S. Pat. No. 2,903,476

\*Sevin, Trade Mark of UCC for 1-naphthyl N-methyl carbamate

**CAUTION: Keep out of the reach of children.**

See side panel for additional cautions.