

ACCEPTED
 Jan 1, 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 2342-481

Comments



NET CONTENTS 3- POUNDS PRODUCT No. 946

THIODAN® 50-WP

ACTIVE INGREDIENTS	BY WT.
*Endosulfan	50.00%
INERT INGREDIENTS	50.00%
TOTAL	100.00%

*Hexachlorohexahydro-2,4,3-benzodioxathiepin oxide
 Thiodan is a registered trademark of Farbwerke Hoechst A.G.

WARNING: Keep Out of the Reach of Children. See below for additional cautions.

DIRECTIONS
 FASCO THIODAN 50-WP is a wettable powder insecticide concentrate for use in water base sprays against insects as directed below.

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

IRISH POTATOES — Against Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper, and Aphids, use 1 to 2 lbs. per 100 Gals., not over 5 lbs. of concentrate per acre. Apply when insects appear and repeat as needed up to harvest.

EGGPLANTS, TOMATOES, PEPPERS — Against Aphids, Colorado Potato Beetle, Green Stink Bug, Blister Beetle, Flea Beetle, and Whitefly on Eggplants and Tomatoes; Aphids and Flea Beetles on Peppers, use 1 pound (no more) per acre in sufficient water for good coverage. Apply when insects first appear and repeat as needed. Spray plants thoroughly. Do not apply within 1 day of harvest.

TOBACCO INSECTS — Against Flea Beetle, Hornworm, Budworm, Green Peach Aphid, and Cabbage Looper.

TOBACCO SEEDBEDS — Use 1 pound per 100 gallons of water. Apply about 6 gallons of spray per 100 square yards.

FIELD — Use 1 to 2 pounds per acre in sufficient water for coverage. Apply when insects first appear, repeat as needed. Do not apply within 5 days of harvest.

BROCCOLI, CABBAGE — Against Cabbage Aphids, Cabbage Looper, Imported Cabbage-worm, and Diamond back moth larvae, use 1 1/2 to 2 pounds (no more) per acre in sufficient water for good coverage. Apply when insects first appear, repeat as needed. Do not apply within 7 days of harvest.

CUCUMBERS, MELONS, SUMMER SQUASH, WINTER SQUASH — Against Aphids, Cucumber beetles, and Squash vine borer, use 1 to 2 pounds (no more) per acre in sufficient water for good coverage. For vine borer apply weekly to flower buds, stems and vines beginning when moths first appear. Apply against Aphids and Cucumber beetles when they first appear, repeat as needed. Against Cabbage Looper, use 2 pounds (no more) per acre. Apply when Looper's first appear, repeat as needed. May be applied up to day of harvest.

PEACHES — (Southern States) — Against Peach Tree Borer use 1 1/2 lbs. per 100 gallons of water. Spray on bark areas from ground to scaffold limbs immediately after harvest and repeat 3-4 weeks later. Do not allow monkeys to graze in treated orchards.

NOTE — The U.S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerance, should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

WARNING

Hazardous if swallowed, inhaled or absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eye contact, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. During normal use or prolonged exposure in spray mixing and loading operations, wear clean clothes, rubber gloves and a mask or respirator of a type passed by the U.S. Department of Agriculture for Thiodan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas on the day of application should wear protective clothing, test for use, storage in or around the home. Do not reuse the container. Do not use when empty.

DAZGE TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not use where runoff is likely to occur. Apply this product only as specified on this label. Do not contaminate water by washing of equipment or disposal of waste.

STATE OF USE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranty, expressed or implied, concerning this product or its use, which extends beyond the description on the label. Read all directions carefully.

DRAFT 1 64-111-101-101-101 U.S. Patent No. 2,772,691

ACCEPTED WITH COMMENTS