

IDEAL

NEMAGON LIQUID is a liquid concentrate which is readily diluted with water for use as a soil treatment against certain Nematodes. Use on Mineral Soils only to control Cyst Nematodes (soybean, golden, sugar beet); free living nematodes (sting, ring, awl, spiral, lance, pin, sheath, stubby root, stilet, dagger); root knot, root-lesion or meadow, burrowing, citrus, and stem and bulb nematodes.

DO NOT USE NEMAGON LIQUID ON SENSITIVE CROPS INCLUDING BEETS, CARNATION, CHRYSANTHEMUM, DWARF PALMS, GARLIC, IRISH POTATO, LANTANA SPP., ONION, PEPPER, SWEET POTATO, AND TOBACCO.

GENERAL DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

Apply NEMAGON LIQUID when soil temperature at a depth of 6 to 8 inches is 40 to 80 F. The soil should be in good tilth and contain adequate moisture for good seed germination. Trash in the soil should be well decomposed. Extremely wet or dry soils should not be treated. Applications can be made before, during, or after planting according to directions only.

HOW TO APPLY:

Pre-planting Overall Application — Use a garden hose sprayer or sprinkler can. Dilute the NEMAGON LIQUID with sufficient water to cover surface evenly (at least 1 gallon of coarse spray to each ten square feet). Follow immediately with irrigation (sprinkling) to soak the fumigant solution to a depth of six inches. Do not disturb the soil after application for 7-14 days for shallow-rooted crops and 4 to 6 weeks for deep rooted crops.

Pre-planting Row Application — Mark out the rows and prepare a narrow trench 6 inches deep directly in the rows to be planted. Apply the NEMAGON LIQUID, diluted in water for coverage, evenly down the center of each trench. Immediately refill trenches with soil and lightly pack by rolling. Do not disturb the soil after application for 7-14 days for shallow-rooted crops and 4 to 6 weeks for deep rooted crops.

Post-planting Row Application to Established Plants — Prepare a narrow trench 6 inches deep and 6 inches wide on each side of the plants growing in the row. Apply NEMAGON LIQUID, diluted in water for coverage, evenly in the bottom of the trench. Immediately cover the trench with 6 inches of soil and lightly pack the soil.

CROPS AND APPLICATION RATES

ORNAMENTALS

Use 4 tablespoons per 100 sq. ft. (pre-planting Overall Application) or 4 tablespoons per 100 ft. of row (Pre-planting or Post-planting Row Application) for Aster, Azalea, Camellia, Daisy (Shasta), Forsythia, Gardenia, Hibiscus, Ligustrum (Privet),

Philodendron, Pittosporum, and Rose. Use 2-2/3 tablespoons per 100 sq. ft. (Pre-planting Overall Application) or 2 tablespoons per 100 ft. of row (Pre-planting or Post-planting Row Application) for African Violet, Caladium, Dieffenbachia, Dogwood, Flowering Cherry, Gladiolus, Hydrangea, and Spirea.
LAWN GRASSES (Bermuda, Bluegrass, Centipede, St. Augustine, Zoysia). Apply as a drench (Pre-planting Overall Application or to established lawns) using 3 to 7 tablespoons per 100 sq. ft., 1/2 to 1 cup per 250 sq. ft., 1/2 to 1 pint per 500 sq. ft. Aerate before application for better penetration.

WARNING! HARMFUL IF INHALED. MAY BURN SKIN.

CAUSES IRRITATION OF EYES, NOSE, AND THROAT. MAY BE ABSORBED THROUGH SKIN.

Do not breathe vapor. Use in well ventilated area. Do not get in eyes. Avoid contact with skin or clothing. Do not take internally. Wash thoroughly with soap and water after handling and before eating or smoking. In case of accidental spillage on person or clothing, immediately remove contaminated shoes and clothing and wash skin with soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Wash clothing and air shoes thoroughly before re-use. Do not contaminate feed and foodstuffs. Keep out of the reach of children. Keep children and pets off treated areas until this material has been washed into the soil. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material. Keep away from heat and open flame.

NON-WARRANTY

Seller makes no warranty of any kind, express or implied concerning this product. Buyer assumes all risk of use or handling, whether in accordance with the directions or not.

Manufactured by

FLORIDA AGRICULTURAL SUPPLY COMPANY

DIVISION — WILSON & TOOMER FERTILIZER COMPANY

JACKSONVILLE, FLORIDA, U.S.A.

DRAFT F-1

875541

U.S.D.A. Reg. No. 2342-612

NEMAGON LIQUID

AGAINST NEMATODES FOR GRASSES, SHRUBS, FLOWERS SOIL DRENCH

ACTIVE INGREDIENTS	BY WEIGHT
1,2-Dibromo-3-Chloropropane	67.87% **
Other Halogenated C ₃ Compounds	3.7% **
INERT INGREDIENTS	28.56%

TOTAL — 100.00%

Licensed under United States Patents 3,049,472 and 2,937,936
Contains 2 Quarts (8.6 pounds) NEMAGON SOIL FUMIGANT per gallon.
*Shell Registered Trademark.
**Equivalent to 71.44% w Nemagon Soil Fumigant.

WARNING: Keep Out of the Reach of Children. See back panel for additional cautions.

NET CONTENTS
1 PINT

PRODUCT NO. 142

MINI LABEL

ACCEPTED
Mar 31 1970
REGISTERED SUBJECT

2342-612