

8-26-69
2342-702

FASCO

SPECIMEN LABEL

NET CONTENTS
4 POUNDS

DIPHENAMID 50-WP

PRODUCT NO.
3-2556-1

A selective herbicide for the pre-emergence control of weed grasses and broadleaf weeds in tobacco, tomato and pepper.

ACTIVE INGREDIENTS	BY WT.
Diphenamid (N,N-Dimethyl-2, 2-diphenylacetamide)	50.00%
INERT INGREDIENTS	50.00%
TOTAL — 100.00%	

CAUTION: Keep Out of the Reach of Children

Harmful if swallowed. Avoid breathing of dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and food-stuffs.

WEEDS CONTROLLED			
ANNUAL GRASSES	ANNUAL GRASSES	ANNUAL BROADLEAF WEEDS	ANNUAL BROADLEAF WEEDS
Annual bluegrass	Fall panicum (Panic grass)	Carpetweed	Lambquarters
Annual sedge	Foxtail	Common chickweed	Moose ear chickweed
Barnyardgrass (Watergrass)	Goosegrass (Crowfoot grass)	Corn spurry	Sheep's pursue
Cheatgrass	Stinkgrass	Eying primrose	Spiny amaranth (Stickerweed)
Crabgrass	Wild oat	Field peppergrass	Thyme leaved sandwort
Johnsongrass (fruit seed)	Witchgrass (Ticklegrass)	Florida pusley	Pigweed
Ryegrass		German moss (Knotgrass)	Purslane
Sandbur (Sandspur)		Groundsel	Red sorrel
		Knotweed	Smartweed

DIRECTIONS

SOIL PREPARATION: Prepare the soil for seeding, planting, or transplanting according to good agricultural practices. All weed growth should be worked into the soil prior to treatment. To insure uniform treatment, all soil clods and organic debris should be eliminated from the rows.

CULTIVATION: Shallow (1/2 to 2 inches) cultivation will not destroy the effectiveness of FASCO Diphenamid 50 WP.

APPLICATION: (See Recommendations) Use standard low pressure (25 to 50 ps) boom spray equipment. Calibrate before use and check frequently during application to insure a uniform spray pattern. Continuous agitation necessary for optimum results.

Broadcast (Over-all) Application: Uniformly apply the recommended rate of FASCO DIPHENAMID 50 WP in 30 to 50 gallons of water per acre. Avoid spray overlap that will increase FASCO DIPHENAMID 50 WP rates above the recommended amount. Use Tee Jet 8003 or 8004 fan type or similar nozzles of equal or greater size. Use 50 mesh or coarser screens in spray nozzle heads. For small areas, apply FASCO DIPHENAMID 50 WP in backpack or other low pressure sprayers at 3 to 5 gallons of water per 1,000 Sq. Ft.

BAND (ROW) APPLICATION: The amount of FASCO DIPHENAMID 50 WP required per crop acre is reduced with band application.

USE THIS FORMULA: Band width (inches) X Broadcast rate = pounds per acre
 Row width (inches) in pounds for band treatment

EXAMPLE: If 12 lb. per acre are recommended for broadcast, and you want to apply a 12-inch band on 36" rows —
 $12/36 \times 12 = 4$ lb. per acre

Use Tee Jet 8003 or 8004 or similar type nozzles. Use 50-mesh or coarser screens in spray nozzle heads.

INCORPORATION: In dry soils, incorporate immediately to depth of 1/2 to 2 inches.

IRRIGATION: Adequate surface soil moisture is essential. If available, apply 1/2 inch overhead irrigation shortly after application, and repeat in 7 to 10 days. FASCO DIPHENAMID 50 WP should be incorporated if furrow irrigation is used.

WARNING

Harmful if swallowed. Induce vomiting and call a physician immediately. Avoid prolonged or repeated contact with skin. If spilled on hands or other skin areas, wash immediately with soap and water. In case of contact with eyes, wash with water for 15 minutes and call a physician immediately. Do not use cover crops for food or cut for feed within 6 months after application. Avoid breathing dust. Avoid contamination of feed or foodstuffs. Do not reuse empty container. Destroy when empty.

GENERAL USE PRECAUTIONS

Read all label directions before using FASCO DIPHENAMID 50 WP. Use only on recommended crops at recommended rates. Do not use FASCO DIPHENAMID 50 WP in direct combination with fertilizers, insecticides or fungicides. Edible crops other than those registered should not be planted in FASCO DIPHENAMID 50 WP treated soils for 6 months after date of last application.

A variety of extreme environmental conditions may have effects on herbicide treatments. The grower should watch to note that the treatment is working and in case of poor control should take appropriate measures. Do not graze treated areas.

Package contains full net weight. Heat space is necessary to allow accurate filling and avoid spilling.

RECOMMENDATIONS: For control of certain annual grasses and broadleaf weeds in all crops listed below. Equivalent 1 1/2 tablespoons = 1 cupful.

Crop	When to Apply	Oz. per 1000 Sq. Ft.		Remarks
		Light Soil	Heavy Soil	
Tobacco (Seeding)	At seeding	3 (1 cupful)	4 1/2 (1 1/2 cupfuls)	Do not mix FASCO DIPHENAMID 50 WP with other chemicals before application.
Crop	When to Apply	Lb. per Acre		Remarks
		Light Soil	Heavy Soil	
Tobacco (Transplants)	At planting	8	12	Apply over top or directly to base of transplants.
Pepper	At direct seeding, transplanting, or with 1 month following transplanting	6	10	Before application, work all weeds growth into soil. May be applied over top of transplants.
Tomato	At direct seeding, transplanting, or with 1 month following transplanting	8	12	Before application, work all weeds growth into soil. May be applied over top of transplants.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

Manufactured by **FLORIDA AGRICULTURAL SUPPLY COMPANY**, Jacksonville, Florida, U.S.A.

DPAFT A

883227

U.S.D.A. Reg. No. 23427-07