

SPRAYING RECOMMENDATIONS

CROP	TO CONTROL	DIRECTIONS
Peanuts	Leaf Spot	Mix 2 to 3 Quarts TC-90 [®] in 10 to 25 gallons of water and apply to one acre. Begin treatment 2 weeks before disease normally appears for best preventive control, or make first application when disease is first noted. Apply at 14 day intervals after first application.
Cucumber, Cantaloupe, Squash, Pumpkin, Watermelon	Powdery Mildew	Mix 2 Quarts TC-90 in 100 gallons of water and apply to one acre. Begin treatment 2 weeks before disease normally appears for best preventative control, or make first application when disease is first noted. Apply at 7 to 10 day intervals after first application.
Cucumber	Downy Mildew (Southeastern States Only)	Same as for powdery mildew of cucurbits
	Angular Leaf Spot (North Central States Only)	
Citrus	Melanose (Florida Only)	Mix 6 to 8 quarts TC-90 in 10 gallons of water and apply to one acre by aircraft. Use 2 quarts TC-90 in 100 gallons of water when applied with ground spray equipment. Apply 1 to 3 weeks after petal fall.
Carrot	Leaf Spot (North Central States Only)	Mix 2 to 3 quarts TC-90 in 100 gallons of water and apply to one acre. Begin treatment 2 weeks

Net Contents
Gals.



Product No.
3-1125-1

TC-90[®]

EMULSIFIABLE LIQUID COPPER FUNGICIDE

ACTIVE INGREDIENTS

Copper salts of fatty and rosin acids
(Copper as Metallic 4.00%)

BY WT.

48.00%

INERT INGREDIENTS

52.00%

TOTAL — 100.00%

CAUTION: Keep Out of the Reach of Children. Harmful if swallowed. Do not induce vomiting. In case of ingestion call a physician immediately. Do not spray into or near fire or open flames.

USDA Reg. No. 2342 703

TC-90 Trade Mark of Tennessee Corporation

for

Pepper Bacterial Leaf Spot (Southeastern States Only) before disease normally appears for best preventative control or make first application when disease is first noted. Apply at 7 to 10 day intervals after first application.

Pour TC-90 into water with spray tank agitation to form best emulsion. After emulsion is formed, agitation is not required.

CONSULT STATE AGRICULTURAL EXPERIMENT STATION OR STATE EXTENSION SERVICE SPECIALISTS FOR ADDITIONAL INFORMATION AS TO TIMING, NUMBER AND RATE OF APPLICATIONS.

TC-90 IS COMPATIBLE WITH MANEB, CAPTAN, PHALTAN, DDT, TOXAPHENE, THIODAN, AND ORGANIC PHOSPHATES. USE WITH OTHER FUNGICIDES AND INSECTICIDES WITH CAUTION. OBSERVE ALL CAUTIONS AND LIMITATIONS ON LABELING OF ALL PRODUCTS USED IN MIXTURES.

WARNING: DECONTAMINATE AND DESTROY USED CONTAINERS

DECONTAMINATE empty containers by washing thoroughly with a strong solution of alkali and detergent and rinsing with water. Do not contaminate lakes, streams, or sewers with this water.

DESTROY containers by crushing or puncturing, then bury.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

12-9-67
2-3-72-703